


November 2018

E

WESTERN CENTRAL ATLANTIC FISHERY COMMISSION (WECAFC)
NINTH SESSION OF THE SCIENTIFIC ADVISORY GROUP (SAG)
Bridgetown, Barbados, 19-20 November 2018
Review of the state of fisheries and fisheries resources in the WECAFC region

Update on Review of the State of Fisheries in FAO Area 31¹

Introduction

1. This document is intended to update the state of fisheries in the region of the Western Central Atlantic Fisheries Commission (WECAFC), as presented in the VIII Session of the Scientific Advisory Group in 2017. The structure and content of the report remains essentially the same, except that FAO landings data have been updated to 2016 and the addition of stock assessment reports not included in the 2017 document. This document only covers FAO Area 31 (Western Central Atlantic, Fig. 1).

¹ Prepared by Jeremy J. Mendoza FAO Consultant




Figure 1. Area of competence of the Western Central Atlantic Fisheries Commission (FAO Statistical Area 31 is delineated in light orange)

2. WECAFC manages nearly 15 million km² of marine area extending from Cape Hatteras in North Carolina, United States of America (35°N) to south of Cape Recife, Brazil (10°S). This area covers the south-east coast of the United States, the Gulf of Mexico, the Caribbean Sea and the north-east coast of South America. Approximately 51% of the mandate area is in areas beyond national jurisdiction (ABNJ) and around 81% corresponds to waters with depths greater than 400 m. Except for Northern Brazil which is included in FAO Area 41, the rest of the management area corresponds to FAO Area 31 (Table 1 and Fig. 1). The Caribbean Sea supports the highest species diversity in the Atlantic Ocean and is a global-scale hot spot of marine biodiversity (Roberts *et al.*, 2002). This species diversity supports industrial and small scale fisheries along the region which target mollusks, crustaceans and fishes inhabiting from shallow coastal areas to deep abyssal slopes. The productivity of the Western Tropical Atlantic is quite heterogeneous; the most productive regions are related to riverine input and upwelling systems, especially in the Gulf of Mexico and along the Northeastern shelf of South America. Also, coral reefs, coastal lagoons, mangrove forests and seagrass beds are highly productive and sustain important fishery resources. The diversity and complexity of these coastal and marine ecosystems represents a serious challenge for fisheries research and management in this area.

Table 1. Location and area of major shelf zones in the WECAFC area (Stevenson, 1981)

LOCATION	AREA (km ² in 1000s)	FAO AREA
Continental shelf		
USA Southeast coast	110	31
Gulf of Mexico	600	31
Yucatan-Eastern Venezuela	250	31
Guyana, Suriname, French Guiana	200	31
Northern Brazil	360	41
TOTAL continental shelf	1520	
Islands		
Islands and offshore banks	380	31
GRAND TOTAL	1900	

Profile of Landings

3. Information on landings are reported to FAO on a yearly basis by member countries. Only commercial landings are reported, excluding information on other components of catch, such as discards and recreational landings. The commercial landings data provided to FAO must be used with caution to infer status of different species and/or species groups and ideally should be complemented with information from stock assessments, trends in effort or catch per unit of effort, and other sources (i.e. expert knowledge). However, due to diverse causes (i.e. high species diversity, limited technical and financial resources, relative importance of diverse small scale fisheries) in many cases such additional information is not available.

4. At present FAO landings data covers the period from 1950 to 2016. As the early years (1950s to 1970s) reflect the period of development for most fisheries in the region, for presentation purposes only the period from 1975 to 2016 will be covered in the present document. Total landings from Area 31 peaked at 2.5 million tonnes in 1984, landings then declined steadily until 1992 when 1.6 million tonnes were reported. Subsequently landings increased and stabilized above 1.7 million tonnes between 1993 and 2004, afterwards landings declined more or less steadily until 2014 when a minimum of 1.2 million tonnes was reported. In 2015 and 2016 total landings have been increasing and reached 1.4 million tonnes in 2015 and 1.6 million tonnes in 2016. To a great extent the evolution of landings has been driven by variations in ISSCAAP group 35 (herrings, sardines and anchovies) which represent the main proportion of total landings (Fig. 2).




Figure 2. Total reported landings (in thousand tonnes) by ISSCAAP groups in the Western Central Atlantic (1975-2015).

5. Landings from ISSCAAP group 35 represented 48% of total landings reported in 2016. The most important species in this group is Gulf menhaden (*Brevoortia patronus*), reported mainly by the United States of America, with reported landings of 618 719 tonnes in 2016, followed by round sardinella (*Sardinella aurita*), landed mainly by Venezuela, with total landings in 2016 of 123 377 tonnes. Both species showed an increase over landings in 2015 that represented overall more than 126 000 tonnes. On the other hand, Atlantic menhaden (*Brevoortia tyrannus*) has presented historically low landings levels over the last decade (Fig. 3a).

6. The Flathead grey mullet (*Mugil cephalus*), reported mainly by the United States of America and Mexico, showed stability in landings over the period 2012-2015 with values around 9 600 tonnes after minima recorded in 2010 and 2011, in 2016 landings increased to 14 584 tonnes. Mulletts nei (family Mugilidae), mainly reported by Mexico and Venezuela, dropped sharply in 2016 to 2 093 tonnes after reaching 8 456 tonnes in 2015. While grunts nei (family Haemulidae), mainly landed in Mexico, Venezuela and Cuba, showed a smaller decrease and reached 5 703 tonnes in 2016 (Fig. 3b).

7. A selection of soft bottom demersal fishes from ISSCAAP group 33, such as Sea catfishes nei, mainly landed in Mexico and Venezuela, showed an increase during the period 2012-2015, but decreased to 6 646 tonnes in 2016. The spotted weakfish (*Cynoscion nebulosus*), reported by Mexico and the United States of America, also showed a slight decrease in 2016 after a recent small peak in landings during the period 2013-2015. Weakfishes nei, mainly from Mexico, Venezuela and French Guiana, reached 6 053 tonnes in 2016. Whitemouth croaker (*Micropogonias furnieri*), reported only by Venezuela, increased to 2 926 tonnes after a decreasing trend since the early 2000s (fig. 3c).




Figure 3a. Selected species in ISSCAAP group 35

Figure 3b. Selected species in ISSCAAP group 33




Figure 3c. Selected soft bottom species

Figure 3d. Selected groupers and snappers

8. Groupers nei (family Serranidae), reported in recent years mainly by Dominican Republic and Venezuela, continued to show a decreasing trend up to 2016; however, this decrease may be partially explained by changes in taxonomic resolution from reports by the United States of America. Snappers nei (family Lutjanidae), reported mainly by Venezuela, Mexico, Dominican Republic, and Antigua and Barbuda, showed a decreasing trend, mainly from Venezuelan landings, in recent years. Northern red snapper (*Lutjanus campechanus*), reported by Mexico and the United States of America, continued the recent increasing trend and reached 8 007 tonnes in 2016. While Red grouper (*Epinephelus morio*), reported by the United States of America and Dominican Republic, shows stable landings around 2 700 tonnes since 2012 (Fig. 3d). Red grouper landings from Mexico, which is a main producer, are reported as part of the group Groupers, seabasses, nei.

9. ISSCAAP group 36 includes tunas and tuna-like fishes. Skipjack tuna (*Katsuwonus pelamis*) is presently reported mainly by Venezuela and recent total landings have remained relatively stable in

recent years and reached 3 004 tonnes in 2016. Yellowfin tuna (*Thunnus albacares*) landings, reported among others by Venezuela, Suriname, Trinidad and Tobago, Grenada and Mexico, have been relatively stable over the last decade around 14 000 tonnes and reached 16 328 tonnes in 2016. Albacore tuna (*Thunnus alalunga*), of which the major producers are Taiwan, Venezuela, and Saint Vincent and the Grenadines, has continued its slight recovery and reached 4 672 tonnes in 2016. Landings of Atlantic bonito (*Sarda sarda*) showed an increase to 3 721 tonnes mainly from reports by Mexico (Fig. 3e).

10. Other species from ISSCAAP group 36, such as Blackfin tuna (*Thunnus atlanticus*), reported mainly by Cuba, Grenada, Venezuela and Saint Lucia, showed a slight increase and reached 1 390 tonnes in 2016. Swordfish (*Xiphias gladius*) landings have remained low at 2 789 tonnes in 2016, after a relatively recent peak just above 5 000 tonnes in 2005. Atlantic sailfish (*Istiophorus albicans*) showed a recovery in 2016 with total landings reaching 1 325 tonnes in 2016, while Wahoo (*Acanthocybium solandri*) continued a decreasing trend in recent years and reached 721 tonnes in 2016 (Fig. 3f).

11. King mackerel (*Scomberomorus cavalla*) landings, mainly from Mexico, United States of America, Venezuela, Trinidad and Tobago, and Guyana, decreased slightly to 9 516 tonnes in 2016. Atlantic Spanish mackerel (*Scomberomorus maculatus*), reported by Mexico and the United States of America, continued its recent increasing trend and reached 10 408 tonnes in 2016. On the other hand, landings of Serra Spanish mackerel (*Scomberomorus brasiliensis*) and cero (*Scomberomorus regalis*) have remained at recent historically low levels in 2014-2016 (Fig. 3g).

12. ISSCAAP group 45 includes several species of highly valued crustaceans. Northern brown shrimp (*Farfantepenaeus aztecus*), Northern pink shrimp (*Farfantepenaeus duorarum*) and Northern white shrimp (*Litopenaeus setiferus*) are reported mainly by Mexico and the United States of America. Northern brown shrimp has been fluctuating around 51 000 tonnes over the last decade and showed a decrease to 48 288 tonnes in 2016 after a recent peak in 2015. After three years (2013-2015) of stable landings around 42 000 tonnes, Northern white shrimp landings reached 53 106 tonnes in 2016. Northern pink shrimp landings showed a decrease to 6 861 tonnes in 2016. Penaeus shrimps *nei*, reported mainly by Mexico, Venezuela, Nicaragua and Honduras, have continued to decline and reached a recent minimum of 6 395 tonnes in 2016, which is partially related to increased taxonomic resolution in landings from Venezuela. Atlantic seabob (*Xiphopenaeus kroyeri*) landings are reported to FAO mainly by Guyana, Suriname, the United States of America and Mexico. Overall landings increased in 2016 and reached 29 981 tonnes. (Fig. 3h).

13. Caribbean spiny lobster (*Panulirus argus*) landings are reported to FAO by 27 coastal and island nations from Area 31, among which the Bahamas, Cuba, Dominican Republic, Honduras, Nicaragua and the United States of America are the major producers. Overall fishery landings peaked in 1999 at 35 787 tonnes and since then decreased to a minimum around 24 000 tonnes in 2009. In recent years landings show an increasing trend with total landings reaching 33 136 tonnes in 2016. Blue crab (*Callinectes sapidus*) landings are reported mainly by Mexico, the United States of America and Venezuela. Total landings of blue crab peaked in 1994 at 74 827 tonnes, since then landings have dropped somewhat and remained relatively stable reaching 54 116 tonnes in 2016. Around 30 coastal and island nations report Stromboid conchs *nei*, among which the most important producers are Nicaragua, Bahamas, Jamaica, Honduras, Dominican Republic, Turks and Caicos, and Belize. Total landings have been stable around 30 000 tonnes in recent years and reached 31 931 tonnes in 2016 (Fig. 3i).




Figure 3e. Selected species in ISSCAAP group 36




Figure 3f. Selected species in ISSCAAP group 36




Figure 3g. Selected species from ISSCAAP group 36




Figure 3h. Selected species from ISSCAAP group 45

14. Common dolphinfish (*Coryphaena hippurus*) landings are reported mainly by France, Venezuela, Saint Lucia, Barbados and the Dominican Republic. Landings have shown a more or less continuous increase and stabilized around 6 000 tonnes from 2013 to 2015, however, landings dropped significantly to 3 943 tonnes in 2016. Flying fishes nei are reported mainly by Barbados, landings stabilized around 2 600 tonnes between 2008 and 2010, but have since dropped significantly and reached 544 tonnes in 2016. Decreased landings of flying fish have been associated with reduced catchability during Sargassum influx events (Ramlogan *et al.* 2017). Jacks, crevalles nei are reported mainly by Mexico, after recent minima observed from to 2005 to 2010 landings have recovered and fluctuated around 12 000 tonnes with registered landings of 10 556 tonnes in 2016 (Fig. 3j).

15. Landings from ISSCAAP group 38 (Sharks, rays, chimaeras) peaked at 37 000 tonnes in 1994 and then declined more or less continuously until 2002 when around 24 000 tonnes were reported; since then landings have fluctuated between 22 000 tonnes and 35 000 tonnes reaching 23 492 tonnes in 2016. Requiem sharks nei (Family Carcharhinidae), reported mainly by Mexico and Venezuela, show an overall decreasing trend since the mid 1990s and reached 1 112 tonnes in 2016; this decrease in recent years is partly due to increased taxonomic resolution for certain species (Blue shark, Atlantic sharpnose shark and Blacktip shark, among others) included within this group. Landings of Rays, mantas nei, reported mainly by Mexico, Venezuela and Cuba, have been decreasing steadily since the early 2000s and reached 1 689 tonnes in 2015; significant recent decreases in reports by Mexico and Venezuela may be attributed to increased taxonomic resolution, as reports for Southern stingray (*Hypanus americanus*) started in 2011 and represented 6 734 tonnes in 2016 (Fig. 3k).




Figure 3i. Selected invertebrate species




Figure 3j. Selected species in ISSCAAP group 37




Figure 3k. Selected species in ISSCAAP group 38

Resource Status and Fisheries Management

16. There are several institutional arrangements among countries in the region in order to promote and facilitate the responsible utilization of the fisheries and other aquatic resources within Area 31. Each organization has a different geographical coverage and mandate: Western Central Atlantic Fisheries Commission (WECAFC) of the FAO, the International Commission for the Conservation of Atlantic Tunas (ICCAT), the Caribbean Regional Fisheries Mechanism (CRFM), the Caribbean Fisheries Management Council (CFMC), the Latin American Organization for Fishery Development (OLDEPESCA), the Central American Organization for the Fisheries and Aquaculture Sector (OSPESCA), the Association of Caribbean States (ACS), the Organization of Eastern Caribbean States (OECS), and the fisheries authorities of the different nations in Area 31. However, despite the relatively large number of existing arrangements, information that can be used for management purposes still needs to be improved.

17. Information on the status of the stocks was collected from all the institutional sources referred to above and by searching the World Wide Web. This information allowed to update the stock status of several species and species groups. Table 2 is used by FAO to synthesize information on the status of stocks in area 31. It includes 33 species/stocks that are allocated both a stock status and a level of uncertainty on status determination. The following paragraphs provide information on stock status determination for these groups, as well as for other species for which recent stock assessments were available.

18. For Flathead grey mullet recent assessments were available for the states of Florida (Chagaris et al., 2014) and Louisiana (West et al., 2016a). In Florida the assessment indicated that stocks (west and east coasts) were not overfished and that overfishing was not occurring. Subsequently, commercial landings in the state of Florida reported by the National Marine fisheries Service (NMFS) have decreased from 5 988 in 2014 to 2 707 tonnes in 2016. For the state of Louisiana assessment results indicated that the grey mullet stock was neither overfished nor experiencing overfishing. In Mexico most landings are from the state of Tamaulipas and have remained relatively stable for the period 2011-2016. The fishery in Mexico is considered to be fully exploited (Carta Nacional Pesquera, 2017). In Venezuela landings in 2016 were 2 814 tonnes, which represented 30 percent of the mean peak landings for the period 1992-1996.

19. For Red grouper (*Epinephelus morio*) the most recent stock assessment indicates overexploitation and overfishing in the USA Southeastern Atlantic coast (SEDAR, 2017a). In the USA Gulf of Mexico with data up to 2013 the stock was neither overfished nor experiencing overfishing (SEDAR, 2015), recent NMFS Gulf of Mexico landings for 2014-2016 showed a slight decrease from 3 007 tonnes in 2014 to 2 404 tonnes in 2016. In Mexican waters of the Gulf of Mexico assessments results indicated that for 2015 the stock was overexploited (Carta Nacional Pesquera, 2017), and recent landings data in the state of Yucatan, which represents around 80 percent of total landings, have fluctuated between 7 000 tonnes and 8 000 tonnes in the period 2010-2015. There are no recent assessments for Nassau grouper (*Epinephelus striatus*), but due to declining trends the Nassau grouper is classified as endangered on the IUCN red list and threatened under the USA Endangered Species Act (Sherman et al., 2016) and is considered overfished in the United States Caribbean region (NOAA, 2017) and in Cuba (Baisre, 2018). A recent assessment of gag grouper (*Mycteroperca microlepis*) in USA Gulf of Mexico waters indicated that the stock was not overfished nor experiencing overfishing (SEDAR, 2016a). A gag grouper stock assessment for the Southeastern

coast of the USA indicated that the stock was not overfished but that it was experiencing overfishing in 2012 (SEDAR, 2014a), and NMFS data show reduced landings for 2013-2016 for this stock. Snowy grouper (*Hyporthodus niveatus*) was assessed in the USA Gulf of Mexico by data-limited methods (SEDAR, 2016b) and most stock indicators showed high probabilities of overfishing and overfished conditions. A recent assessment of black seabass (*Centropristis striata*) in USA South Atlantic waters indicated that the stock was not overfished nor undergoing overfishing (SEDAR, 2018a).

20. Grunts (family Haemulidae) are reported mainly by the Mexico, Venezuela and Cuba. There are no recent stock assessments available for species within this family. Landing histories from Cuba and Colombia show significant declines and are considered overexploited in Cuba (Baisre, 2018), while landings from Mexico show significant recoveries in recent years. Data from commercial catch rates of the grunts complex in the state of Florida, USA, where most of the grunts are landed from this country, suggests full exploitation (FWRI, 2016a).

21. Several taxonomic groups are included in Mulletts nei. These include Mulletts nei, White (*Mugil curema*), Lebranche (*Mugil liza*), Bobo (*Joturus pichardi*) and Parassi (*Mugil incilis*) mulletts. These species are reported by several countries including the United States of America, Mexico, Venezuela, Colombia, Dominican Republic and Cuba, among others. The overall trend in the time series for Mulletts nei is decreasing since 1980, when a maximum of 25 457 tonnes was reached, until 2016 when 2 093 tonnes were reported. However, this trend in recent years is misleading as it is apparently mainly the result of improved taxonomic reporting to FAO by the United States of America and Mexico. However, Mulletts nei show significant decreases in Colombia and Cuba (see also, Baisre, 2018). Landings of Lebranche mullet are mainly from Venezuela and considering the landings history it would seem that Lebranche mullet is fully exploited in Venezuela. White mullet is mainly exploited in the state of Veracruz in Mexico and landings have oscillated around 3 000 tonnes between 2003 and 2016, and the species is considered fully exploited (Carta Nacional Pesquera, 2017).

22. The largest proportion of landings from the family Sciaenidae are reported as Weakfishes nei by Venezuela, Mexico and French Guiana. Landings from Venezuela reached a peak of 20 045 tonnes in 1995 and decreased to 3 518 tonnes in 2015. Additionally, for 2016 landings from Venezuela decreased significantly due to increased taxonomic resolution for sciaenids. Mexico and French Guiana data only cover the period 2005-2016 and show relatively stable landings in recent years. In the United States the black drum (*Pogonias cromis*) has been assessed using data-limited methods along the Atlantic coast (Maine to Florida) as a single stock, and it is considered not overexploited nor overfished (ASFMC, 2015a). In USA waters of the Gulf of Mexico, where most of the landings for this species occur, it is considered abundant and not overfished by the Gulf States Marine Fisheries Commission (GSMFC), but no formal recent assessments were available. A stock assessment of Spotted weakfish (*Cynoscion nebulosus*) in waters of the state of Louisiana (USA) indicated that the stock was neither overfished nor experiencing overfishing (West *et al.*, 2014). On the other hand a stock assessment in waters of the state of Mississippi indicated overexploitation (Leaf *et al.*, 2016). Squeteague (*Cynoscion regalis*) is reported only by the United States of America in Area 31. Landings in 1980 reached 4 682 tonnes and then decreased significantly and reached 15 tonnes in 2015. A stock assessment for the Atlantic coast of the United States (FAO Areas 21 and 31) indicated that the stock was overfished but not experiencing overfishing in 2014 (ASMFC, 2016). Red drum (*Sciaenops ocellatus*) is only reported by the United States of America. Landings reached a peak of 2 510 tonnes in 1976 and then decreased abruptly and reached 58 tonnes in 2015. The most recent assessment indicates that the Southern Atlantic stock (from South Carolina to Florida)

is likely not experiencing overfishing, but it was not possible to establish if the stock was overfished or not (ASMFC, 2017a).

23. For Northern red snapper (*Lutjanus campechanus*) the most recent assessments estimated that the USA South Atlantic stock was overfished and experiencing overfishing (SEDAR, 2017b), while the Gulf of Mexico stock has been rebuilding and is no longer considered overfished nor undergoing overfishing (SEDAR, 2018b). In Mexico Northern red snapper landings averaged around 3 000 tonnes for the period 2010-2015. The resource is considered fully exploited for the state of Tabasco, but deteriorating for other states in the Gulf of Mexico (Carta Nacional Pesquera, 2017). Vermillion snapper (*Rhomboplites aurorubens*) is reported by Mexico, Puerto Rico, Venezuela and USA. Total landings showed a peak in 2005 at 4 055 tonnes and then declined to 3 435 in 2016. The most recent assessment in USA waters of the Gulf of Mexico indicated that the stock was not overfished nor undergoing overfishing (SEDAR, 2016c). An assessment from the South Atlantic coast of the USA indicated that the stock was not overfished nor experiencing overfishing (SEDAR, 2018c). An assessment report for Mutton snapper (*Lutjanus analis*) from the Southeast Atlantic and Gulf of Mexico USA waters indicated that the stock was neither overexploited nor experiencing overfishing in 2013 (O’Hop et al., 2015). Recent landing reports from NMFS have been below the Allowable Biological Catch limit. In Cuba Mutton snapper is considered overfished based on national landing reports (Baisre, 2018). A recent assessment of the Gulf of Mexico Grey snapper (*Lutjanus griseus*) indicated that the stock is not overfished but that it is undergoing overfishing (SEDAR, 2018d).

24. Atlantic menhaden (*Brevoortia tyrannus*) is only exploited by the USA in Area 31. FAO landing reports indicated a maximum of 139 874 tonnes in 1981, which then declined sharply and reached 216 tonnes in 2016. However, the stock of Atlantic menhaden extends beyond Area 31 as it is distributed from the waters of the state of Maine to those of the state of Florida. The most recent assessment for this stock, with data up to 2016, indicated that the stock was not overfished nor undergoing overfishing (ASMFC, 2017b). Atlantic thread herring (*Opisthonema oglinum*) is reported to FAO in Area 31 mainly by the USA, Venezuela and Cuba. Landings from Venezuela and the USA show significant declines, but there are no assessments available for these stocks. In Cuba it is estimated to be fully fished (Baisre, 2018). Gulf menhaden (*Brevoortia patronus*) is only reported to FAO by the United States of America. The most recent benchmark stock assessment (SEDAR, 2013a) used data for the period 1977-2011 and indicated that the stock was not overfished and overfishing was not occurring. A recent update of the benchmark assessment with data up to 2015 arrived at the same conclusions (Schueller, 2016). Round sardinella (*Sardinella aurita*) is reported to FAO in Area 31 by Venezuela, the USA and Mexico. Venezuela is the major producer and for the period 1950-2016 accounted on average for more than 99 percent of total reported landings. Landings have been increasing since a minimum of 34 753 tonnes in 2011 to 122 646 tonnes in 2016. This recent trend may indicate that the stock is recovering from an overfished condition.

25. The most recent assessment for Albacore tuna (*Thunnus alalunga*) indicated that stock status in the North Atlantic had improved and was not overfished nor experiencing overfishing (ICCAT, 2016a). In the case of Western Atlantic Skipjack tuna (*Katsuwonus pelamis*) the stock was assessed in 2014 and results indicated that the stock was unlikely to be overexploited (ICCAT, 2014). For Yellowfin tuna (*Thunnus albacares*) a single stock for the Atlantic Ocean is used for management purposes by ICCAT and most of the landings are reported from the Tropical Eastern Atlantic (FAO Areas 34 and 47). The most recent assessment (ICCAT, 2016b) indicated that the Yellowfin tuna stock was slightly overfished (biomass 5% below B_{MSY}) but not undergoing overfishing in 2014. Atlantic bonito (*Sarda sarda*) landings are reported by a number of countries in Area 31, among

which the most important are Mexico, Venezuela, and to a lesser extent Trinidad and Tobago, the United States of America and France (Martinique). Overall landings show an increasing trend until the 1990s when peak production reached 4 740 tonnes in 1994, landings then dropped to a low of 1 187 in 2008, and later increased to 3 721 in 2016 mainly due to landings from Mexico, while landings from Venezuela and Trinidad and Tobago remained at historically low levels. There are no data on migratory movements or stock structure of Atlantic bonito in the Western Central Atlantic. Atlantic bonito is within the different species under ICCAT, but no stock assessments are available due to insufficient data (ICCAT, 2017a).

26. Atlantic sailfish (*Istiophorus albicans*) is reported mainly by Venezuela, Grenada, Dominican Republic and Cuba. In the most recent assessment of the Western Atlantic stock (ICCAT, 2016c), due to uncertainty in different model estimates, the working group recommended that landings should not exceed current levels. The stock is considered not likely to be overexploited nor overfished (ICCAT, 2018). Blue marlin (*Makaira nigricans*) is exploited by several mainland and island countries in Area 31 with total landings in 2016 of 705 tonnes. A stock assessment with data up to 2016 estimated that the Atlantic stock of Blue marlin was overfished and undergoing overfishing (ICCAT, 2018). Bigeye tuna (*Thunnus obesus*) landings are reported to FAO by several regional and Asian countries. Most of the landings are associated with longliners from Japan, Taiwan (Prov. Of China) and Korea, while main producing countries from the region include Venezuela, and Saint Vincent and the Grenadines. For management purposes ICCAT considers a unit stock for the Atlantic Ocean. The most recent assessment estimated that the stock was overfished and undergoing overfishing (ICCAT, 2018). Atlantic bluefin tuna (*Thunnus thynnus*) is reported mainly by the United States of America and Mexico, the most recent assessment for the Western Atlantic stock indicates that the resource is not undergoing overfishing but it was not possible to estimate biomass reference points (ICCAT, 2017b). Swordfish (*Xiphias gladius*) is reported mainly by Spain and the United States of America in Area 31. Landings peaked at 5 255 tonnes in 1989, after a decline in the 1990s landings increased to exceed 4 000 tonnes during most of the period 2003-2013, but then declined to 2 789 tonnes in 2016. For management purposes ICCAT considers three stocks: Mediterranean, North Atlantic and South Atlantic. The most recent assessment for the North Atlantic stock estimated that the stock was neither overfished nor undergoing overfishing (ICCAT 2017c). Blackfin tuna (*Thunnus atlanticus*) is reported to FAO by several mainland and island nations in Area 31. Main producers are Venezuela, Dominican Republic, Cuba, France (Martinique and Guadeloupe) and Saint Lucia. Overall landings showed a peak in 1992 at 5 480 tonnes and later declined to reach 1 390 tonnes in 2016. Several country series showed declining trends, except for Cuba and to a lesser extent Saint Lucia. ICCAT considers that available data are insufficient to assess the resource (ICCAT, 2017a).

27. Atlantic Spanish mackerel (*Scomberomorus maculatus*) landings are reported by Mexico and the United States of America in Area 31, with Mexico being the major producer. Atlantic Spanish mackerel assessments have been done for the South Atlantic and Gulf of Mexico regions of the USA. In the South Atlantic assessment results indicated that the stock was neither overfished nor experiencing overfishing (SEDAR, 2012b) and recent NMFS landings have been below estimated MSY levels. In the Gulf of Mexico, results indicated that the stock was neither overfished nor experiencing overfishing; however, no clear determination of stock status was made from the assessment, as the independent reviewers differed on the appropriateness of the assessment for establishing these determinations (SEDAR, 2013b). In Mexican waters no assessments are available, but based on landing statistics Atlantic Spanish mackerel is considered fully exploited (Carta Nacional Pesquera, 2017), which is consistent with recent FAO landings data. Cero (*Scomberomorus regalis*) landings are only reported by island states in Area 31, among which the most important are Puerto Rico,

Dominican Republic and France (Martinique and Guadeloupe). However, most of the country time series are incomplete. Overall landing reports showed that production exceeded 500 tonnes from 1958 to 1986 with peaks of 800 tonnes in 1959-1960 and 1971-1972; however, landings have declined considerably since 1987 and reached 250 tonnes in 2016. King mackerel (*Scomberomorus cavalla*) landings are reported to FAO by a number of countries, among which the most important are Mexico, USA, Venezuela, Dominican Republic, and Trinidad and Tobago. Total landings increased during most of the time series and reached a peak of 13 214 tonnes in 2004, and later decreased somewhat reaching 9 516 tonnes in 2016. Stock assessments of King mackerel have been made in USA waters of the South Atlantic and Gulf of Mexico coasts. In the South Atlantic assessment results indicated that the stock was not overfished nor undergoing overfishing (SEDAR, 2014b). In the Gulf of Mexico assessment results indicated that the King mackerel stock was not overfished nor undergoing overfishing (SEDAR, 2014c). Data for 2014-2016 from NMFS landing reports for both stocks have remained relatively stable. In Mexican waters the stock is considered to be fully exploited in accordance with the landings history of the species in the Gulf of Mexico area (Carta Nacional Pesquera, 2017), which is supported by recent FAO landings data. There are no data on migrations or stock structure for King mackerel in the Southeastern Caribbean, if a single stock is assumed in this area it is likely that the stock is fully exploited based on the combined landings histories of Venezuela and Trinidad and Tobago. Landings of Serra Spanish mackerel (*Scomberomorus brasiliensis*) are reported to FAO in Area 31 by Venezuela, Trinidad and Tobago, Guyana and Grenada, with Venezuela and Trinidad and Tobago being the major producers. Total landings increased more or less steadily until 1991 when peak landings of 7 419 tonnes were reached, landings remained above 4 000 tonnes until 2007, but later decreased more or less steadily until 2016 when 1 975 tonnes were reported. This general pattern is more or less similar for Venezuela and Trinidad and Tobago and the landing histories would suggest that Serra Spanish mackerel is overexploited in these two countries.

28. Sharks, rays, chimaeras are reported by several countries, among which the major producers are Mexico, Venezuela, Cuba and Spain. Total landings from this group peaked at 36 982 tonnes in 1994 and have since then declined to reach 23 492 tonnes in 2016. Blue shark (*Prionace glauca*) landings are mainly reported by Spain. Total landings peaked at 11 569 tonnes in 2011 and declined sharply to 1 939 in 2016. This highly migratory species has been assessed by ICCAT and results indicated that Blue shark in the North Atlantic is not likely to be overfished nor experiencing overfishing (ICCAT, 2015b). Shortfin mako shark (*Isurus oxyrinchus*) landings are mainly reported by Spain. Overall landings reached 537 tonnes in 2012 and decreased to 159 tonnes in 2016. The most recent assessment by ICCAT of the North Atlantic stock of Shortfin Mako shark estimated that the stock was overfished and undergoing overfishing (ICCAT, 2017d). Additionally, at present there are 2 species of sharks (Daggernose shark, *Isogomphodon oxyrinchus*, and Oceanic whitetip shark, *Carcharhinus longimanus*) and three species of batoids (Small-tooth sawfish, *Pristis pectinata*, Large-tooth sawfish, *Pristis pristis*, and Bancroft's numbfish, *Narcine bancroftii*) which are classified as Critically Endangered by the IUCN Red List of Threatened Species in the region. Furthermore, the Scalloped hammerhead, *Sphyrna lewini*, the Great hammerhead, *Sphyrna mokarran*, and the Bigeye thresher shark, *Alopias superciliosus*, are classified as Endangered by the IUCN (WECAFC, 2017).

29. The United States of America is the major producer of Blue crab (*Callinectes sapidus*) in FAO Area 31. A stock assessment was carried out for blue crab in USA waters of the Gulf of Mexico (GDAR, 2013) based on 2 stocks: an eastern stock centered on Tampa Bay in the state of Florida and a western stock centered on the coast of the state of Louisiana, and also including the coasts of the states of Alabama, Mississippi and Texas. The base model found that both stocks in 2011 were not

overfished nor undergoing overfishing, although the Western stock was in a depressed state and approaching an overfished limit. An assessment of blue crab in Louisiana waters, where most of the landings of the Western Gulf of Mexico stock come from, indicated that in 2015 the stock was overfished and experiencing overfishing (West et al., 2016b). An update assessment with data up to 2017 estimated that the stock was no longer overfished nor experiencing overfishing (West et al., 2018). Blue crab is considered fully exploited in Mexican waters of the Gulf of Mexico (Carta Nacional Pesquera, 2017), FAO data show an increasing trend with maximum historical catches of nearly 18 000 tonnes in 2016. In Venezuela blue crab landings reported to FAO cover the period 1989-2016, landings increased significantly from 2 353 tonnes in 1989 to a maximum of 18 701 tonnes in 2007, landings remained above 14 900 tonnes until 2010 but later declined steeply to reach 3 900 tonnes in 2014, but have recently recovered and reached 7 119 tonnes in 2016. This landing history suggests that blue crab is fully exploited or overexploited in Venezuela.

30. Caribbean spiny lobster (*Panulirus argus*) landings are reported to FAO by 27 coastal and island nations from Area 31, among which the United States of America, Mexico, Cuba, the Bahamas, Nicaragua and Honduras are the major producers. Overall fishery landings peaked in 1999 at 35 787 tonnes and since then have decreased slightly to 33 136 tonnes in 2016. Several country reports (Anguilla, Antigua and Barbuda, Belize, Bahamas, Mexico, Nicaragua, Cuba) indicate a fully fished or stable status, while for other countries (Caribbean Netherlands, Dominican Republic, France, Honduras, Panama and United States of America) the status of stocks is unknown (FAO, 2015). In Puerto Rico the stock is considered to be undergoing overfishing but it is unknown whether the stock is overfished (NOAA, 2017). A major issue regarding assessment and management of Caribbean spiny lobster is stock connectivity through larval drift. For example, in the most recent stock assessment in waters of the United States of America it was not possible to clearly establish stock status due to uncertainty in recruitment sources (SEDAR, 2010). At present efforts by regional and national institutions are underway in order to collect data and assess the resource from a Caribbean wide perspective (FAO, 2015).

31. The most recent assessment for northern brown shrimp (*Farfantepenaeus aztecus*) in USA waters of the Gulf of Mexico (Hart, 2016a) indicated that the resource was not overfished nor undergoing overfishing. The Northern brown shrimp is considered to be fully exploited in Mexican waters of the Gulf of Mexico (Carta Nacional Pesquera, 2012, DOF, 2014b), which is consistent with recent reported landings to FAO. Northern pink shrimp (*Farfantepenaeus duorarum*) is reported to FAO by the United States of America, Cuba and Mexico. The United States of America reports the largest proportion of landings in Area 31. The most recent assessments in Gulf of Mexico waters of the USA determined that Northern pink shrimp was neither overexploited nor undergoing overfishing (Hart, 2016b; Hart, 2017). In Mexico Northern pink shrimp is considered overfished (Carta Nacional Pesquera, 2012: DOF, 2014c), landings showed an increase in 2014 and 2015 but declined in 2016. In Cuba landings of Northern pink shrimp have been declining consistently since the mid 1970s, and landings in 2016 represented around 12 percent of average maximum landings. Northern white shrimp (*Litopenaeus setiferus*) landings are reported to FAO by the United States of America, which is the main producer in these records, and Mexico. The most recent assessment in Gulf of Mexico waters of the USA indicated that the resource was not overfished nor experiencing overfishing (Hart, 2016c).

32. *Penaeus* shrimps *nei* are reported to FAO by 15 countries bordering the Western Central Atlantic. The major producers are Mexico, Venezuela, Suriname, Guyana, Nicaragua and French Guiana. Overall landings show a steep decline over the last decade which may be partly explained

by increased taxonomic resolution in reports from Mexico. Venezuela has been traditionally the second largest producer of this group in the area and after minima observed in 2009 and 2010, following the industrial trawl ban, recent landings have increased and reached 2 276 tonnes in 2015. Landings from Suriname peaked during 1970-1975, when landings exceeded 6 900 tonnes, landings have declined significantly in recent years and reached 377 tonnes in 2016. In Guyana landings peaked at 8 652 tonnes in 1978 and since then have declined considerably to 411 tonnes in 2016, and in French Guiana landings peaked in 1995 at 4 455 tonnes and started to decline sharply after the late 2000s and reached 700 tonnes in 2016. Other nations such as Honduras and Colombia also show apparent overexploitation of stocks, as landings in 2016 were below 30% of landings in years of maxima. Redspotted shrimp (*Farfantepenaeus brasiliensis*) are landed by several countries in Area 31, but are only reported to FAO by the USA and French Guiana, with only one year of data each in the time series. However, the species is exploited by other countries in the region and, for example, in Mexico it is considered overexploited (DOF, 2014d). Rock shrimp (*Sicyonia brevirostris*) landings are reported to FAO only by the USA. Landings show strong yearly fluctuations with a period of high catches during the 1990s, when landings peaked at 4 409 in 1998, since then there has been a decrease to 297 tonnes in 2016, the southern Atlantic stock is not considered to be undergoing overfishing, but stock status is unknown (NOAA, 2017). In Mexico the rock shrimp is considered overexploited (DOF, 2014d). Royal red shrimp (*Pleoticus robustus*) is only reported by the USA in Area 31. The landings history shows important fluctuations with a peak in 2013 at 589 tonnes and then decreased to 138 tonnes in 2016. Royal red shrimp in the Gulf of Mexico is not considered to be undergoing overfishing, but the status of the stock is unknown (NOAA, 2017).

33. Several species of Stromboid conchs are exploited in Area 31 of which the Queen conch (*Lobatus gigas*) is the most important fishery resource. Total landings reached a maximum of 41 727 tonnes in 1994 and later declined to 22 334 tonnes in 2007, but have recovered in recent years and reached 31 931 tonnes in 2016. To a great extent this increase is related to landings from Nicaragua which represented only 890 tonnes in 2007 and reached 11 400 tonnes in 2016. For some major producers such as Bahamas, Belize and Nicaragua the stocks are considered to be fully exploited or stable, however for other major producers such as Honduras, and the Turks and Caicos stock status is unknown (FAO, 2016). In Dominican Republic, also a major producer, the shallow water fraction is considered overexploited (FAO, 2016). In Caribbean waters of the United States of America the species is overfished but not undergoing overfishing (NOAA, 2017). In Mexico there are fishing bans for Yucatan and Chinchorro Bank (Carta Nacional Pesquera, 2012) and the species is considered overfished (Carta Nacional Pesquera, 2017). In Panama a fishing ban was extended for an additional five years in 2009, and in Colombia the fishery was closed in 2010-2012 and reopened during a short period in 2013-2014 under a quota system (FAO, 2016). Recently, a draft Regional Queen Conch Management plan has been reviewed and expert agreement has been reached on harmonized conversion factors for different forms of conch meat processing in the region (FAO, 2016).

34. American cupped oyster (*Crassostrea virginica*) is only reported by the USA and Mexico. The USA has traditionally been the major producer and landings peaked at 147 012 tonnes in 2000, since then landings have decreased substantially and reached 10 007 tonnes in 2016. Biomass estimates of American cupped oysters in Louisiana, which accounts for most of the landings from the USA in Area 31, indicate resource overexploitation (Louisiana Wildlife and Fisheries, 2015). In Mexico, landings have been much more stable and peaked at 56 599 tonnes in 1989 and later decreased somewhat and reached 44 313 tonnes in 2016. The resource is considered fully exploited (Carta Nacional Pesquera, 2012), which is supported by relatively stable landings in recent years in FAO data. Mangrove cupped oyster (*Crassostrea rhizophorae*) landings are reported mainly by

Venezuela and Cuba, but also by Colombia and Dominican Republic. Overall landings show a peak of 6 878 tonnes in 1990 and later decreased sharply to 9 tonnes in 2016. The landings history in Venezuela suggests overexploitation. On the other hand, Cuba stopped reporting the species in 1997, albeit the series in the previous years were above 1 800 tonnes, Baisre (2018) reported a reduction of national landings from Cuba of 68%. For this species, along its range, habitat degradation and coastal pollution are also important factors affecting fisheries production.

35. Calico scallop (*Argopecten gibbus*) is only reported by the USA in Area 31. The data is characterized by several pulses related to recruitment variations. Landings reached a peak of 400 000 tonnes in 1984 and the last pulse was observed in 1994-1995 with peak landings of 74 325 tonnes in 1994; since then there have been no landing reports to FAO for this species. Among Other Scallops, Pectens, American sea scallop (*Placopecten magellanicus*) is reported only by the USA in Area 31. Landings of this species are highly irregular and showed peak catches of 3 966 tonnes in 1978 and of 5 749 tonnes in 1987, since then landings have dropped markedly and only 2 tonnes were reported in 2015. However, it should be noted that the northern part of Area 31 is the southern distributional range of this temperate species and the population is assessed as a single stock for FAO areas 21 and 31. The most recent stock assessment indicated that the stock was not overfished nor experiencing overfishing (ASMFC, 2014). The Atlantic bay scallop (*Argopecten irradians*) is also reported only by the USA. The landing history shows strong fluctuations with peaks of 2 917 tonnes in 1968 and 2 625 tonnes in 1986, landings have since then dropped steeply and no landings were reported in 2014 and 2015. This would indicate an overfished condition for the species in area 31. However, habitat loss and toxic algal blooms have also been indicated as drivers of population decline in FAO areas 21 and 31 (Oreska et al., 2017).

References

- ASMFC (Atlantic States Marine Fisheries Commission). 2014. Stock assessment for Atlantic sea scallops in 2014, updated through 2013. 59th SAW Assessment Report. 300 pp.
- ASMFC. 2015a. Black Drum Stock Assessment and Peer Review Reports. 351 pp.
- ASMFC. 2016. Weakfish Benchmark Stock Assessment and Peer Review Report. 270 pp.
- ASMFC. 2017a. Red Drum Benchmark Stock Assessment & Peer Review Report. 126 pp.
- ASMFC. 2017b. Atlantic Menhaden Stock Assessment Update. 180 pp.
- Baisre, J.A. 2018. An overview of Cuban commercial marine fisheries: the last 80 years. *Bull. Mar. Sci.* 94. <https://doi.org/10.5343/bms.2017.1015>
- Carta Nacional Pesquera. 2012. Acuerdo por el que se da a conocer la Actualización de la Carta Nacional Pesquera. Publicado en el Diario Oficial de la Federación 24 de agosto de 2012. 236 pp.
- Carta Nacional Pesquera. 2017. Acuerdo por el que se da a conocer la Actualización de la Carta Nacional Pesquera. Publicado en el Diario Oficial de la Federación 11 de junio de 2018. 268 pp.

Cass-Calay, S.L., Porch, C.E., Goethel, D.R., Smith, M.W., Matter, V. & McCarthy, K.J. 2015. Stock Assessment of Red Snapper in the Gulf of Mexico 1872 – 2013 - with Provisional 2014 Landings. SEDAR Update Assessment. 242 pp.

Chagaris, D., Addis, D. & Mahmoudi, B. 2014. The 2014 stock assessment update for striped mullet, *Mugil cephalus*, in Florida. Florida Fish and Wildlife Conservation Commission, 76 pp.

CRFM (Caribbean Regional Fisheries Mechanism) 2012. Report of Eighth Annual Scientific Meeting – Kingstown, St. Vincent and the Grenadines, 20 - 30 June 2012 – Fishery Management Advisory Summaries. CRFM Fishery Report – 2012, Volume 2. 78p.

DOF (Diario Oficial de la Federación). 2014a. Acuerdo por el que se da a conocer el Plan de Manejo Pesquero de Mero (*Epinephelus morio*) y especies asociadas en la Península de Yucatán. Publicado en el Diario Oficial de la Federación 25 de noviembre de 2014. 76 pp.

DOF. 2014b. Acuerdo por el que se da a conocer el Plan de Manejo Pesquero de Camarón Café (*Farfantepenaeus aztecus*) y Camarón Blanco (*Litopenaeus setiferus*) en las costas de Tamaulipas y Veracruz. Publicado en el Diario Oficial de la Federación 12 de marzo de 2014. 58 pp.

DOF. 2014c. Acuerdo por el que se da a conocer el Plan de Manejo Pesquero de Camarón Rosado (*Farfantepenaeus duorarum*) de la Sonda de Campeche. Publicado en el Diario Oficial de la Federación 28 de marzo de 2014. 68 pp.

DOF. 2014d. ACUERDO por el que se da a conocer el Plan de Manejo Pesquero para las especies de camarón rojo (*Farfantepenaeus brasiliensis*) y de Roca (*Sicyonia brevirostris*) de los Caladeros de Contoy, Quintana Roo. Publicado en el Diario Oficial de la Federación 25 de marzo de 2014. 40 pp.

FAO. 2015. Report of the first meeting of the OSPESCA/WECAFC/CRFM/CFMC Working Group on Caribbean Spiny Lobster, Panama City, Panama, 21–23 October 2014. FAO Fisheries and Aquaculture Report, No. 1095. Bridgetown, Barbados. 112 pp.

FAO. 2016. Report of the second meeting of the CFMC/OSPESCA/WECAFC/CRFM Working Group on Queen Conch, Panama City, Panama, 18–20 November 2014. FAO Fisheries and Aquaculture Report, No. 1097. Bridgetown, Barbados. 435 pp.

FWRI (Fish and Wildlife Research Institute) 2016a. Grunts, Family Haemulidae. Fish and Wildlife Research Institute. Florida Fish and Wildlife Conservation Commission. 7 pp.

GDAR (Gulf Data, Assessment, and Review). 2013. GDAR 01 – Stock Assessment Report. Gulf of Mexico Blue Crab. Gulf States Marine Fisheries Commission. Ocean Springs, MS. 313 pp.

Hart, R.A. 2016a. Stock Assessment Update for Brown Shrimp (*Farfantepenaeus aztecus*) in the U.S. Gulf of Mexico for 2015. NOAA Fisheries, Southeast Fisheries Science Center, Galveston Laboratory, Galveston, TX. 19 pp.

Hart, R.A. 2016b. Stock Assessment Update for Pink Shrimp (*Farfantepenaeus duorarum*) in the U.S. Gulf of Mexico for 2015. NOAA Fisheries, Southeast Fisheries Science Center, Galveston Laboratory, Galveston, TX. 17 pp.

Hart, R.A. 2016c. Stock Assessment Update for White Shrimp (*Litopenaeus setiferus*) in the U.S. Gulf of Mexico for 2015. NOAA Fisheries, Southeast Fisheries Science Center, Galveston Laboratory, Galveston, TX. 20 pp.

Hart, R.A. 2017. Stock Assessment Update for Pink Shrimp (*Farfantepenaeus duorarum*) in the U.S. Gulf of Mexico for the 2016 Fishing Year. NOAA Fisheries, Southeast Fisheries Science Center, Galveston Laboratory, Galveston, TX. 17 pp.

ICCAT (International Commission for the Conservation of Atlantic Tunas). 2014. Report of the 2014 ICCAT East and West Atlantic Skipjack Stock Assessment Meeting. ICCAT, Madrid. 97 pp.

ICCAT. 2015b. Report of the 2015 ICCAT Blue Shark Stock Assessment Session. ICCAT, Madrid. 116 pp.

ICCAT. 2016a. Report of the 2016 ICCAT North and South Atlantic Albacore Stock Assessment Meeting. ICCAT, Madrid. 99 pp.

ICCAT. 2016b. Report of the 2016 ICCAT Yellowfin Tuna Stock Assessment Meeting. ICCAT, Madrid. 103 pp.

ICCAT. 2016c. Report of the 2016 Sailfish Stock Assessment. ICCAT, Madrid. 68 pp.

ICCAT. 2017a. Report of the 2017 Small Tunas Species Group Intersessional Meeting. Collect. Vol. Sci. Pap. ICCAT, 74(1): 1-75.

ICCAT. 2017b. Report of the 2017 ICCAT Bluefin Stock Assessment Meeting. ICCAT, Madrid. 106 pp.

ICCAT. 2017c. Report of the 2017 ICCAT Atlantic Swordfish Stock Assessment Session. ICCAT, Madrid. 85 pp.

ICCAT. 2017d. Report of the 2017 ICCAT Shortfin Mako Assessment Meeting. ICCAT, Madrid. 64 pp.

ICCAT. 2018. Report of the Standing Committee on Research and Statistics. ICCAT, Madrid. 469 pp.

Leaf, R., Dippold, D. & Hendon, R. 2016. 2016 Stock Assessment for Spotted Seatrout, *Cynoscion nebulosus*, in Mississippi. Mississippi Department of Marine Resources. Office of Marine Fisheries. 97 pp.

Louisiana Wildlife and Fisheries. 2015. 2015 Oyster Stock Assessment Report of the Public Oyster Areas of Louisiana. Seed Grounds and Seed Reservations. Oyster Data Report Series No. 21. 116 pp.

NOAA (National Oceanic and Atmospheric Administration). 2017. Status of Stocks 2017. Annual Report to Congress on the Status of U.S. Fisheries. National Oceanic and Atmospheric Administration, National Marine Fisheries Service.

O'Hop, J., Muller, R.G. & Addis, D.T. 2015. Stock Assessment of Mutton Snapper (*Lutjanus analis*) of the U.S. South Atlantic and Gulf of Mexico through 2013. SEDAR Update Assessment. Florida Fish and Wildlife Conservation Commission. 144 pp.

Oreska, M.P.J., Truitt, B., Orth, R.J. & Luckenbach M.W. 2017. The bay scallop (*Argopecten irradians*) industry collapse in Virginia and its implications for the successful management of scallop-seagrass habitats. *Marine Policy*, 75: 116-124.

Ramlogan, N.R., P. McConney and H.A. Oxenford. 2017. Socio-economic impacts of Sargassum influx events on the fishery sector of Barbados. Centre for Resource Management and Environmental Studies, the University of the West Indies, Cave Hill Campus, Barbados. CERMES Technical Report No. 81: 86pp.

Roberts, C.M., McClean, C.J., Veron, J.E., Hawkins, J.P., Allen, J.R., et al., 2002. Marine biodiversity hotspots and conservation priorities for tropical reefs. *Science* 295, 1280–1284.

Schueller, A. 2016. GDAR 02. Gulf menhaden stock assessment. 2016 Update. GSFMC No. 254. Gulf States Marine Fisheries Commission. Ocean Springs, MS. 75 pp.

SEDAR (Southeast Data, Assessment, and Review). 2010. Stock assessment of spiny lobster, *Panulirus argus*, in the Southeast United States. SEDAR 8 Update Assessment Workshop Report. Key West, Florida. 128 pp.

SEDAR. 2012. SEDAR 28 – South Atlantic Spanish mackerel Stock Assessment Report. SEDAR, North Charleston SC. 444 pp.

SEDAR. 2013a. SEDAR 32A - Gulf of Mexico menhaden Stock Assessment Report. SEDAR, North Charleston SC. 422 pp.

SEDAR. 2013b. SEDAR 28 – Gulf of Mexico Spanish Mackerel Stock Assessment Report. SEDAR, North Charleston SC. 712 pp.

SEDAR. 2014a. Stock Assessment of Gag off the Southeastern United States. SEDAR Update Assessment. Southeast Fisheries Science Center. National Marine Fisheries Service. 112 pp.

SEDAR. 2014b. SEDAR 38 – Stock Assessment Report. South Atlantic King Mackerel. SEDAR, North Charleston SC. 502 pp

SEDAR. 2014c. SEDAR 38 – Stock Assessment Report. Gulf of Mexico King Mackerel. SEDAR, North Charleston SC. 465 pp.

SEDAR. 2015. SEDAR 42 – Gulf of Mexico Red Grouper Assessment Report. SEDAR, North Charleston SC. 612 pp.

SEDAR. 2016a. SEDAR 33 – Update Report. Gulf of Mexico Gag Grouper. SEDAR, North Charleston SC. 123 pp.

SEDAR 2016b. SEDAR 49 – Stock Assessment Report. Gulf of Mexico Data-limited Species: Red Drum, Lane Snapper, Wenchman, Yellowmouth Grouper, Speckled Hind, Snowy Grouper, Almaco Jack, Lesser Amberjack. SEDAR, North Charleston SC. 618 pp.

SEDAR. 2016c. SEDAR 45 – Stock Assessment Report. Gulf of Mexico Vermillion Snapper. SEDAR, North Charleston SC. 188 pp.

SEDAR. 2017a. SEDAR 53 – South Atlantic Red Grouper Assessment Report. SEDAR, North Charleston SC. 159 pp.

SEDAR. 2017b. Stock Assessment of Red Snapper off the Southeastern United States. SEDAR Benchmark Assessment. Southeast Fisheries Science Center. National Marine Fisheries Service. 145 pp.

SEDAR. 2018a. South Atlantic Black Seabass. SEDAR 56 Stock Assessment Report. SEDAR, North Charleston SC. 164 pp.

SEDAR. 2018b. Gulf of Mexico Red Snapper. SEDAR 52 Stock Assessment Report. North Charleston SC. 434 pp.

SEDAR. 2018c. South Atlantic Vermillion Snapper. SEDAR 55 Stock Assessment Report. North Charleston SC. 170 pp.

SEDAR. 2018d. Gulf of Mexico Gray Snapper. SEDAR 51 Stock Assessment Report. North Charleston SC. 428 pp.

Sherman, K.D., Dahlgren, C.P., Stevens, J.R. & Tyler, C.R. 2016. Integrating population biology into conservation management for endangered Nassau grouper *Epinephelus striatus*, Mar. Ecol. Prog. Ser., 554: 263-280.

Stevenson, D.K. 1981. A review of the marine resources of the WECAFC region. FAO Fisheries Technical Paper, No. 211. 134 pp.

WECAFC. 2017. Draft recommendation on applying a precautionary approach to fisheries of threatened species of sharks and rays in the WECAFC area. 1st meeting of the WECAFC/OSPESCA/CRFM Working Group on Sharks. Barbados, 17-19 October 2017.

West, J., Decossas, G., Melancon, A., Potts, S. & Powers, J.E. 2014. Update Assessment of Spotted Seatrout *Cynoscion nebulosus* in Louisiana Waters 2014 Report. Louisiana Department of Wildlife and Fisheries. 54 pp.

West, J., Decossas G. & Powers, J. 2016a. Update Assessment of Striped Mullet *Mugil cephalus* in Louisiana Waters 2016 Report. Louisiana Department of Wildlife and Fisheries, 38 pp.

West, J., Blanchet, H., Marx, J. & Powers, J.E. 2016b. Update Assessment of Blue Crab in Louisiana Waters 2016 Report. Louisiana Department of Wildlife and Fisheries. 30 pp.

West, J. Blanchet, H. & Cagle, P. 2018. Update Assessment of Blue Crab in Louisiana Waters. 2018 Report. Louisiana Department of Wildlife and Fisheries. 33 pp.

Table 2. Stock status based on FAO classification. (F: fully exploited; O: overexploited; U: non-fully exploited; L: low; M: medium; H: high)

Name	Species	Main fishing countries in 2015	State 2015	Uncertainty 2015	State 2016	Uncertainty 2016	Sources
Flathead grey mullet	<i>Mugil cephalus</i>	Mexico, USA, Venezuela	F,F, O	M	F,F, O	M	Carta Nacional Pesquera, 2017; Chagaris et al. 2014; West et al. 2016a
Groupers		Mexico, USA, Venezuela	F/O	M	F/O	L	DOF, 2014a; SEDAR 2017a; SEDAR 2015; SEDAR 2018a; Baisre 2018
Grunts, sweetlips nei	<i>Haemulidae</i>	Venezuela, Mexico, Cuba	F/O	M	F/O	H	FWRI 2016a; Baisre 2018
Mullets nei	<i>Mugilidae</i>	Mexico, USA, Venezuela	F/O	H	F/O	H	Based on landing history for Cuba
Sciaenids	<i>Sciaenidae</i>	Venezuela, USA, Mexico	F/O	M	F/ O	M	ASMFC, 2015a; West et al. 2014; ASMFC, 2016; ASMFC 2017a
Snappers	<i>Lutjanidae</i>	Mexico, USA, Venezuela	F/O	M	F/O	M	SEDAR, 2017b; Cass-Calay et al., 2015; O'Hop et al., 2015; Baisre 2018; SEDAR 2018b and 2018c
Other Miscellaneous coastal fishes			F/O	M	F/ O	H	Based on landing histories
33-Miscellaneous coastal fishes					F	M	Landing histories
34-Miscellaneous demersal fishes							
Atlantic menhaden	<i>Brevoortia tyrannus</i>	USA	F	L	F	L	ASMFC 2017b
Atlantic thread herring	<i>Opisthonema oglinum</i>	Venezuela, Cuba	F/?	H	F/?	H	Landing histories
Gulf menhaden	<i>Brevoortia patronus</i>	USA	F	M	F	L	SEDAR, 2013 ^a ; Schueller, 2016
Round sardinella	<i>Sardinella aurita</i>	Venezuela	F/O	H	F/O	H	Landing history Venezuela
Other Herrings, sardines, anchovies			O,F	H	O, F	H	Landing histories
35-Herrings, sardines, anchovies							
Albacore	<i>Thunnus alalunga</i>	Taiwan, Venezuela	F	L	F	L	ICCAT, 2016a
Atlantic bonito	<i>Sarda sarda</i>	Mexico	F/O	H	F/O	H	Landing histories
Atlantic Spanish mackerel	<i>Scomberomorus maculatus</i>	Mexico, USA	F	M	F	M	SEDAR, 2012b; SEDAR, 2013b, Carta Nacional Pesquera, 2017
Cero	<i>Scomberomorus regalis</i>		?		?		
King mackerel	<i>Scomberomorus cavalla</i>	Mexico, Venezuela, USA, Trinidad Tob	F	M	F	M	SEDAR, 2014b; SEDAR 2014b;

Serra Spanish mackerel	<i>Scomberomorus brasiliensis</i>	Trinidad Tob, Venezuela, Guyana	O	H	O	H	Landing histories
Skpkack tuna	<i>Katsuwonus pelamis</i>	Venezuela, Cuba			F	L	ICCAT, 2014
Yellowfin tuna	<i>Thunnus albacares</i>	Mexico, USA, Venezuela	O	L	O	L	ICCAT, 2016b
Other Tunas, bonitos, billfishes			O,F	M	O, F	M	ICCAT, 2012a; ICCAT, 2015a; 2017b; 2017c
36-Tunas, bonitos, billfishes							
38-Sharks, rays, chimaeras		Cuba, Mexico, Spain, Venezuela	F/O	M	F/O	M	ICCAT, 2015b; ICCAT, 2017d
39-Marine fishes not identified			F	H	F	H	Global landing history Area 31
42-Crabs, sea-spiders	<i>Callinectes sapidus</i>	Mexico, USA, Venezuela	F/O	M	F/O	M	West et al., 2016b; West et al., 2018; Carta Nacional Pesquera, 2017
Caribbean spiny lobster	<i>Panulirus argus</i>	Bahamas, Cuba, Honduras, Nicaragua	F	H	F	H	FAO, 2015; Landing histories
Other Lobsters, spiny-rock lobsters							
43-Lobsters, spiny-rock lobsters							
Atlantic seabob	<i>Xiphopenaeus kroyeri</i>	Guyana, Suriname, Mexico	F	M	F	M	CRFM, 2014; CRFM, 2012; Landing histories
Northern brown shrimp	<i>Penaeus aztecus</i>	Mexico, USA	F	L	F	L	Hart, 2016a; Carta Nacional Pesquera, 2012; DOF, 2014
Northern pink shrimp	<i>Penaeus duorarum</i>	Mexico, USA	F/O	M	F/O	M	Hart, 2016b; Hart, 2017 and landing histories
Northern white shrimp	<i>Penaeus setiferus</i>	USA	F	L	F	L	Hart, 2016c
Penaeus shrimps nei	<i>Penaeus spp</i>	Mexico, Nicaragua, Venezuela	F/O	H	F/O	H	Landing histories
Redspotted shrimp	<i>Penaeus brasiliensis</i>		O	H	O	H	DOF, 2014d
Rock shrimp	<i>Sicyonia brevirostris</i>	USA, Mexico	O	H	O/?	H	DOF, 2014d, NOAA, 2017
Royal red shrimp	<i>Pleoticus robustus</i>	USA	F	H	F	H	Landing history
Other Shrimps, prawns					F	H	Landing histories
45-Shrimps, prawns							
Stromboid conchs nei	<i>Strombus spp</i>	Bahamas, Nicaragua, Jamaica	F/O	H	F/O	H	CRFM, 2012; NOAA, 2016; FAO 2016
Other Abalones, winkles, conchs		Mexico	F	H	F	H	Landing history
52-Abalones, winkles, conchs							
American cupped oyster	<i>Crassostrea virginica</i>	Mexico, USA	F/O	M	F/O	M	Louisiana, Wildlife and Fisheries, 2015; Carta Nacional Pesquera, 2012
Other Oysters			O	H	O	H	Landing histories
53-Oysters							

Calico scallop	<i>Argopecten gibbus</i>	USA	?		?		Pulse fishery highly dependent on recruitment.
Other Scallops, pectens		USA	O,F	M	O,F	M	ASMFC, 2014 and landing history
55-Scallops, pectens							

Table 3. Stock assessment methods and status for selected stocks or locations in FAO Area 31 (F=Fully fished; O= Overfished; ?= Unknown)

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
<i>ISSCAAP 33</i>				
Flathead Grey Mullet <i>Mugil cephalus</i>	Florida West coast	F	Stock Synthesis.	Data up to 2013. Aprox. 25% of FAO total landings. Not overfished nor overfishing in 2013. NMFS landings in 2014 similar to 2013. However, significant drop (~50%) for 2015 and 2016.
	Florida East coast	F	Stock Synthesis.	Data up to 2013. Aprox. 4% of FAO total landings. Not overfished nor overfishing in 2013. NMFS landings in 2014 similar to 2013. Significant drop (~50%) in 2015 and 2016.
	Mexico (mainly Tamaulipas state)	F	Landings history	Aprox. 40% of FAO total landings. Considered fully exploited by INAPESCA. Increasing trend for Tamaulipas state from 2011 to 2015.
	Venezuela	F/O	Landings history	Aprox. 30% of FAO total landings. Landings in 2016 were 30% of average maximum landings (AML)
Mulletts nei	Cuba	O	Landings history	Aprox. 6% of FAO total landings. Landings in 2016 were 14% of AML
Lebranche mullet <i>Mugil liza</i>	Venezuela	F	Landings history	100% of FAO total landings. Landings in 2016 were 65% of AML
Red Grouper <i>Epinephelus morio</i>	USA Southeast Atlantic	O	Statistical Catch at Age	Data up to 2015. Landings in 2016 < 1% of total landings. Stock overfished and undergoing overfishing.
	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2013. Aprox. 38% of total landings. Not overfished nor overfishing. Landings from 2014 to 2016 relatively stable.
	Mexican Gulf of Mexico	O	Virtual Population Analysis	Data up to 2015. Mexican FAO landings of Groupers nei have remained at high levels. Aprox. 60% of total landings.
Nassau Grouper <i>Epinephelus striatus</i>	Bahamas	O	Landings history	Aprox. 60% of total FAO landings. Landings in 2016 7% of AML. Considered Endangered by IUCN

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
Nassau Grouper <i>Epinephelus striatus</i>	Cuba	O	Landings history	Aprox. 39% of total FAO landings. Landings in 2016 < 2% of AML.
Gag Grouper <i>Mycteroperca microlepis</i>	USA Southeast Atlantic	F	Statistical Catch at Age	Data up to 2012. Aprox. 20% of total FAO landings. Not overfished but overfishing in 2012. NMFS landings decreasing for 2013-2016.
	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2015. Aprox. 80% of total FAO landings. Not overfished nor overfishing in 2015. NMFS landings increased from 294 t in 2015 to 491t in 2016.
Snowy grouper <i>Hyporthodus niveatus</i>	USA Gulf of Mexico	O	Data limited methods	Data up to 2014. Aprox. 54% of total landings. High probability of overfished and overfishing. NMFS landings stable between 2015 and 2016.
	USA Southeast Atlantic	O	Statistical Catch at Age	Data up to 2012. Aprox. 45% of total landings. Overfished but no overfishing in 2012. NMFS landings increasing from 2013 to 2016.
Black Seabass <i>Centropristis striata</i>	USA Southeast Atlantic	F	Statistical Catch at Age	Data up to 2016. Aprox. 100% of total landings. Not overfished nor overfishing.
Red Drum <i>Sciaenops ocellatus</i>	USA Southeast Atlantic	?	Statistical Catch at Age	Data up to 2013. Aprox. 11% of total landings. No overfishing but not possible to estimate overfished status. Reduction in landings from 2014 to 2016.
Squeteague <i>Cynoscion regalis</i>	USA Atlantic coast	O	Bayesian Catch at Age	Data up to 2014. 100% of total landings. Stock overfished but no overfishing.
Spotted Weakfish <i>Cynoscion nebulosus</i>	Louisiana state, USA	F	Statistical Catch at Age	Data up to 2013. Aprox. 35% of total landings. Not overfished nor overfishing in 2013.
	Mississippi state, USA	O	Statistical Catch at Age	Data up to 2014. Aprox. 5% of total landings. Overfished and overfishing in 2014. NMFS landings relatively stable from 2014 to 2016.
	Mexican Gulf of Mexico	O	Landings history	Aprox. 60% of total landings. FAO landings in 2016 were 13% of AML

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
Black Drum <i>Pogonias cromis</i>	USA Atlantic coast	F	Depletion Based Stock Reduction Analysis	Data up to 2012. Aprox. 4% of total landings. Not overfished nor experiencing overfishing. NMFS landings for 2013-2016 below MSY estimates.
	USA Gulf of Mexico	F	Landings history	Aprox. 92% of FAO landings. Landings in 2016 12% higher than AML. Considered abundant and not overfished by GSMFC.
	Mexican Gulf of Mexico	O	Landings history	Aprox. 4% of FAO landings. Landings in 2016 were 8% of AML.
Lane snapper <i>Lutjanus synagris</i>	Mexico	O	Landings history	Aprox. 10% of total FAO landings. Landings in 2016 were 11% of AML.
	Cuba	O	Landings history	Aprox. 78% of total FAO landings. Landings in 2016 were 32% of AML.
Mutton Snapper <i>Lutjanus analis</i>	USA Gulf of Mexico	F	Statistical Catch at Age	Data up to 2013. Aprox. 84% of total landings. Not overfished nor overfishing. Landings in 2014-2016 below Allowable Biological Catch
Northern Red Snapper <i>Lutjanus campechanus</i>	USA Southeast Atlantic	O	Statistical Catch at Age	Data up to 2014. Less then 1% of total landings. Overfished and overfishing. At F=0 rebuilding not before 2025.
	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2016. Aprox. 37% of total landings. Not Overfished nor overfishing.
	Mexican Gulf of Mexico	O	Landings history	Aprox. 60% of total landings. Landings in 2010 about 1000 t and considered overfished. Landings in 2016 >4000 t from non traditional areas.
Vermillion Snapper <i>Rhomboplites aurorubens</i>	USA Southeast Atlantic	F	Statistical Catch at Age	Data up to 2016. Aprox. 10% of total landings. Not overfished nor overfishing.
	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2014. Aprox. 27% of total landings. Not overfished nor overfishing. NMFS landings in 2015 and 2016 lower than in 2014.

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
Grey snapper <i>Lutjanus griseus</i>	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2015. Not overfished but overfishing occurring.
<i>ISSCAAP 35</i>				
Atlantic menhaden <i>Brevoortia tyrannus</i>	USA Atlantic coast	F	Statistical Catch at Age	Data up to 2015. Not overfished nor overfishing. Most landings from Area 21.
Gulf menhaden <i>Brevoortia patronus</i>	USA Gulf of Mexico	F	Statistical Catch at Age	Data up to 2016. Not overfished nor overfishing.
Round sardinella <i>Sardinella aurita</i>	Northeast Venezuela	F	Landings history	Landings have recovered since minimum in 2011. Landings in 2016 at 84% AML.
Atlantic thread herring <i>Opisthonema oglinum</i>	USA South Atlantic and Gulf of Mexico	?	Landings history	Aprox. 33% of total landings. Landings in 2016 at 20% AML. Apparently factors other than fishing affecting abundance.
	Cuba	F	Landings history	Aprox. 58% of total landings. Landings in 2016 at 83% AML.
	Northeast Venezuela	?	Landings history	< 6% total landings. Landings in 2016 at less than 2% AML. Closed reduction factories.
<i>ISSCAAP 36</i>				
Albacore tuna <i>Thunnus alalunga</i>	North Atlantic	F	Biomass Dynamics	Data up to 2014. Not overfished nor overfishing (96.8% probability).
Skipjack tuna <i>Katsuwonus pelamis</i>	Western Atlantic	F	ASPIC and Catch-MSY	Data up to 2013. Not overfished nor overfishing.
Yellowfin tuna <i>Thunnus albacares</i>	Atlantic	O	ASPIC, Virtual Population Analysis and Stock Synthesis	Data up to 2014. Stock is slightly overexploited but overfishing was not occurring
Atlantic bluefin tuna <i>Thunnus thynnus</i>	Western Atlantic	?	Virtual Population Analysis and Stock Synthesis	Data up to 2015. Overfished status was not assessed but biomass has increased from 2004 to 2015. No overfishing.

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
Bigeye tuna <i>Thunnus obesus</i>	Atlantic	O	ASPIC and Stock Synthesis	Data up to 2014. Overfished and overfishing
Swordfish <i>Xiphias gladius</i>	North Atlantic	F	Bayesian biomass dynamics	Data up to 2015. Not overfished nor overfishing occurring
Atlantic sailfish <i>Istiophorus albicans</i>	Western Atlantic	F	Stock Synthesis	Data up to 2014. Not likely to be overfished nor experiencing overfishing.
Blue marlin <i>Makaira nigricans</i>	Atlantic	O	Surplus production and age structured models	Data up to 2016. Stock overfished and experiencing overfishing.
Atlantic Spanish Mackerel <i>Scomberomorus maculatus</i>	USA South Atlantic	F	Stock Synthesis	Data up to 2012. Aprox. 10% of total landings. Not overfished nor overfishing. Landings 2013-2016 below MSY
	Mexican Gulf of Mexico	F	Landings history	Aprox. 81% of total landings. Landings in 2016 at 91% AML
King mackerel <i>Scomberomorus cavalla</i>	USA Southeast Atlantic	F	Stock Synthesis	Data up to 2012. Aprox. 10% of total landings. Not overfished nor overfishing. NMFS landings 2013-2016 below MSY
	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2012. Aprox. 10% of total landings. Not overfished nor overfishing during fishery history.
	Mexico	F	Landings history	Aprox. 58% of total landings. Landings in 2016 16% above AML. Relatively stable since late 1990s
Serra Spanish mackerel <i>Scomberomorus brasiliensis</i>	Venezuela	O	Landings history	Aprox. 45% of total landings. Landings in 2016 21% of AML.
	Trinidad and Tobago	O	Landings history	Aprox. 35% of total landings. Landings in 2016 30% of AML.
Atlantic bonito <i>Sarda sarda</i>	Mexico	F	Landings history	Aprox. 99% of total landings. Landings in 2016 at maximum values.
	Trinidad and Tobago	O	Landings history	Less than 1% of total landings. Landings in 2016 at < 6% AML
	Venezuela	O	Landings history	Less than 1% of total landings. Landings in 2015 at < 1% AML

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
<i>ISSCAAP 38</i>				
Blue shark <i>Prionace glauca</i>	North Atlantic	F	Bayesian Surplus Production	Data up to 2013. Overfished status and overfishing not likely.
Shortfin Mako shark <i>Isurus oxyrinchus</i>	North Atlantic	O	Bayesian Surplus Production	Data up to 2015. Overfished and overfishing
<i>ISSCAAP 42</i>				
Blue crab <i>Callinectes sapidus</i>	Louisiana state	F	Catch Survey Analysis	Data up to 2017. Aprox. 37% of total landings. Not overfished nor overfishing
	Mexican Gulf of Mexico	F	Landings history	Aprox. 33% of total landings. Landings peaked in 2016 at 39% above AML. Considered fully fished.
	Lake Maracaibo, Venezuela	F/O	Landings history	Aprox. 13% of total landings. Landings in 2016 at 43% AML. Landings increasing after sharp decrease from 2009 to 2014.
<i>ISSCAAP 43</i>				
Caribbean spiny lobster <i>Panulirus argus</i>	Bahamas	F	Landings history	Aprox. 26% of total landings. Landings in 2016 at 89% AML.
	Nicaragua	F	Landings history	Aprox. 19% of total landings. Peak landings in 2015 and 2016.
	Cuba	F/O	Landings history	Aprox. 14% of total landings. Landings in 2016 at 37% AML.
<i>ISSCAAP 45</i>				
Atlantic seabob <i>Xiphopenaeus kroyeri</i>	Suriname	F	Bayesian Biomass Dynamics	Data up to 2011. Aprox. 26% of total landings. Not overfished nor overfishing. Recent relatively stable landings and landings in 2016 at 70% AML.
	Guyana	F	Statistical age and weight structured	Data up to 2012. Aprox. 68% of total landings. Not overfished nor overfishing. Landings in 2016 at 10% above AML.
Northern Brown shrimp <i>Farfantepenaeus aztecus</i>	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2015. Aprox. 78% of total landings. Not overfished nor overfishing

ISSCAAP Group and Species	Stock or location	Status	Assessment Method	Comments
Northern Pink Shrimp <i>Farfantepenaeus duorarum</i>	Cuba	O	Landings history	Aprox. 11% of total landings. Landings in 2016 at 11% AML.
Northern White shrimp <i>Litopenaeus setiferus</i>	USA Gulf of Mexico	F	Stock Synthesis	Data up to 2015. Aprox. 97% of total landings. Not overfished nor overfishing
Penaeus shrimps nei	Nicaragua	O	Landings history	Aprox. 24% of total landings. Landings in 2015 at 31% AML
	French Guiana	O	Landings history	Aprox. 11% of total landings. Landings in 2016 at 16% AML
	Suriname	O	Landings history	Aprox. 6% of total landings. Landings in 2016 at 5% AML
<i>ISSCAAP 52</i>				
Stromboid conchs	Nicaragua	F	Landings history	Aprox. 36% of total landings. Landings peaked in 2015-2016 at 22% above AML.
	Bahamas	F	Landings history	Aprox. 8% of total landings. Landings in 2016 at 51% AML.
	Jamaica	F	Density Survey and landings history	Data up to 2011. Aprox. 12% of total landings. In 2011 not overfished nor experiencing overfishing.
<i>ISSCAAP 53</i>				
American cupped oyster <i>Crassostrea virginica</i>	Louisiana state	O	Density survey	Data up to 2014. Aprox. 14% of total landings. Public oyster beds in Louisiana at 34% of target biomass. USA landings at 10% AML in 2016.
	Mexican Gulf of Mexico	F	Landings history	Aprox. 86% of total landings. Landings in 2016 at 87% AML.