



***Pervagor melanocephalus* (Bleeker, 1853)**

En - Blackheaded leatherjacket.

Maximum total length about 10.7 cm. Inhabits coral reefs to depths of 40 m. Widespread in the West Pacific region, from Indonesia to Tonga, and from northeastern Australia to Okinawa.




(after Randall, Allen, and Steene, 1990)




***Pervagor nigrolineatus* (Herre, 1927)**

En - Blacklined leatherjacket.

Maximum total length about 9.7 cm. Inhabits shallow areas near coral reefs, but has also been trawled in deeper waters. Occurs on atolls off the northwestern coast of Australia, in Indonesia, New Guinea, the Solomons, Palau, the Philippines, and Viet Nam.




(after Hutchins, 1986)




***Pseudaluteres nasicornis* (Temminck and Schlegel, 1846)**

En - Rhino leatherjacket.

Maximum standard length about 18 cm. Inhabits trawling grounds to 75 m, but also found in shallow weedy areas. Found in the Indo-West Pacific, from South Africa to eastern Australia and New Caledonia, and north to Japan.





(after Gloerfelt-Tarp and Kailola, 1984)



Pseudomonacanthus elongatus Fraser-Brunner, 1940**En** - Fourbanded leatherjacket.



Maximum total length about 30 cm. Inhabits trawling grounds at depths between 32 and 55 m. Occurs in northwestern Australia to as far east as Cape York.



(after Sainsbury et al., 1985)

Pseudomonacanthus macrurus (Bleeker, 1857)**En** - Smallspotted leatherjacket.



Maximum total length about 24 cm. Inhabits shallow weedy areas and also taken by trawlers. Found in New Guinea, Indonesia, Singapore, and the Philippines.



(after Gloerfelt-Tarp and Kailola, 1984)

Pseudomonacanthus peroni (Hollard, 1854)**En** - Potbellied leatherjacket.

Maximum total length about 45.7 cm. Inhabits trawling grounds at depths between 15 and 114 m. Occurs in northern Australia only.




(after Sainsbury et al., 1985)


Rudarius minutus Tyler, 1970

En - Minute leatherjacket.

Maximum total length about 3 cm. Inhabits shallow areas of dead and live corals. Found in northeastern Australia, New Guinea, Indonesia, Palau, and the Philippines.




(after Kuitert, 1992)




Scobinichthys granulatus (Shaw, 1790)

En - Rough leatherjacket.

Maximum total length 35 cm. Inhabits areas of seagrass and rocky reef. Occurs in Australia's southern half.




(after Allen, 1985)




Stephanolepis cirrhifer (Temminck and Schlegel, 1850)

En - Fringed leatherjacket.

Maximum total length about 24.5 cm. Inhabits shallow areas of weed and rocky reef. Marketed in Japan. Found in Japan, China (including Taiwan Province), and northern Philippines.




(after Masuda et al., 1984)




Thamnaconus hypargyreus* (Cope, 1871)*En** - Yellowspotted leatherjacket.


Maximum total length about 21.2 cm. Inhabits offshore waters at depths between 130 and 235 m where it is taken by trawlers. Marketed in Japan. Occurs in northern Australia, China (including Taiwan Province), and southern Japan.




(after Sainsbury et al., 1985)

***Thamnaconus modestoides* (Barnard, 1927)****En** - Modest filefish; **Fr** - Bourse mingo; **Sp** - Cachúa mingo.


Maximum total length about 38 cm. Inhabits moderately deep water at depths between 72 and 248 m. Found in many scattered locations in the Indian and western Pacific oceans including southeastern Africa, Madagascar, Chagos Archipelago, Andaman Islands, western and eastern Australian coasts, New Caledonia, and southern Japan.



(after Masuda et al., 1984)

***Thamnaconus tessellatus* (Günther, 1880)****En** - Manyspotted leatherjacket.

Maximum total length about 28 cm. Inhabits moderately deep waters at depths between 120 and 236 m. Found in northwestern and northeastern Australia, New Caledonia, Indonesia, the Philippines, and Japan.



(after Sainsbury et al., 1985)

