

CODEX ALIMENTARIUS COMMISSION



Food and Agriculture
Organization of the
United Nations



World Health
Organization

E

Viale delle Terme di Caracalla, 00153 Rome, Italy - Tel: (+39) 06 57051 - E-mail: codex@fao.org - www.codexalimentarius.org

Agenda item 8

CRD04

October 2024

JOINT FAO/WHO FOOD STANDARDS PROGRAMME

CODEX COMMITTEE ON RESIDUES OF VETERINARY DRUGS IN FOODS

27th Session

21-25 October 2024

Omaha, Nebraska, United States of America

REPORT OF THE PHYSICAL WORKING GROUP ON CRITERIA AND PROCEDURES FOR THE ESTABLISHMENT OF ACTION LEVELS FOR VETERINARY DRUGS IN FOOD OF ANIMAL ORIGIN RESULTING FROM UNAVOIDABLE AND UNINTENTIONAL VETERINARY DRUG CARRYOVER IN NON-TARGET ANIMAL FEED

(Prepared by the Physical Working Group chaired by Australia and co-chaired by Canada)

INTRODUCTION

1. A Physical Working Group (PWG) meeting chaired by Australia and co-chaired by Canada took place to further refine the discussion paper on the possible requirements or criteria for developing action levels for compounds in edible tissues/commodities due to the unintended and unavoidable carryover of authorized veterinary drugs in feed and their transfer from feed into food of animal origin and to use nicarbazin and lasalocid as a pilot cases.

WORK PROCESS: PARTICIPATION AND METHODOLOGY

2. The PWG was attended by various Member countries and organizations and observers.
3. The PWG Chairs discussed the last draft discussion document that was shared as part of CX/RVDF 24/27/8. In line with the terms of reference (TOR) of the PWG, the document contained proposed criteria for establishing action levels, a proposed procedure, as well as a pilot studies estimating action levels for the unavoidable and unintended carry-over of nicarbazin and lasalocid in chicken eggs. Several comments were received on this draft as included in CX/RVDF 24/27/8 Add.1, CRD6, CRD7, CRD9, CRD12, CRD13 and CRD14.
4. On the basis of these comments, the PWG Chairs prepared a revised document, which is attached as an appendices to this report (Appendix I) for consideration by Codex members and observers at CCRVDF27.
5. In addition to the comments received, one member provided an alternative approach for consideration.
6. The PWG chairs outlined three potential options for addressing carryover:
 - a. Option 1: Proposed criteria and procedures for the establishment of action levels (Appendix I).
 - b. Option 2: Proposal to develop carry-over risk management information and Risk Management Decision Tool (RMDT) (Appendix III).
 - c. Option 3: Hybrid approach combining options 1 and 2 (Appendix IV).
7. The PWG agreed to refine the criteria for establishing action levels as outlined in the discussion paper, as this is required for the advancement of both options 1 and 3.

SUMMARY OF DISCUSSION

8. The PWG chairs provided an overview of the document and went through the four questions that were posed in CL2024/68-RVDF. The following were the main highlights of the discussion:
 - a. Retain paragraphs 6a and 7 in the general criteria for the proposed approach.
 - b. Retain paragraph 12 while deleting subpoints c, d and e, related to CCRVDF conducting the TMDI calculation.
 - c. Retain paragraph 13 relating to the sources of residue transfer and monitoring data.
 - d. Retain subpoints c and d in paragraph 15 on considerations related to carryover.
 - e. Retain paragraph 23 and all associated notes (1-5) related to the human dietary exposure assessment.
 - f. One member had questions pertaining to the previously agreed paragraph 18 regarding options for assessment of animal dietary exposure.

CONCLUSIONS

9. The WG completed its task as per its TOR. The outcome is presented in the discussion paper attached in Appendix I of this document.
10. The proposal for action levels put forward in the discussion paper aims to provide a pragmatic approach for the establishment of action levels. The criteria proposed (described in the discussion paper), which, when satisfied, supports the estimation of action levels while maintaining protection of the consumer. The use of this approach recognises that unavoidable and unintended carry-over of veterinary drugs from medicated to un-medicated feed occurs and sometimes leads to detectable residues in commodities currently without an MRL.

RECOMMENDATIONS

11. Codex members and observers are invited to consider:
 - i. The proposed approach for the establishment of action levels as presented in the discussion paper (Appendix I).
 - ii. Pilot studies using nicarbazin and lasalocid residues in chicken eggs, as presented in the discussion paper, which illustrates the proposed approach for estimating action levels as presented in Appendix II.
 - iii. The alternate proposal for the *Develop carry-over risk management information and risk management decision tool* (Appendix III).
 - iv. The potential hybrid approach as outlined in Appendix IV.

APPENDIX I**PROPOSED APPROACH FOR ESTABLISHING ACTION LEVELS FOR VETERINARY DRUG RESIDUES IN FOOD PRODUCTS FROM NON-TARGET ANIMALS LINKED TO THE UNINTENDED AND UNAVOIDABLE VETERINARY DRUG CARRY-OVER IN NON-TARGET ANIMAL FEED****Proposal by the Electronic Working Group (For comments)****DEFINITIONS AND TERMS USED IN THIS DOCUMENT**

Action level: Maximum concentration of residue resulting from unintended and unavoidable carry-over in a feed of a veterinary drug (expressed in mg/kg or µg/kg on a fresh weight basis) in a non-target animal that is recommended by the Codex Alimentarius Commission to be legally permitted or recognized as acceptable in or on a food.

Transfer Factor (TF): The ratio between the veterinary drug residue in the tissue or commodity of interest (fat/skin, muscle, liver, kidney, milk, or eggs) and the concentration of veterinary drug in animal feed.

Unavoidable and unintended veterinary drug carry-over in a non-target animal feed: The presence of a veterinary drug in a non-target animal feed caused by the previous manufacture of feed using the same equipment after one or more mitigation procedures have been performed (e.g., flushing, sequencing, or physical clean-out).

Non-target Animal: An animal unintentionally exposed to a veterinary drug not authorized or registered for use in that animal species or production class.

1. Action levels for unavoidable and unintended presence of veterinary drug residues in food products from non- target animals exposed to veterinary drug carry-over in animal feed will be established based on a scientific risk assessment taking into account food safety and whether the best practice has been followed (e.g., *Code of Practice on Good Animal Feeding* (CXC 54-2004), Good Manufacturing Practices (GMP) and Hazard Analysis and Critical Control Point (HACCP)) to minimize the unavoidable and unintended veterinary drug carry-over in non- target animal feed, to a level that is achievable after having implemented mitigation measures according to the Code of Practice on Good Animal Feeding.

General criteria for the proposed approach

2. Action levels for the unintended and unavoidable carry-over of veterinary drugs in non-target animal feed to food should be based on the 'As Low as Reasonably Achievable' concept and only be derived where the framework of the *Code of Practice on Good Animal Feeding*, GMPs, and/or HACCP has been used to minimize the veterinary drug carry-over.
3. Action levels should be developed only to cover situations where low-level residues of an approved/registered veterinary drug used according to good veterinary practices are consistently detected by a [competent] authority in edible commodities from non-target animals and investigations by the [competent] authority confirm the source to be unintended and unavoidable carry-over of a veterinary drug in animal feed.
4. Action levels for non-target animals should be derived only for veterinary drugs authorized for use in a target class of animals.
5. The residues in food resulting from the authorized or registered use of the veterinary drug plus the residues in food resulting from unavoidable and unintended veterinary drug carry-over in animal feed should not result in an exposure that exceeds the established health-based guidance value (HBGV) for the veterinary drug.
6. Action levels should be derived only for residues of veterinary drugs that have adopted (or JECFA recommended) maximum residue limits (MRLs).
 - a) [Action levels should not be established for veterinary drugs for which the Joint FAO/WHO Expert Committee on Food Additives (JECFA) was unable to establish an HBGV or recommend MRLs due to specific human health concerns or inadequate toxicological data.]
7. [Transfer factors (TFs) can be used to estimate the concentration of residues in edible commodities from non- target animals.]

8. Action levels in [edible commodities] may be [derived] from the transfer factors and concentration of unintended and unavoidable veterinary drugs in non-target animal feed after appropriate mitigation steps have been performed (e.g., flushing, sequencing, or physical clean-out) following the manufacture of feed containing the maximum authorized concentration of the drug for the target class of animals.
9. Analytical methods should be available for the veterinary drug residue in the edible commodity for which action levels are proposed.

Procedure

10. The following four steps should be followed for setting action levels for residues of veterinary drugs detected in foods of animal origin determined to be caused by unavoidable and unintended veterinary drug carry-over in non-target animal feed based on the Guidelines on the Application of Risk Assessment for Feed (CX/G 80-2013) and risk assessment approaches.

Step 1. Assess animal dietary exposure assessment.

Step 2. Estimate anticipated residue levels in food commodities of animal origin.

Step 3. Set Action levels.

Step 4. Evaluate human dietary exposure assessment.

11. CCRVDF will perform **Steps 1, 2, and 3**, and then for **Step 4**, CCRVDF may request that JECFA conduct an appropriate exposure assessment based on the proposed action level derived under **Step 3**.
12. [CCRVDF will do an initial Theoretical Maximum Daily Intake (TMDI) calculation, and where there are exceedances, can request such an exposure assessment from JECFA under **Step 4**. CCRVDF may:
 - a) provide JECFA with the proposed action level(s) in the applicable commodity(ies) from **Steps 1-3** and any data that might help with conducting an exposure assessment.
 - b) request JECFA to conduct an exposure assessment that considers exposure from the proposed action level(s) and sources of exposure from the authorized use(s) of the veterinary drug.
13. [Data such as residue transfer and residue monitoring data from peer-reviewed scientific literature and/or data previously reviewed by regulatory authorities may be used by CCRVDF in setting action levels for residues in food products from non-target animals, where it can be concluded that it was due to the unavoidable and unintended veterinary drug carry-over in non-target animal feed.]
14. Residue monitoring data, including trace-back information from a [competent authority], demonstrating that residues are caused by unavoidable and unintended veterinary drug carry-over in non-target animal feed, should be made available to CCRVDF.
15. CCRVDF may consider the following when evaluating the data:
 - a) Does the data demonstrate that unavoidable and unintended carry-over occurs even when mitigation steps are followed (e.g., flushing, sequencing)?
 - b) Does the data demonstrate that unavoidable and unintended carry-over concentrations of the veterinary drug in the feed of non-target animals cause residues in edible commodities from non-target animals?
 - c) [Are the data representative of the various formulations of the veterinary drug available globally?
 - d) Are the data representative of feed mixing practices used globally?
16. The details of the four general steps for setting action levels for residues of veterinary drugs detected in foods of animal origin determined to be caused by unavoidable and unintended veterinary drug carry-over in non-target animal feed are discussed below.

Step 1: Assess animal dietary exposure assessment.

17. The veterinary drug carry-over present in non-target feed or feed ingredients will be identified.
18. The anticipated exposure levels for non-target animals will be estimated considering:
- a) **Option 1:** A default hypothetical carry-over of 1% can be applied to the highest authorized dose of the veterinary drug in feed for the target class of animals in situations where:
 - i) Unintended and unavoidable carry-over has been demonstrated; and
 - ii) Suitable data is not available to establish with certainty that unintended and unavoidable carry-over would occur at a level higher (or lower) than 1%.
 - b) **Option 2:** The maximum observed concentration of unavoidable and unintended veterinary drug carry-over in non-target feed determined in feed mills operating under routine good manufacturing conditions. Monitoring data where investigations cannot verify GMP is unsuitable for this purpose.

Step 2: Estimate anticipated residue levels in food of animal origin.**a) Calculating the Transfer Factors (TFs)**

19. The potential transfer of a veterinary drug from feed to food can be estimated by calculating TFs based on suitable feeding studies on non-target animals fed feed containing the veterinary drug at levels close to the unavoidable and unintentional carry-over levels (e.g., feed, oral capsule).
20. TF can be calculated as follows:

$$TF = \frac{\text{residue level in food of animal origin (milk, eggs or tissues) (fresh weight), expressed in mg/kg}}{\text{level in total feed ration (dry weight), expressed in mg/kg}}$$

Notes (1):

1. The highest individual animal tissue residue level will be used in the TF calculations. The average residue will be used if the highest residue is not reported.
2. In the case of residue levels that are below the limit of quantification of the analytical method (LOQ) and above the limit of detection (LOD) of the analytical method, the TF will be reported as $LOQ \div \text{feed concentration}$.
3. Residue values are less than the LOD of the analytical method will not be used.
4. If there are multiple feeding studies for a particular animal species, studies that fed the veterinary drug at concentrations most representative of the carry-over level should be used preferentially to calculate the TFs.
5. If multiple TFs are derived from drug concentrations in feed close to the carry-over level, the median transfer factor will be used to estimate the anticipated residue levels in edible animal commodities.
6. Survey/monitoring data from national regulatory bodies or reported in the scientific literature may be used to increase confidence in the estimated residue levels in edible tissues resulting from veterinary drug carry-over under good manufacturing practices.
7. TFs should be calculated for one food commodity (e.g., liver) and should not be applied to a different commodity (e.g., eggs).
8. TFs should be calculated for one species and should not be applied to a different species.

b) Calculating the anticipated veterinary drug residue transfer level.

21. Anticipated veterinary drug residue transfer levels in food of animal origin (including muscle, liver, kidney, skin/fat, milk or egg) of non-target animals can be calculated using the TFs, and the level of veterinary drug in the animal's feed estimated either by hypothetical carry-over rates of the highest authorized dose of the veterinary drug in feed for the target-class of animals or the maximum observed carry-over level as measured in non-target feed from feed mills operating under routine good manufacturing conditions. Monitoring data where investigations cannot verify GMP is unsuitable for this purpose.

Anticipated residue level = TF x veterinary drug carry-over level in animals total feed ration (dry weight)

Step 3: Action levels

22. Action levels for food of animal origin from non-target animals can be estimated based on the anticipated residue levels in food of animal origin from animals exposed under practical conditions.

Notes (2):

TF based on a relatively high drug concentration in feed might overestimate the residue concentration in food of animal origin caused by unavoidable and unintended veterinary drug carry-over in animal feed. To account for this, the anticipated residue level in food of animal origin from non-target animals can be the lesser of either:

1. the concentration estimated by using the TF, or
2. the residue concentration determined to be caused by unavoidable and unintended veterinary drug carry-over in animal feed that satisfied bullet point #2 of the General Criteria.

"Action levels should be developed only to cover situations where low-level residues of an approved/registered veterinary drug used according to good veterinary practices are consistently detected by a [competent] authority in edible commodities from non-target animals, and investigations by the [competent] authority confirm the source to be unintended and unavoidable carry-over of a veterinary drug in animal feed".

Step 4 Human dietary exposure assessment

23. An estimate of consumer dietary exposure from residues present at action levels in food of animal origin (eggs, milk, meat, edible offal) from non-target animals will be calculated following approaches for both chronic exposure (based on the ADI) and acute exposure (based on the ARfD, when established).]



[Notes (3):

1. In performing the dietary exposure assessment, exposure to the relevant foods containing residues at the proposed action level(s) and the other sources of dietary exposure from the authorized use(s) of the veterinary drug (e.g., exposure originating from the current Codex MRLs) should be considered.
2. An estimate of the ratios for marker residues to total residues of toxicological or microbiological concern (M:T) may be required.
3. Extrapolation of M:T ratios from one species to a related species (i.e., ruminant to ruminant) is likely feasible if:
 - a) Identical or very similar M:T ratios exist for tissues/commodities of two related species; and/or
 - b) The M:T ratios in tissues/commodities of one related species = 1.
4. Dietary exposure estimates based on the intended use of the veterinary drug plus the residues in food resulting from the proposed action level(s) should not exceed the established health-based guidance value (HBGV).
5. Seek advice from JECFA if the exposure from residues in food resulting from the intended use of the veterinary drug plus the residues in food resulting from the proposed action level(s) exceeds the established health-based guidance value (HBGV).]

APPENDIX II**(For information to support comments on Appendix I)****PILOT STUDY A: ESTIMATING ACTION LEVELS FOR UNAVOIDABLE AND UNINTENTIONAL NICARBAZIN CARRY-OVER IN CHICKEN EGG**

Nicarbazin is a non-ionophoric coccidiostat administered in feed to broiler chickens to prevent and control coccidiosis caused by *Eimeria* spp. Nicarbazin is an equimolar mixture of 4,4'-dinitrocarbanilide (DNC) and 2-hydroxy-4,6-dimethylpyrimidine (HDP). DNC is also known as N,N'-bis(4-nitrophenyl urea) and 1,3-bis(4-nitrophenyl urea). After oral ingestion, the complex dissociates into two major metabolites, DNC and HDP, and both components undergo metabolism via different routes and at different rates. **Table 1** gives a summary of nicarbazin details.

Table 1: Summary of nicarbazin details

Chemical name	an equimolar amount of 4,4'-dinitrocarbanilide (DNC) and 2-hydroxy-4,6-dimethylpyrimidine (HDP). DNC is also known as N,N'-bis(4-nitrophenyl)urea.
Marker residue	4,4'-dinitrocarbanilide (DNC)
Structure	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>DNC</p>  </div> <div style="text-align: center;"> <p>HDP</p>  </div> <div style="text-align: center;"> <p>1 : 1</p> </div> </div> <p>(Tarbin et al., 2005)</p>
Water solubility (20 °C)	DNC - 0.02 mg/L and HDP >10000 mg/L
log K_{ow}	DNC - 3.6 and HDP - 0.94 at pH 5-9 (EFSA, 2003).
Target animal	chickens for fattening, turkeys for fattening
Authorized maximum content in complete feed and Withholding Period (WHP)	125 ppm in the feed, 1 day 40-50 ppm, 0 days (nil) when co-formulated with ionophores (AUS) <125 ppm with 4 days, >125 ppm with 5 days (US) 30-50 ppm with 0 days when co-formulated with narasin (US) 125 ppm, 1 day (EU), 40-50 ppm, 5 days when co-formulated with narasin (EU)
LOQ	0.02 - 0.1 mg/kg for all tissues
ADI	0.9 mg/kg bw (DNC) (JECFA/94/SC, 2022)
MRLs for chicken (broilers) (mg/kg)	AUS muscle 5, liver 35, kidney 20, fat/skin 10, egg 0.3 EU muscle 4, liver 15, kidney 6, fat/skin 4 Canada muscle 4, liver 15, kidney 8, fat/skin 4 Codex muscle 0.2, liver 0.2, kidney 0.2, fat/skin 0.2 JECFA (2022) muscle 4, liver 15, kidney 8, fat/skin 4 US liver 52 UK VMD egg 0.100 (Differential Action Limit, DAL)

Maximum content in feed for non-target species (mg/kg)	EU Regulation EU 574/2011 Feed materials- 1.25 Compound feed for equine species, laying birds and chickens reared for laying (> 16 weeks) – 1.25, other animal species – 3.75 Brazilian Regulation (MAPA 2016) Feed materials - 1.25
---	---

2. Laying hens are identified as the most likely non-target animals to be exposed to unavoidable and unintended carry-over of nicarbazin in non-target animal feed, as feed for chickens and laying hens is often prepared at the same feed mill. Survey or residue monitoring data on nicarbazin in poultry eggs (**Table 2**) and feeding studies on laying hens (**Table 4**) provide evidence for the detectable nicarbazin levels in eggs from laying hens fed feed produced in accordance with good manufacturing practices. **Attachment 1** summarizes the residue data for nicarbazin measured in edible tissues from poultry fed nicarbazin-containing medicated feed.
3. A wide range of nicarbazin levels in eggs with the highest DNC level of 900 µg/kg (**Table 2**) were reported in surveys/residue monitoring of egg samples. As listed in **Table 4**, feeding studies on laying hens resulted in nicarbazin levels in eggs ranging from 226 to 15300 µg/kg. The variation may be explained in part due to the differences in the authorized use-patterns for nicarbazin in broiler chickens. Feeding the broiler chickens with diet containing nicarbazin (in the form of nicarbazin or co-formulated with other ionophores) resulted in liver residue concentrations ranging from around 20-39770 µg/kg, in kidney of 230-5400 µg/kg, in muscle 2-6560 µg/kg, and skin/fat of <10-7750 µg/kg depending on the different feeding levels, WHPs and analytical methodologies (**Attachment 1**). The highest levels of nicarbazin were measured in eggs and poultry liver compared to the other edible poultry tissues.
4. In terms of possible sources of nicarbazin residues in edible commodities of chicken, carry-over of nicarbazin to non-target animal feed during feed manufacturing (Cannavan et al., 2000; Cannavan and Kennedy, 2000; McEvoy et al., 2003) has been identified as a source of nicarbazin residues in egg. Several authors have also highlighted the ingestion of the droppings containing the excreted (non-absorbed) nicarbazin as a possible cause of nicarbazin residues in broiler chicken tissues (Cannavan and Kennedy, 2000; Kan et al., 1996). They have shown that residue levels in the liver but not in the muscle could exceed the concentration of 200 µg/kg in field conditions following the use of 125 mg nicarbazin/kg feed.

Table 2: Survey or residue monitoring data on nicarbazin in poultry egg⁴

Country	Year	Commodity	LOQ (mg/kg)	MRL (mg/kg)	No. of samples tested	Positive sample n>LOQ (n>LOD)	Residue levels (µg/kg)	Highest residue level (µg/kg)	Reference
Australia	2011-2021	egg	0.01	0.3	301	13 (28)	<10-66	66	Australia NRS data
Belgium	2005	egg			320	13		10	Mortier et al., 2005
Belgium	2002-03	egg	a)	a)	a)	a)	3-197 (4), >	a)	Mortier et al., 2005
	2005	b) Poultry, egg, rabbit	b)	b)	b) 6	b)	10 (2)) > 10	b)	
Croatia	2009-2011	poultry egg	0.00015 0.015	0.0005 0.05	307 275		1.85 21.1	122.8 314.4	Bilandžić et al., 2013
EU	2004 - 2005	egg	0.001-0.1		3314	23			EFSA 2008
Ireland	2002-2004	poultry egg			546	9	14-122	122	Danaher et al., 2008
North Ireland	1996-1997	egg (190)		0.001	190	39	4-342	342	Cannavan and Kennedy 2000
Italy	2012	a) Poultry, ovine, eggs	0.001		49 (28, 1, 20)	4	1.4-96	96	Roila et al., 2019
	2013		LOQ		49 (31, 18)	9	12-21	21	
	2014	Poultry, eggs			80 (33, 47)	14	13-238	238	
	2016	Poultry, eggs			58 (34, 24)	20	13-516	516	
	2017	Poultry, eggs Poultry, eggs			46 (34, 12)	d) 13	1-321	321	
UK	1995-2004	chicken egg			2178	123	> 10 DNC	900	UK-VMD, 1995-2004 EFSA 2018
UK	2007	egg		0.025	234	2	40, 60	60	UK, 2007

⁴ Nicarbazin is authorized in the EU and Australia for use in broiler chickens but not approved for laying hens. Residues in eggs are assumed to be from carry-over.

Nicarbazin presence in eggs due to unavoidable and unintended nicarbazin carry-over in animal feed

Step 1. Animal dietary exposure assessment

Option 1

5. A maximum approved rate of 125 mg/kg in broiler feed is considered for proposing action levels in eggs from laying hens utilizing hypothetical carry-over rates. Carry-over of nicarbazin in laying hen feed at a hypothetical level of 1% of the maximum authorized level of 125 mg/kg for broiler chicken would result in carry-over levels of nicarbazin in laying hens feed of 1.25 mg/kg.

Option 2

6. Table 3 summarizes the carry-over levels of nicarbazin in non-medicated animal feed during medicated feed manufacturing. The controlled feed mill studies of Martinez et al. (2018) demonstrated that following medicated feed manufacture at 125 mg/kg nicarbazin and subsequent cleaning and flushing procedures (representing good manufacturing practices), carry-over levels up to 2.2 mg/kg were found in non-medicated feed. This study compared various flushing procedures that reduced the carry-over levels in non-medicated feed. They further claimed that due to the nicarbazin's high electrostatic potential, it tends to cling to the bin walls where the product moisture and environmental conditions may also play roles in its adhesion to the bin walls.

Table 3: Carry-over levels of nicarbazin in non-medicated animal feed during medicated feed manufacturing

Level in medicated feed (mg/kg)	Flushing procedure	Level in flush (mg/kg)	Level in non-medicated diet (mg/kg)	Reference
125	Five flush-size treatments 2.5, 5.0, 10, 15, and 20% of the mixer's total capacity (Forberg 454.5 kg capacity drop bottom paddle mixer)	19.2 14.8 12.0 6.5 5.6	1.8 2.1 2.2 1.4 1.5	Martinez et al., 2018
125	Three sequential 3-tonne cleaning batches, sampling before pelleting and at one point post-pelleting		Pre-pelleting (first tonne milled) - 3.4 ± 0.26 Post pelleting (after 8 tonnes) - 7.2 ± 1.29	McEvoy et al., 2003

7. Another study (McEvoy et al., 2003) showed that feed batches produced after the intentional incorporation of nicarbazin into feed result in carry-over levels as high as 8.49 mg/kg in the subsequent feed. A study of German feed-production plants (n=450) showed less than 4% carry-over levels in more than half of the examined production plants (W. Strauch, 2002 from EFSA, 2008). Another survey of Belgian compound-feed production companies reported the same level of carry-over in pelleted feeds, whereas the mash feeds showed a carry-over level of less than 5% (EFSA, 2008). Studies on carry-over in feed conducted in Italy in 2015 and 2017 reported 0.1-0.8 mg/kg of nicarbazin in poultry non-medicated feed (Roila et al., 2019). In 2006, the Czech Republic reported 43.5 mg/kg of nicarbazin in one sample of non-medicated pre-mixture for pigs out of 254 samples of different feed commodities (EFSA, 2008). Data for nicarbazin residues from a 2010-2012 Italian survey of non-medicated feedstuff showed a highest carry-over level of 0.46 mg/kg (Moretti et al., 2013), whereas another survey conducted in feedstuffs from feed mills or animal farms in Italy from 2010-2017 showed nicarbazin residues as high as 1.46 mg/kg (Annunziata et al., 2018). Nicarbazin is authorized in the EU and Australia for use in broiler chickens but not approved for laying hens, so it is assumed residues in eggs are due to carry-over.

8. “The CGMP regulations require medicated feed manufacturers to use one or more of the approved cleanout procedures, such as cleaning, sequencing, and/or flushing to prevent unsafe contamination by drug carryover (Food and Drug Administration, Department of Health and Human Services, 1976). The most effective cleanout procedure is considered the thorough cleaning of the feed manufacturing equipment. However, given its time-consuming nature and the downtime needed to clean the equipment, sequencing and flushing thoroughly are the most commonly used in the feed industry.” [...] “When it comes to flushing, the FDA recommends using 50– 100 g/kg of the mixer’s total capacity as the flush material.” (Martinez et al., 2018).
9. Based on the controlled feed mill study of Martinez et al., 2018, under practical conditions (following cleaning and flushing representing GMP), a maximum nicarbazin level of 2.2 mg/kg would be expected in non-medicated feedstuff due to the unavoidable and unintended carry-over of nicarbazin in non-target animal feed.

Step 2. Estimates of anticipated residue levels in food commodities of animal origin

a) Calculating TF for egg

10. As given in **Table 4**, feeding studies with laying hens were used to assess the potential for residues to transfer from feed to egg. DNC is contained predominantly in egg yolk, whereas the HDP is found mainly in albumin (Cannavan et al., 2000, Mortier et al., 2005). DNC is the marker residue for nicarbazin. In whole egg residues were 226 µg/kg on feeding at 1 mg/kg (Oishi and Oda, 1989), 7.69 µg/kg at 0.2 mg/kg, 17.96 µg/kg at 0.4 mg/kg, 64.10 µg/kg at 1.3 mg/kg, 192.3 µg/kg at 3.8 mg/kg and 631 µg/kg at 12.1 mg/kg (Cannavan et al., 2000), 300 µg/kg at 2 mg/kg and 6500 µg/kg at 40 mg/kg (Mortier et al., 2005), 10000 µg/kg at 200 mg/kg (Nose et al 1982) and 15300 µg/kg at 147 mg/kg (Johnston et al., 2001).
11. From **Table 4**, feeding studies with laying hens only fed nicarbazin at levels close to the carry-over level of 2.2 mg/kg were used to assess the potential for veterinary drug carry-over to transfer from feed to egg (Cannavan et al., 2000 and Mortier et al., 2005). As summarized in **Table 4**, TFs for eggs are 0.051 and 0.150, so the median TF is **0.10** (the Nose et al., 1982 study was not used as issues were observed with animal health and the Oishi and Oda et al., 1989 study was excluded as it is unknown if the nicarbazin values are measured as DNC).

Table 4: Compilation of feeding studies of nicarbazin on laying poultry

Species	Feed level (mg/kg)	Duration (days)	LOD (mg/kg)	LOQ (mg/kg)	Residue monitored	Residue level in eggs (µg/kg)	TF _{egg}	Reference
Laying hens*	2	14	NS	0.001 CC α	DNC	300	0.150	Mortier et al., 2005
	40					6500	0.162	
Laying hens	200	14	NS	NS	DNC	10000	0.05 ^C	Nose et al 1982
Laying hens	1.0	10	0.010	NS	DNC	226	0.226	Oishi and Oda, 1989
	0.5					-		
	0.1					-		
	0.05					-		
Laying hens*	0.2	16	0.0003	0.001	DNC	7.69	0.038 ^D	Cannavan et al., 2000
	0.4					17.96	0.045 ^D	
	1.3					64.10	0.050 ^D	
	3.8					192.3	0.051 ^D	
	12.1					631	0.052	
Laying hens	34.9	14	0.035 ^A	0.117 ^B	DNC	4300	0.123	Johnston et al., 2001
	54.2					9400	0.173	
	92.5					13900	0.150	
	147					15300	0.104	

*Feeding studies used to calculate TFs. NS – Not Specified.

^A LOD = $3 \times S/N$ (Primus et al., 2003)

^B LOQ = $10 \times S/N$

^C laying ceased after 7 days of dosing and restarted after 12 days on non-medicated feed.

^D TFs were calculated by applying “ $Y = 0.0195x + 0.05$ equation” derived by Mortier et al., 2005

b) Calculating the anticipated veterinary drug carry-over level in egg

Option 1

12. Considering the carry-over of nicarbazin in laying hen feed at 1% and assuming a median transfer factor of 0.10, the expected nicarbazin residue level in the egg at 1% carry-over would be 125 µg/kg ($TF_{\text{egg}} \times \text{residue level in the feed} = 0.10 \times 125 \text{ mg/kg feed} \times 1\%$).
13. Cannavan et al. (2000) showed a linear relationship between nicarbazin feed intake and DNC levels in eggs that could be described by the equation below. They further demonstrated that nicarbazin levels in feed above 2 mg/kg result in DNC levels in eggs greater than the UK differential action limit (DAL) of 100 µg/kg.

$$\text{Feed-nicarbazin (mg/kg)} = 0.0195 \times \text{whole egg residue DNC (}\mu\text{g/kg)} + 0.05$$

so

$$\text{whole egg residue DNC (}\mu\text{g/kg)} = (\text{feed nicarbazin (mg/kg)} - 0.05)/0.0195$$

14. From the above equation, DNC residues in eggs of 61.5 µg/kg would be anticipated at feed carry-over of 1% of the maximum authorized level of 125 mg/kg.

Option 2

15. Based on feed mill studies under practical conditions, the maximum nicarbazin carry-over in non-medicated feed is 2.2 mg/kg (Martinez et al., 2018). Utilizing this level, the expected nicarbazin residue level in the egg would be 220 µg/kg ($TF_{\text{egg}} \times \text{residue level in the feed} = 0.10 \times 2.2 \text{ mg/kg feed}$).

Step 3. Action levels

16. Table 5 summarizes the anticipated nicarbazin residue levels in eggs calculated using the median TF and assuming a hypothetical carry-over rate of 1% (Option 1) and at the maximum concentration in the feed from feed mill studies (Option 2). The nicarbazin residue level in egg at the 1% hypothetical carry-over level would be 125 µg/kg (Option 1), and at 2.2 mg/kg of carry-over level in feed, it would be 220 µg/kg (Option 2).

Table 5: Summary of the anticipated residue levels in chicken egg

Commodity	TF	Anticipated residue level (µg/kg)	
		Option 1	Option 2
		1% (1.25 mg/kg feed)	2.2 mg/kg feed
Egg	0.10	125	220

17. In the current example, the anticipated nicarbazin residue level of 220 µg/kg was chosen as the appropriate value for human exposure assessment based on the available feed mill data (Option 2) over the conservative default carry-over level of 1%.
18. If no data demonstrate the amount of unavoidable and unintentional veterinary drug carry-over in feed occurring after mitigation steps have been performed, then a discussion is needed to determine whether CCRVDF should consider setting action levels.

Step 4. Human dietary exposure assessment

19. Noting that JECFA is the appropriate committee to perform **Step 4** (Human dietary exposure assessment), in this pilot study, dietary exposure to nicarbazin residues in food resulting from unavoidable and unintended nicarbazin carry-over in non-target animal feed was assessed using the JECFA TMDI (Theoretical Maximum Daily Intake) as a conservative approach.
20. The 2022 JECFA established an ADI of 900 µg/kg bw (DNC) based on toxicological effects (JECFA/94/SC). Based on intended use in broilers considered by the 2022 JECFA, incurred DNC residues in chicken muscle, offal, and skin with fat, at 24 hours withdrawal time and 125 mg/kg feed, the highest Global Estimates of Chronic Dietary Exposure (GECDE) for infants and toddlers estimated by the 2022 JECFA was 210 µg/kg bw per day representing 23% of the upper bound of the ADI of 900 µg/kg bw.
21. For the expected carry-over residues in eggs, a dietary exposure assessment was performed using the 220 µg/kg nicarbazin residue level in eggs, food consumption factor of 100 g of egg, and ADI value of 900 µg/kg bw/day (**Table 6**).
22. As a marker residue to total residue (M:T) ratio is unavailable for eggs, the lowest M:T ratio identified by JECFA in the target animal species (kidney – 0.25) has been used to complete the human dietary exposure assessment.

Table 6: Estimation of dietary exposure to nicarbazin (DNC) residues in chicken eggs using JECFA TMDI approach

Commodity	Daily consumption (g)	Anticipated residue level (µg/kg)	M:T	TMDI (mg)
Egg	100	220	0.25	0.088
TMDI as %ADI				0.16%

dietary exposure estimate (TMDI) = 0.088 mg ÷ 60 kg person/day

= 0.00147 mg/kg bw/day

= 0.00147 mg/kg bw/day ÷ 0.9 mg/kg bw/day × 100%

= 0.16% of the ADI

23. The dietary exposure estimates for nicarbazin residues in eggs from non-target animals represent 0.16% of the ADI. Therefore, it can be considered that there is no appreciable risk to consumers' health from the consumption of eggs produced from laying hens consuming a feed with a carry-over level of up to 2.2 mg/kg, regardless of other sources of dietary exposure.
24. Alternatively, JECFA can be requested to advise on an estimate of the appropriate M:T ratio for eggs based on the established M:T ratios in the target animal species, applying safety factors as deemed necessary.
25. In this pilot study, an action level of 0.220 mg/kg for nicarbazin in eggs from laying hens as non-target animals is proposed to accommodate the presence of nicarbazin because of unavoidable and unintended nicarbazin carry-over in animal feed (Table 7). This aligns with similar limits established by the EU and New Zealand for nicarbazin in eggs (0.220 mg/kg).


Table 7: Proposed action level for nicarbazin in chicken egg

Commodity	Proposed action level (mg/kg)	[For comparison] Maximum content (mg/kg)
Egg	0.220	0.3 (EU) 0.3 (New Zealand)
Marker residue - 4,4'-dinitrocarbanilide (DNC)		

PILOT STUDY B: ESTIMATING ACTION LEVELS FOR UNAVOIDABLE AND UNINTENTIONAL LASALOCID CARRY-OVER IN CHICKEN EGG

26. Lasalocid sodium (referred to as lasalocid hereafter) is a monocarboxylic polyether ionophore obtained via fermentation of a strain of *Streptomyces*. It is used to control coccidiosis in chickens for fattening, chickens reared for laying, turkeys, and minor avian species (EFSA, 2017). **Table 8** provides a summary of lasalocid details.
27. Laying hens are identified as the most likely non-target animals to be exposed to unavoidable and unintended carry-over of lasalocid in non-target animal feed, as feed for broiler chickens and laying hens is often prepared at the same feed mill. Survey or residue monitoring data on lasalocid in poultry eggs (**Table 9**) and feeding studies on laying hens (**Table 11**) provide evidence for detectable lasalocid levels in eggs from laying hens fed feed produced in accordance with good manufacturing practices. **Attachment 2** summarizes the residue data for lasalocid measured in edible tissues from poultry fed lasalocid-containing medicated feed.
28. Lasalocid is classified as the most potent ionophore in terms of causing unavoidable residues in eggs (Olejnik et al., 2014; Anadon and Martinez-Larranaga, 2014). This is attributed to the fact that eggs are a major route of excretion for lasalocid residues in laying hens. Therefore, even low levels of lasalocid in feed, resulting from unintentional and unavoidable carry-over, can result in the accumulation of residues at reasonably high concentrations in eggs (Wong and Roxburgh, 2010; Kennedy et al., 1996; Vandenberg et al., 2012).
29. A wide range of lasalocid levels were reported in surveys/residue monitoring of eggs (<1-3450 µg/kg) (**Table 9**). As listed in **Table 11**, feeding studies on laying hens resulted in lasalocid levels in whole eggs ranging from 6.36 – 12,000 µg/kg. The variation is likely explained by the concentration of lasalocid incorporated into feed to conduct each study, as lasalocid deposition into egg has been shown to have a nearly linear relationship with the concentration in feed (EFSA, 2007).
30. In terms of possible sources of lasalocid residues in edible commodities of chicken, carry-over of lasalocid to non-target animal feed during feed manufacturing has been identified as a source of lasalocid residues in egg (Kennedy et al., 1996; Vandenberg et al., 2012). Even if all preventative measures are followed, such as the use of rinsing batching, cross-contamination of residues is unavoidable under practical conditions (EFSA, 2007). Additionally, lasalocid is a very dusty compound, which can easily contaminate feed during the manufacturing process (Rokka et al., 2013).

Table 8: Summary of lasalocid details

Chemical Name	Lasalocid sodium
Marker Residue	Lasalocid A
Structure	 (EFSA, 2017)
Water Solubility (20 °C)	1,060 mg/L (EFSA, 2017)
log K_{ow}	2.3 (EFSA, 2017)
Target Animal	Chickens for fattening, chickens reared for laying, turkeys for fattening
Authorized maximum content in complete feed and Withholding Period (WHP)	100 mg/kg (broiler chickens and turkeys; Canada) – zero-day (meat) 75-125 mg/kg (broiler chickens and turkeys; US) – zero-day (meat) 75-125 mg/kg (turkeys; UK) – 5-day (meat) 75-100 mg/kg – (broiler chickens and replacement pullets; Australia) – 3- day (meat) and 14-day (egg) 90-125 mg/g – (turkeys; Australia) – zero-day (meat) 75-100 mg/kg – (broiler chickens and replacement pullets; New Zealand) – zero-day (meat) 14-day (egg) 90-125 mg/g – (turkeys; New Zealand) – zero-day (meat)

LOQ	5 µg/kg for all tissues
ADI	0-5 µg/kg bw/day (JECFA/78/SC, 2013)
MRLs for chicken (broilers/layers) (mg/kg)	<p>AUS fat/skin 0.6, kidney 0.7, liver 1.2, muscle 0.4, eggs 0.05</p> <p>Canada kidney 0.4, liver 0.4, muscle 0.1, skin/fat 0.4</p> <p>Codex muscle 0.4, liver 1.2, kidney 0.6, skin/fat 0.6</p> <p>EU muscle 0.06, skin/fat 0.3, liver 0.3, kidney 0.15, eggs 0.15</p> <p>UK VMD muscle 0.06, liver 0.3, kidney 0.15, skin/fat 0.3, eggs 0.1</p> <p>US skin/fat 1.2, liver 0.4</p>
Maximum content in feed for non-target species (mg/kg)	<p>EU Regulation EU 574/2011</p> <p>Feed materials - 1.25</p> <p>Compound feed for dogs, calves, rabbits, equine species, dairy animals, laying birds, turkeys (> 16 weeks), and chickens reared for laying (> 16 weeks) – 1.25</p> <p>Compound feed for chickens for fattening, chickens reared for laying (< 16 weeks), and turkeys (< 16 weeks) for the period before slaughter in which the use of lasalocid sodium is prohibited (withdrawal feed) – 1.25</p> <p>Other animal species – 3.75</p>
Maximum content in food from non-target species (mg/kg)	<p>EU Regulation EU 610/2012</p> <p>Food of animal origin from animal species other than poultry and bovine: Milk – 0.001</p> <p>Liver – 0.05</p> <p>Kidney – 0.02 Other food – 0.005</p>

Table 9: Survey or residue monitoring on lasalocid in poultry egg

Country	Year	Commodity	LOQ (mg/kg)	MRL (mg/kg)	No. of samples tested	Positive sample n>LOQ (n>LOD)	Residue levels (µg/kg)	Highest residue level (µg/kg)	Reference
Australia	2021-22	Whole egg	Limit of Reporting (LOR) – 0.01	0.05	30	1	>LOR to ≤½MRL	NS	NRS, 2021-2022
EU	2005	Egg	CCα = 0.001	MRL NS AL – 0.01 (Belgian Food Agency)	320	24	<1-49	49	Mortier et al., 2005
(Northern) Ireland	1994	Egg	0.001	-	161	107	0.3-129	129	Kennedy et al., 1996; Kennedy et al., 1998
(Northern) Ireland	1995	Egg	0.001	-	220	45 (= 220*0.205)	<1 to >10	NS	Kennedy et al., 1998
Belgium	2004	Egg	-	-	190	-	4-90	90	Mortier et al., 2005
UK	1995-2005	Egg	-	0.01	2855	138	<2-3,450	3450	EFSA, 2007
Italy	2012 2013 2014 2015 2016 2017	Egg	0.001	-	353 animal tissue and egg samples	2 - 1 4 (2) 1 -	1.5-2.0 - 1.2 2.8-1,002 26 -	1,002	Roila et al., 2019

UK	1998	Egg	-	0.002 ^a	221	5	<1-3500	3500	Wong and Roxburgh, 2010
	1999			0.002	208	21			
	2000			0.002	212	20			
	2001			0.002	222	12			
	2002			0.04	280	18			
	2003			0.05	275	33			
	2004			0.05	283	8			
	2005			0.05	294	4			
	2006			0.05	249	4			
	2007			0.15	218	0			

NS – Not Specified. AL – Action Limit.

^a Defined as action level/maximum residue limit for each specific year.

Lasalocid presence in eggs due to unavoidable and unintended lasalocid carry-over in animal feed

Step 1. Animal dietary exposure assessment

Option 1

31. A maximum approved rate of 125 mg lasalocid/kg in broiler feed is considered for proposing action levels in eggs from laying hens utilizing hypothetical carry-over rates. Carry-over of lasalocid in laying hen feed at a hypothetical level of 1% of the maximum authorized level of 125 mg lasalocid/kg of feed for broiler chickens would result in a carry-over level of lasalocid in laying hen feed of 1.25 mg/kg.

Option 2

32. Table 10 summarizes the carry-over levels of lasalocid in non-medicated animal feed during medicated feed manufacturing, as reported by Kennedy et al. (1996, 1998). In one study (Kennedy et al., 1996), a feed mill prepared a four-ton batch of medicated turkey grower meal using a powdered lasalocid drug premix to provide a final therapeutic concentration of 100 mg lasalocid/kg of feed. The level of lasalocid carry-over was assessed in nine subsequent batches of non-medicated feed. The first batch of non-medicated feed contained 6 mg lasalocid/kg of feed, while the ninth batch of non-medicated feed contained lasalocid concentrations ranging from 0.5-1 mg/kg. In an analogous experiment, cross-contamination of lasalocid from a medicated premix to successive batches of non-medicated premix was measured, with only the first batch of non-medicated premix containing appreciable levels of lasalocid (level not reported) (Kennedy et al., 1996).
33. In a second study conducted by Kennedy et al. (1998), the carry-over of lasalocid into non-medicated feed was evaluated following the manufacturing of medicated feed (100 mg lasalocid/kg of feed) using a granular lasalocid premix. Lasalocid remained present in the first four batches of non-medicated feed but could not be detected in the subsequent five batches. When compared to the results of the study conducted by Kennedy et al. in 1996, the extent of carry-over into non-medicated feed was significantly reduced when medicated feed was manufactured using the granular premix.
34. In both studies described above, there was no indication of any mitigation steps representing good manufacturing practices being performed.

Table 10: Carry-over levels of lasalocid in non-medicated animal feed during medicated feed manufacturing

Level in medicated feed (mg/kg)	Flushing procedure	Level in flush (mg/kg)	Level in non-medicated diet (mg/kg)	Reference
100	-	-	Batch one – 6 Batch nine – 0.5-1	Kennedy et al., 1996
100	-	-	Batch 1 – 3.2 Batch 4 – 0.25	Kennedy et al., 1998

35. A study of German feed-production plants (n=450) showed less than 4% carry-over levels in more than half of the examined production plants (W. Strauch, 2002 from EFSA, 2007). Another survey of Belgian compound-feed production companies reported the same level of carry-over in pelleted feeds, whereas the mash feeds showed a carry-over level of less than 5% (EFSA, 2007). Studies on carry-over in feed conducted in Italy from 2012 to 2017 reported lasalocid levels in non-medicated poultry feed between 0.1-5.9 mg/kg (Roila et al., 2019). A Swiss feed plant that produces feed for chickens for fattening and laying hen meal found that one production passage without the addition of lasalocid was not sufficient to reduce the contents of the drug in laying hen meal below 30 µg/kg (Noser et al., 2006).
36. In 2007, following a request from the European Commission, the Panel on Contaminants in the Food Chain was asked to deliver a scientific opinion on cross-contamination of non-target feedingstuffs by lasalocid authorized for use as a feed additive. The Czech Republic analyzed 254 samples from feed materials for non-target animal species in 2006. One positive sample, a complete feeding stuff for pigs, contained 8.41 mg lasalocid/kg feed (EFSA, 2007). Denmark reported the analyses of 111 feed samples for non-target species collected between 2004 and 2007 and found one positive sample containing 0.26 mg lasalocid/kg feed (EFSA, 2007). Information from the Rapid Alert System for Food and Feed (RASFF) between April 2002 and April 2006 showed nine incidences where feed for non-target animal species contained lasalocid. The amounts detected were between 0.003 and 12.07 mg lasalocid/kg feed, with one outlier containing 64.6 mg/kg feed. The outlier is most likely due to accidental contamination (EFSA, 2007).

37. Based on the feed mill study of Kennedy et al., 1998, a maximum lasalocid level of 3.2 mg/kg may be expected in non-medicated feedstuff. However, whether mitigation steps representing GMP were performed during this study is unclear. Indeed, “the CGMP regulations require medicated feed manufacturers to use one or more of the approved cleanout procedures, such as cleaning, sequencing, and/or flushing to prevent unsafe contamination by drug carryover (Food and Drug Administration, Department of Health and Human Services, 1976). The most effective cleanout procedure is considered the thorough cleaning of the feed manufacturing equipment. However, given its time-consuming nature and the downtime needed to clean the equipment, sequencing and flushing thoroughly are the most commonly used in the feed industry.” [...] “When it comes to flushing, the FDA recommends using 50–100 g/kg of the mixer’s total capacity as the flush material.” (Martinez et al., 2018). Consequently, the anticipated exposure level for non-target animals could not be estimated.

Step 2. Estimates of anticipated residue levels in food of animal origin

a) Calculating TF for egg

38. As given in **Table 11**, feeding studies with laying hens were used to assess the potential for residues to transfer from feed to egg. Lasalocid A is the marker residue for lasalocid. For the study performed by Kennedy et al. (1996), residues in whole eggs were calculated using the following equation:

$$\text{Concentration in eggs (ug/kg)} = 63.6 \times \text{Concentration in feed (mg of lasalocid/kg of feed)}$$

39. In whole egg, residue concentrations were 6.36 µg/kg on feeding at 0.1 mg/kg, 19.08 µg/kg at 0.3 mg/kg, 31.8 µg/kg at 0.5 mg/kg, 63.6 µg/kg at 1.0 mg/kg, and 318 µg/kg at 5.0 mg/kg (Kennedy et al., 1996), 370 µg/kg at 3.125 mg/kg, 780 µg/kg at 6.25 mg/kg, and 1410 µg/kg at 12.5 mg/kg (Vandenberge et al., 2012), 11000-12000 µg/kg at 125 mg/kg (EMA, 2006) and 2.0 µg/kg at 1 mg/kg (Rokka et al., 2005).
40. As suitable data is not available to establish, with certainty, that unintended and unavoidable carry-over of lasalocid would occur at a level higher than 1% of the maximum authorized dose (125 mg/kg), a default hypothetical carry-over of 1% (1.25 mg/kg) was considered. From **Table 11**, feeding studies with laying hens only fed lasalocid at levels close to the carry-over level of 1.25 mg/kg were used to assess the potential for veterinary drug carry-over to transfer from feed to egg (Kennedy et al., 1996 and Vandenberge et al., 2012).
41. As summarized in **Table 11**, the TFs for eggs are 0.0636 and 0.118, resulting in a median TF of 0.0908. The TF determined by Rokka et al. (2005) was excluded as it did not state the concentration of lasalocid in the whole egg but only in the egg yolk. The EMA (2006) study was excluded as the laying hens were fed lasalocid at a therapeutic concentration. Therefore, the TF may be overestimated.

Table 11: Compilation of feeding studies of lasalocid on laying poultry

Species	Feed level (mg/kg)	Duration (days)	LOD (µg/kg)	LOQ (µg/kg)	Residue monitored	Residue levels in eggs (µg/kg)	TF _{egg}	Reference
Laying hens ^{*,a}	0.1 0.3 0.5 1.0 5.0	16	0.3	1.0	Lasalocid A	6.36 19.08 31.8 63.6 318	0.0636	Kennedy et al., 1996
Laying hens ^{*,a}	3.125 6.25 12.5	14d + 17d depletion period	-	2	Lasalocid A	370 (d16) 780 (d11) 1410 (d14)	0.118 0.125 0.113	Vandenberge et al., 2012
Laying hens	125	12	0.093	10	Lasalocid A	11000-12000 (whole egg; total residues) 291 (egg white)	0.088- 0.096	EMA, 2006

						32500 (egg yolk)		
Laying hens	1.0	21	CCB – 2.0		Lasalocid A	2.0 (egg yolk)	0.002	Rokka et al., 2005
* The peak residue concentration was used to determine the residue levels in eggs.								
^a Feeding studies used to calculate TFs.								
NS – Not Specified.								

b) Calculating the anticipated veterinary drug carry-over level in egg

Option 1

42. Considering the carry-over of lasalocid in laying hens feed at 1% and assuming a mean transfer factor of 0.0908, the expected lasalocid residue level in the egg would be 113.5 µg/kg ($TF_{\text{egg}} \times \text{residue level in the feed} = 0.0908 \times 125 \text{ mg/kg feed} \times 1\%$).

Option 2

43. The anticipated exposure level for non-target animals could not be estimated based on the feed mill study of Kennedy et al., 1998, as GMP could not be verified.

Step 3. Action levels

44. Table 12 summarizes the anticipated lasalocid residue level in eggs calculated using the median TF and assuming a hypothetical carry-over rate of 1% (Option 1). The lasalocid residue level in the egg at the 1% hypothetical carry-over level would be 113.5 µg/kg (**Option 1**).

Table 12: Summary of the anticipated residue levels in chicken egg

Commodity	TF	Anticipated residue level (µg/kg)	
		Option 1 1% (1.25 mg/kg feed)	Option 2
Egg	0.0908	113.5	-

45. In the current example, an anticipated lasalocid residue level of 113.5 µg/kg, based on a default hypothetical carry-over of 1%, was chosen as the appropriate value for the human exposure assessment (**Option 1**). A default carry-over level of 1% was considered as no data demonstrates the amount of unavoidable and unintentional veterinary drug carry-over in feed after mitigation steps have been performed.
46. *If no data demonstrate the amount of unavoidable and unintentional veterinary drug carry-over in feed occurring after mitigation steps have been performed, then a discussion is needed to determine whether CCRVDF should consider setting action levels.*

Step 4. Human dietary exposure assessment

47. Noting that JECFA is the appropriate committee to perform **Step 4** (Human Dietary Exposure Assessment), in this pilot study, dietary exposure to lasalocid residues in food resulting from unavoidable and unintended lasalocid carry-over in non-target animal feed was assessed using the JECFA TMDI (Theoretical Maximum Daily Intake) as a conservative approach.
48. The 2013 JECFA established an ADI of 0-5 µg/kg bw for lasalocid based on a NOAEL of 0.5 mg/kg body weight per day based on a toxicity study in rabbits (JECFA/78/SC, 2013). The estimated dietary exposure was calculated as 80 µg/person per day, based on median residues for a 1-day withdrawal in chicken, which represents approximately 27% of the upper bound of the ADI, based on a 60 kg individual (JECFA/78/SC, 2013).

49. "The Global Estimates of Chronic Dietary Exposure (GECDE) for the general population is 1.85 µg/kg bw per day, which represents 37% of the upper bound of the ADI. The GECDE for children is 3.38 µg/kg per day, which represents 67% of the upper bound of the ADI. The GECDE for infants is 2.99 µg/kg bw per day, which represents 60% of the upper bound of the ADI." (FAO/WHO, 2015).
50. For the expected carry-over of residues in eggs, a dietary exposure assessment was performed using the 113.5 µg/kg lasalocid A residue level in eggs, food consumption factor of 100 g of egg, marker residue to total residue (M:T) ratio of 0.38 (EMA, 2006) and ADI value of 5 µg/kg bw/day (**Table 13**).

Table 13: Estimation of dietary exposure to total lasalocid residues in chicken eggs using JECFA TMDI approach

Commodity	Daily consumption (g)	Anticipated residue level (µg/kg)	M:T	TMDI (mg)
Egg	100	113.5	0.38	0.03
TMDI as % ADI				10 %

Dietary exposure estimate (TMDI) = 0.03 mg ÷ 60 kg person/day

= 0.0005 mg/kg bw/day

= 0.0005 mg/kg bw/day ÷ 0.005 mg/kg bw/day x 100%

= 10 % of the ADI

51. The dietary exposure estimates for lasalocid residues in eggs from non-target animals represent 10% of the ADI. Therefore, it can be considered that there is no appreciable risk to consumers' health from the consumption of eggs produced from laying hens consuming a feed with a carry-over level of up to 1.25 mg lasalocid/kg, regardless of other sources of dietary exposure.
52. In this pilot study, it is proposed to establish an action level of 0.1 mg/kg (rounded down from 0.113 mg/kg) for lasalocid in eggs from laying hens as non-target animals to accommodate the presence of lasalocid because of unavoidable and unintended lasalocid carry-over in animal feed (**Table 14**).

Table 14: Proposed action level for lasalocid in chicken egg

Commodity	Proposed action level (mg/kg)	[For comparison] Maximum content (mg/kg)
Egg	0.1	0.15 (MRL - EU) 0.05 (MRL - Australia)
Marker residue – Lasalocid A		

ANNEX**National and Regional Guidelines Canada**

1. Medication Sequencing Guideline for Management of Drug Carryover (<http://inspection.gc.ca/animals/feeds/inspection-program/medication-sequencing/eng/1389362488069/1389362490053>)
2. Validation studies for Modification of Sequencing Guidelines (<http://inspection.gc.ca/animals/feeds/inspection-program/sequencing-guidelines/eng/1373325944197/1373325944713>)
3. Measurement of Feed Carryover Level (<http://inspection.gc.ca/animals/feeds/inspection-program/measurement-of-feed/eng/1373325386112/1373325437132>)
4. Medication Residues Validation Testing Procedures for Equipment Cleanout Procedures (<http://inspection.gc.ca/animals/feeds/inspection-program/equipment-cleanout/eng/1373325971995/1373325972541>)

European Union

1. European Commission. 2011. Regulation (EC) No 574/2011 of 16 June 2011 amending Annex I to Directive 2002/32/EC of the European Parliament and of the Council as regards maximum levels for nitrite, melamine, Ambrosia spp. and carryover of coccidiostats or histomonostats in non-target feed. Off J Eur Commun. L40:19–25.
2. European Commission. 2012. Regulation (EC) No 124/2009 of 10 February 2009 setting maximum levels for the presence of coccidiostats or histomonostats in food resulting from the unavoidable carry-over of these substances in non-target feed. Off J Eur Commun. L178:1-3.
3. COUNCIL DIRECTIVE (90/167/EEC) of 26 March 1990 laying down the conditions governing the preparation, placing on the market, and use of medicated feedingstuffs in the Community.
4. DIRECTIVE 2002/32/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL
5. of 7 May 2002 on undesirable substances in animal feed (<http://eur-lex.europa.eu/legal-content/DE/TXT/HTML/?uri=CELEX:32002L0032&from=DE>)
6. COMMISSION DIRECTIVE 2009/8/EC of 10 February 2009 amending Annex I to Directive 2002/32/EC of the European Parliament and of the Council as regards maximum levels of unavoidable carry-over of coccidiostats or histomonostats in nontarget feed [LexUriServ.do \(europa.eu\)](http://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32009L0008&from=DE)
7. COMMISSION REGULATION (EC) No 124/2009 of 10 February 2009 setting maximum levels for the presence of coccidiostats or histomonostats in food resulting from the unavoidable carry-over of these substances in non-target feed.
8. Document EUR-Lex - 52014PC0556
Proposal for a REGULATION OF THE EUROPEAN PARLIAMENT AND THE COUNCIL on the manufacture, placing on the market, and use of medicated feed and repealing Council Directive 90/167/EEC (<http://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:52014PC0556&qid=1444290152009&from=DE>)
9. Feed Hygiene Regulation (EC) No 1831/2003: Annex II provides, among others, that “Technical or organizational measures must be taken to avoid or minimize, as necessary, any cross-contamination and errors.”
10. European Feed Manufacturers Guide for Good Hygiene Practice for the Manufacturing of Feed for Food-Producing Animals (EFMC): This guide is meant to help operators meet the requirements of the EU Feed Hygiene Regulation. It includes provisions for the prevention and minimization of carry-over, including guidance for measuring premises-bound carry-over. It also includes definitions of carry-over and cross-contamination.

References

1. ANADA 200-027. Abbreviated New Animal Drug Applications (ANADAs). FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. NICARMIX (nicarbazine) 25%. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/960>
2. Annunziata L., Visciano P., Stramenga A., Colagrande M. N., Campana G., Scortichini G., Migliorati G., Compagnone D. 2018. Investigation of Nonionophoric Coccidiostat Residues in Feed as a Consequence of Carryover. *J Food Prot*, 23:482-489. <https://doi.org/10.4315/0362-028X.JFP-17-414>

3. Bilandžić N., Dolenc J., Gačnik K.Š., Varenina I., Kolanović B.S. 2013. Feed additives diclazuril and nicarbazin in egg and liver samples from Croatian farms. *Food Addit Contam Part B Surveill*, 6(2):90-7. <https://doi.org/10.1080/19393210.2012.727474>
4. Cannavan A., Kennedy D.G., 2000. Possible causes of nicarbazin residues in chicken tissues. *Food Addit Contam*, 17:1001–1006 <https://doi.org/10.1080/0265203001001325>
5. Cannavan A., Ball G., Kennedy D.G. 2000. Nicarbazin contamination in feeds as a cause of residues in eggs. *Food Addit Contam*, 17:829-836. <https://doi.org/10.1080/026520300420394>
6. Codex Alimentarius Code of Practice on Good Animal Feeding (CAC/RCP 54-2004) <https://www.fao.org/fao-who-codexalimentarius/thematic-areas/animal-feed>
7. Codex Alimentarius Guidelines on the Application of Risk Assessment for Feed (CAC/GL 80-2013). Codex Alimentarius Commission. <https://www.fao.org/feed-safety/resources/resources-details/en/c/1054051/>
8. Danaher M., Campbell K., O'Keeffe M., Capurro E., Kennedy G., Elliott C.T. 2008. Survey of the anticoccidial feed additive nicarbazin (as dinitrocarbanilide residues) in poultry and eggs. *Food Addit Contam*. 25:32-40. <https://doi.org/10.1080/02652030701552956>
9. EFSA (European Food Safety Authority), 2003. Opinion of the Scientific Panel on Additives and Products or Substances used in Animal Feed on the request from the Commission on the efficacy and safety of the coccidiostat Koffogran. The EFSA Journal, 16, 1-40. <https://doi.org/10.2903/j.efsa.2004.16>
10. EFSA (European Food Safety Authority), 2008. Cross-contamination of non-target feedingstuffs by nicarbazin authorised for use as a feed additive. Scientific opinion of the Panel on Contaminants in the Food Chain. EFSA J. 690, 1–34. <https://doi.org/10.2903/j.efsa.2008.690>
11. EFSA (European Food Safety Authority), 2010a. Scientific Opinion on the safety and efficacy of Koffogran (nicarbazin) as a feed additive for chickens for fattening. EFSA Journal, 8(3):1551. <https://doi.org/10.2903/j.efsa.2010.1551>
12. EFSA (European Food Safety Authority), 2010b. Scientific opinion on the safety and efficacy of Maxiban G160 (narsin and nicarbazin) for chickens for fattening. EFSA Panel on Additives and Products or Substances used in Animal Feed). EFSA J. 8 (4), 1754. <https://doi.org/10.2903/j.efsa.2010.1574>
13. EFSA (European Food Safety Authority), 2017. Safety and efficacy of Monimax® (monensin sodium and nicarbazin) for turkeys for fattening. EFSA Journal 2017;15(12):5094. <https://doi.org/10.2903/j.efsa.2017.5094>
14. EFSA (European Food Safety Authority), 2018a. Safety and efficacy of Coxar® (nicarbazin) for turkeys for fattening. EFSA Journal 2018;16(4):5214.
15. EFSA (European Food Safety Authority), 2018b. Safety and efficacy of Monimax® (monensin sodium and nicarbazin) for chickens for fattening and chickens reared for laying. EFSA Journal 2018;16(11):5459. <https://doi.org/10.2903/j.efsa.2018.5214>
16. EFSA (European Food Safety Authority), 2021. Safety and efficacy of a feed additive consisting of lasalocid A sodium and nicarbazin (Nilablend™ 200G) for chickens for fattening (Zoetis Belgium SA). EFSA Journal 2021;19(3):6466. <https://doi.org/10.2903/j.efsa.2021.6466>
17. FAO/WHO (Food and Agriculture Organisation/World Health Organization), 1999. Evaluation of certain veterinary drug residues in food. Fiftieth report of the Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series 888. http://whqlibdoc.who.int/trs/WHO_TRS_888.pdf.
18. FAO/WHO EXPERT COMMITTEE ON FOOD ADDITIVES Ninety-fourth meeting (Residues of veterinary drugs) Virtual meeting, 16–27 May 2022 (JECFA/94/SC)
19. Johnston J. J., Britton W.M., MacDonald A., Primus T. M., Goodal M. J., Yoder C. A., Miller L. A., Fagerstone K. A. 2001. Quantification of plasma and egg 4, 4'-dinitrocarbanilide (DNC) residues for the efficient development of a nicarbazin based contraceptive for pest waterfowl. *Pest Manage Sci*, 58, 197–202.
20. Kan C. A., Keukens H. J., Boers E. (1995) Nicarbazin residues in broiler muscle, liver and litter/faeces: early exposure or recirculation. 12th European Symposium on the Quality of Poultry Meat 1: 481-485.
21. Konieczka P., 2.31 - Validation and Regulatory Issues for Sample Preparation, Editor(s): Janusz Pawliszyn, Comprehensive Sampling and Sample Preparation, Academic Press, 2012, Pages 699-711, ISBN 9780123813749, <https://doi.org/10.1016/B978-0-12-381373-2.00064-8>.
22. Kramer H. T. 1990. Determination of Nicarbazin in Chicken Tissues by High Performance Liquid Chromatography. Report by Hazelton Laboratoories for Planalquimica Industrial, LTDA Sao Paulo, Brazil.

23. Martinez A., Jones C. K., Stark C. R., McKinney L. J., Behnke K. C., Paulk C. B. 2018. Evaluating flushing procedures to prevent nicarbazin carryover during medicated feed manufacturing, *Animal Feed Science and Technology*, 242:1-7, ISSN 0377-8401, <https://doi.org/10.1016/j.anifeedsci.2018.04.017>.
24. Martinez-Kawas A. 2008. Evaluating Flushing Procedures to Prevent Drug Carryover during Medicated Feed Manufacturing, Master Science Thesis, Universidad Autonoma de Nuevo Leon,
25. McEvoy J.D., Smyth W.G., Kennedy D.G. 2003. Contamination of animal feedingstuff with nicarbazin: investigations in a feed mill. *Food Addit Contam.* 20: 136-40. <https://doi.org/10.1080/0265203021000050608>
26. MAPA 2016a. Ministério da Agricultura, Pecuária e Abastecimento. 2016. Instrução Normativa Nº 14: anexo II: critérios para manipulação de medicamento veterinário em fábricas de produtos destinados à alimentação animal. [Ministry of Agriculture, Livestock and Supply (MAPA). Normative Instruction No. 14: annex II: criteria for handling of veterinary medicinal products in factories of products intended for animal feed]. [accessed 2016 Jul 08]. http://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/21288772/do1-2016-07-08-instrucao-normativa-n-14-de-6-de-julho-de-2016-21288711.
27. MAPA 2016b. Ministério da agricultura pecuária e abastecimento. 2016b. Instrução Normativa Nº 14: anexo II: critérios para manipulação de medicamento veterinário em fábricas de produtos destinados à alimentação animal. Brasília, 15 de julho de 2016 [Normative Instruction No. 14: Annex II: criteria for handling of veterinary medicine in factories of products intended for animal feed. Brasília, July 15, 2016].
28. Mortier L., Huet A. C., Charlier C., Daeselaire E., Delahaut P., Van Peteghem , C. 2005a. Incidence of residues of nine anticoccidials in eggs. *Food Addit Contam* 22: 1120-1125. <https://doi.org/10.1080/02652030500199355>
29. Mortier L., Huet A. C., Daeseleire E., Huyghebaert G., Fodey T., Elliott C., Delehaut P. and Van Peteghem C. 2005b. Deposition and depletion of five anticoccidials in eggs. *J Agric Food Chem.* 53: 7142-7149. <https://doi.org/10.1021/jf050836r>
30. Moretti S., Fioroni L., Giusepponi D., Pettinacci L., Saluti G., Galarini R. 2013. Development and validation of a multiresidue liquid chromatography/tandem mass spectrometry method for 11 coccidiostats in feed. *J AOAC Int*, 96:1245-1257. <https://doi.org/10.5740/jaoacint.12-440>
31. NADA 138-952a. New Animal Drug Application (NADA). FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. 1989. MAXIBAN® Narasin and Nicarbazin Type A Article. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/439>
32. NADA 138-952b. New Animal Drug Application (NADA). FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. 2018. Maxiban™ 72 narasin and nicarbazin Type A medicated article Broiler chickens. Residue Depletion of Nicarbazin and Narasin in Edible Tissues from Chickens Following Administration of Maxiban®2G160 via Feed (Study No. T4HAUK0703). <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/3875>
33. NADA 141-124. New Animal Drug Application (NADA). FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary, 2002. MAXIBAN® + BMD®(narasin and nicarbazin + bacitracin methylene disalicylate). <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/650>
34. NADA 141-113. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Combined use of MAXIBAN® and 3-NITRO® in Chicken Feeds. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/644>
Note: This product is no longer approved for use in the United States.
35. NADA 141-112. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Combined use of MAXIBAN®, BMD®, and 3-NITRO® in Broiler Chicken Feeds. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/643> Note: This product is no longer approved for use in the United States.
36. NADA 140-926. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Combined use of MAXIBAN® and BMD® in Chicken Feeds. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/489>
37. NADA 140-339. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Nicarbazin (NICARB®) plus Bambermycins (FLAVOMYCIN®) <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/478>
38. NADA 140-947. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and

- Drug Administration), Freedom of Information Summary, 1998. Maxiban®, Lincomix®. Experiment No. AAC8814. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/2501> Note: This product is no longer approved for use in the United States.
39. NADA 141-529. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary, 2020. Pennitracin MD® and Maxiban™. Study No. AAC8721. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/9748>
40. Nose N., Hoshino Y., Kikuchi Y., Masaki H., Horie S., Kawauchi S.(1982) Residues of synthetic antibacterial feed additives in tissues and eggs of chickens. *Shokuhin Eiseigaku Zasshi*, 23, 246–252 O'Keeffe M., Capurro E., Danaher M., Campbell K., Elliott C. T. 2007. Investigation of the causes for the occurrence of residues of the anticoccidial feed additive nicarbazin in commercial poultry, *Food Addit Contam*, 24:9, 923-934, <https://doi.org/10.1080/02652030701258778>
41. Oishi Y., Oda T. 1989. Studies on residues of nicarbazin in eggs of laying hens. *Shokuhin Eiseigaku Zasshi*. Food Hygiene and Safety Science) 30: 542-547.
42. Penz A. M. Jr., Vieira S. L., Ludke J. V. 1999. Nicarbazin residues in broiler tissue and litter. *J Poult Sci*, 8, 292-297 <https://doi.org/10.1093/japr/8.3.292>
43. Pietruk K., Olejnik M., Jedziniak P., Szprengier-Juszkiewicz T. 2015. Determination of fifteen coccidiostats in feed at carry-over levels using liquid chromatography-mass spectrometry. *J Pharm Biomed Anal*, 112:50-59. <https://doi.org/10.1016/j.jpba.2015.03.019>
44. van Rhijn H., Yedema E., Beek W., Kan C. A. (Kees) & Sebek, Leon & Keukens, H. (2000). Investigation into the excretion of nicarbazin in cows milk after administration of feeds containing low-levels of nicarbazin. Proceedings of the EuroResidue IV Conference, Veldhoven, The Netherlands.
45. Roila R., Branciaro R., Pecorelli I., Cristofani E., Carloni C., Ranucci D., Fioroni L. 2019. Occurrence and Residue Concentration of Coccidiostats in Feed and Food of Animal Origin; Human Exposure Assessment. *Foods*, 8(10):477 <https://doi.org/10.3390/foods8100477>
46. Strauch W. 2002. Causes and control of carry-over and cross-contamination. *Kraftfutter/Feed Magazine* 85 (4): 151-159 (Part 1). *Kraftfutter/Feed Magazine* 85 (6): 239-249 (Part 2).
47. Tarbin J. A., Bygrave J., Bigwood T., Hardy D., Rose M., Sharman M. 2005. The effect of cooking on veterinary drug residues in food: Nicarbazin (Dinitrocarbanilide component). *Food Addit Contam*, 22(11), 1126–1131. <http://dx.doi.org/10.1080/02652030500357193>.
48. UK-VMD (United Kingdom-Veterinary Medicines Directorate), 1995-2004. The Veterinary Medicines Directorate Annual Reports on Surveillance for Veterinary Residues for 1995-2004. Published by the Veterinary Medicines Directorate, New Haw, Surrey, England. <http://www.vmd.gov.uk/Publications/AnnReps/AnnReps.htm>
49. UK 2007 Annual report on surveillance for veterinary residues in food in the UK 2007 <https://www.gov.uk/government/publications/annual-report-on-surveillance-for-veterinary-residues-in-food-in-the-uk-2007>
50. Wood J. S., Downing G. G. 1980. Modified Pulse Polarographic Determination of Nicarbazin in Chicken Tissue at the 0.1 Part-per-Million Level, *J. Agric and Food Chem*, 28:452-454
51. Anadon, A., Martinez-Larranaga, M.R. 2014. Veterinary drugs residues: coccidiostats. *Encyclopedia of Food Safety*, 3:63-75.
52. Australia New Zealand Food Standards Code – Schedule 20 – Maximum residue limits. [Australia New Zealand Food Standards Code – Schedule 20 – Maximum residue limits \(legislation.gov.au\)](https://www.foodstandards.gov.au/australia-new-zealand-food-standards-code-schedule-20-maximum-residue-limits)
53. Bello, A., Henri, J., Viel, A., Mochel, J. P., Pozniak, B. 2023. Ionophore coccidiostats – disposition kinetics in laing hens and residues transfer to eggs. *Poultry Science*, 102(1):102280.
54. Codex Alimentarius Maximum Residue Limits (MRLs) and Risk Management Recommendations (RMRs) for Residues of Veterinary Drugs in Foods (CXM 2-2021). [MAXIMUM RESIDUE LIMITS \(MRLs\) AND RISK MANAGEMENT RECOMMENDATIONS \(RMRs\) \(fao.org\)](https://www.fao.org/codexalimentarius/cxm-2-2021)
55. Codex Alimentarius Veterinary Drug Details Lasalocid Sodium. [Veterinary Drug Detail | CODEXALIMENTARIUS FAO-WHO](https://www.fao.org/codexalimentarius/veterinary-drug-detail-lasalocid-sodium)
56. Dorne, J.L.C.M., Fernandez-Cruz, M.L., Bertelsen, U., Renshaw, D.W., Peltonen, K., Anadon, A., Feil, A., Sanders, P., Wester, P., Fink-Gremmels, J. 2013. Risk assessment of coccidiostats during feed cross-contamination: animal and human health aspects. *Toxicology and Applied Pharmacology*, 270(3):196-208.

57. EFSA (European Food Safety Authority). 2004. Opinion of the Scientific Panel on additives and products or substances used in animal feed (FEEDAP) on the re-evaluation of coccidiostat Avatec in accordance with article 9G of Council Directive 70/524/EEC.
58. EFSA (European Food Safety Authority). 2007. Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the European Commission on Cross-contamination of non-target feedingstuffs by lasalocid authorized for use as a feed additive. [Cross-contamination of non-target feedingstuffs by lasalocid authorised for use as a feed additive - Scientific Opinion of the Panel on Contaminants in the food Chain \(wiley.com\)](#)
59. EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed (FEEDAP), Rychen, G., Aquilina, G., Azimonti, G., Bampidis, V., Bastos, M.L., Bories, G., Chesson, A., Cocconcelli, P.S., Flachowsky, G., Kolar, B., Kouba, M., Lopez-Alonso, M., Lopez Puente, S., Mantovani, A., Mayo, B., Ramos, F., Saarela, M., Villa, R.E., Wallace, R.J., Wester, P., Brantom, P., Halle, I., van Beelen, P., Holczknecht, O., Vettori, M.V., Gropp, J. 2017. Scientific Opinion on the safety and efficacy of Avatec® 150G (lasalocid A sodium) for chickens for fattening and chickens reared for laying, and modification of the terms of authorization for chickens for fattening, chickens reared for laying, turkeys for fattening, minor avian species (pheasants, guinea fowl, quails and partridges) except laying birds. EFSA J. 2017; 15(8):4857, 32 pp. <https://doi.org/10.2903/j.efsa.2017.4857>
60. EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Bampidis, V., Azimonti, G., Bastos, M.L., Christensen, H., Dusemund, B., Fasmon Durjava, M., Kouba, M., Lopez-Alonso, M., Lopez Puente, S., Marcon, F., Mayo, B., Pechova, A., Petkova, M., Ramos, F., Sanz, Y., Villa, R.E., Woutersen, R., Gropp, J., Rychen, G., Anguita, M., Holczknecht, O., Innocenti, M.L., Navarro-Villa, A., Rossi, B., Vettori, M.V. 2022. Scientific Opinion on the safety and efficacy of a feed additive consisting of lasalocid A sodium (Avatec® 150G) for chickens for fattening and chickens reared for laying (Zoetis Belgium SA). EFSA J. 2022; 20(12):7715, 14 pp. <https://doi.org/10.2903/j.efsa.2022.7715>
61. EMA (European Medicines Agency), 2004. Lasalocid Sodium. Summary Report of the Committee for Veterinary Medicinal Products. EMEA/MRL/912/04-FINAL, October 2004. https://www.ema.europa.eu/en/documents/mrl-report/lasalocid-sodium-summary-report-committee-veterinary-medicinal-products_en.pdf
62. EMA (European Medicines Agency), 2006. Lasalocid Sodium (extension to eggs). Summary Report of the Committee for Medicinal Products for Veterinary Use. EMEA/CVMP/46049/2006-FINAL-corr, July 2006. https://www.ema.europa.eu/en/documents/mrl-report/lasalocid-sodium-extension-eggs-summary-report-committee-veterinary-medicinal-products_en.pdf
63. EMA (European Medicines Agency), 2012. Lasalocid (bovine species). European public MRL assessment report (EPMAR). EMEA/CVMP/504089/2010. https://www.ema.europa.eu/en/documents/mrl-report/lasalocid-bovine-species-european-public-mrl-assessment-report-epmar-cvmp_en.pdf
64. FAO (Food and Agriculture Organization of the United Nations), 2015. Online Edition: Residues of some veterinary drugs in foods and animals. [Food safety and quality: Details \(fao.org\)](#)
65. FAO/WHO (Food and Agriculture Organization/World Health Organization), 2015. Evaluation of certain veterinary drug residues in food. Eighty-first report of the Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series 997, Rome 17-25 November 2015. https://apps.who.int/iris/bitstream/handle/10665/204670/9789240695504_eng.pdf?sequence=1
66. FDA (Food and Drug Administration) CFR – Code of Federal Regulations Title 21 Volume 6 Part 556 Tolerances for Residues of New Animal Drugs in Food (21CFR556) [CFR - Code of Federal Regulations Title 21 \(fda.gov\)](#)
67. Government of Canada. List of Maximum Residue Limits (MRLs) for Veterinary Drugs in Foods. August 26, 2022. [List of Maximum Residue Limits \(MRLs\) for Veterinary Drugs in Foods - Canada.ca](#)
68. JOINT FAO/WHO EXPERT COMMITTEE ON FOOD ADDITIVES Seventy-eighth meeting (Residues of veterinary drugs) Geneva, 5-14 November 2013 (JECFA/78/SC)
69. Kennedy, D.G., Blanchflower, W.J., O’Dornan, B.C. 1995. Development of an ELISA for lasalocid and depletion kinetics of lasalocid residues in poultry. Food Addit. Contam., 12:83-92.
70. Kennedy, D.G., Blanchflower, W.J., Hughes, P.J., McCaughey, W.J. 1996. The incidence and cause of lasalocid residues in eggs in Northern Ireland. Food Addit. Contam., 13(7):787-794.
71. Kennedy, D.G., Hughes, P.J., Blanchflower, W.J. 1998. Ionophore residues in eggs in Northern Ireland: incidence and cause. Food Additives and Contaminants, 15(5):535-541.
72. Martinez A., Jones C. K., Stark C. R., McKinney L. J., Behnke K. C., Paulk C. B. 2018. Evaluating flushing procedures to prevent nicarbazine carryover during medicated feed manufacturing, Animal Feed Science and Technology, 242:1-7,

ISSN 0377-8401. <https://doi.org/10.1016/j.anifeedsci.2018.04.017>

73. Mortier, L., Huet, A.C., Charlier, C., Daeseleire, E., Delahaut, P., Van Peteghem, C. 2005. Incidence of residues of nine anticoccidials in eggs. *Food Additives and Contaminants*, 22(11):1120-1125.
74. NADA 096-298. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. 1995. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/342>
75. NADA 107-996. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. 2002. Lasalocid (AVATEC®) plus Bacitracin methylene disalicylate (BMD®). <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/362>
76. NADA 141-109. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Combined use of AVATEC® and BACIFERM® in turkey feeds. <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/638>
77. NADA 141-488. New Animal Drug Application (NADA), FDA-HHS (Department of Health and Human Services, Food and Drug Administration), Freedom of Information Summary. Lasalocid type A medicated article and lincomycin type A medicated article. 2018 <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/3944>
78. National Residue Survey (NRS), Department of Agriculture, Fisheries, and Forestry. Australian Government. [eggs-residue-testing-annual-datasets-2021-22 \(agriculture.gov.au\)](https://www.agriculture.gov.au/eggs-residue-testing-annual-datasets-2021-22)
79. Noser, J., Wenk, P., Sutter, A., Schneider, P. 2006. Kokzidiostatika in Eiern, Geflügel-fleisch under Futtermitteln: Eine Stichproben-Kontrolle über den Nordwestschweizer Markt. *Mitt Lebensm Hyg* 97:107-120.
80. Olejnik, M., Szprengier-Juskiewicz, T., Jedziniak, P. 2014. Distribution of semduramicin in hen eggs and tissues after administration of cross-contaminated feed. *Food Additives and Contaminants: Part A*, 31(8):1393-1398.
81. Roila, R., Branciarri, R., Pecorelli, I., Cristofani, E., Carloni, C., Ranucci, D., Fioroni, L. 2019. Occurrence and residue concentration in feed and food of animal origin; human exposure assessment. *Foods* 8(10):477. [Occurrence and Residue Concentration of Coccidiostats in Feed and Food of Animal Origin; Human Exposure Assessment \(mdpi-res.com\)](https://www.mdpi.com/2304-8158/8/10/477)
82. Rokka, M., Eerola, S., Perttilä, U., Rossow, L., Venäläinen, E., Valkonen, E., Valaja, J., Peltonen, K. 2005. The residue levels of narasin in eggs of laying hens fed with unmedicated and medicated feed. *Molecular Nutrition and Food Research*, 49(1):38-42.
83. Rokka, M., Jestoi, M., Peltonen, K. 2013. Trace level determination of polyether ionophores in feed. *Biomed Research International*, vol. 2013, Article ID 151363, 12 pages.
84. Strauch, W. 2002. Causes and control of carry-over and cross-contamination. *Kraftfutter/Feed Magazine*, 85(4):151-159 (Part 1). *Kraftfutter/Feed Magazine*, 85(6):239-249 (Part 2).
85. UK-VMD (United Kingdom-Veterinary Medicines Directorate), 2022. Maximum residue limits in Great Britain. [Maximum Residue Limits in Great Britain \(publishing.service.gov.uk\)](https://www.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/108422/maximum-residue-limits-in-great-britain-2022.pdf)
86. Vandenberghe, V., Delezie, E., Huyghebaert, G., Delahaut, P., Pierret, G., De Backer, P., Croubels, S., Daeseleire, E. 2012. Transfer of the coccidiostats monensin and lasalocid from feed at cross-contamination levels to whole egg, egg white and egg yolk. *Food Additives and Contaminants: Part A*, 29(12):1881-1892.
87. Wong, R.Y., Roxburgh, J.W. 2010. Lasalocid awareness and sampling in Scotland. *International Journal of Environmental Health Research*, 20(3):159-169.
88. Young, R., Nunan, C. 2004. Too hard to crack? Eggs with drug residues. Report Four in The use and misuse of antibiotics in UK agriculture series. Soil association.

Attachment 1

Table: Compilation of residues data for nicarbazin measured in edible tissues from poultry fed nicarbazin medicated feed

Species	Feed level (mg/kg)	Dosing period (days)	LOD (mg/kg)	LOQ (mg/kg)	WHP (days)	Residue level (mg/kg)				Reference
						Liver	Kidney	Muscle	Skin/Fat	
Chicken	100 (WF or DL)	28	NS	0.001	9	0.2238± 0.0742 (DL) 0.0237±0.0039 (WF)	-	0.0024±0.0003 (WF) 0.014±0.0084 (DL)	-	Cannavan and Kennedy, 2000
Chicken	13.5 (WF or DL)	32	NS	0.001	5	1.157 (DL) 0.992 (WF)	-	0.055 (DL) 0.028 (WF)	-	Cannavan and Kennedy, 2000
Chicken	125	28	NS	0.05 liver 0.1 kidney 0.025 muscle 0.025 skin/fat	1 (24 h)	9.249±1.804	3.007±1.095	2.110±0.506	2.327±0.37 2	EFSA 2010a
Chicken	125	42	0.03	0.1	1 (24 h)	14.4-21	2.8-5.4	1.4-2.2	1.6-3.0	Wood and Dowling, 1980; JECFA 1999
Chicken	125	49	0.02 ^A	0.1 ^A	1 (24 h)	2.69-9.12 6.62±1.08	-	0.85-1.23 0.98±0.088	0.66-0.99 0.88±0.042	Kramer, 1990; JECFA 1999; ANADA 200-027
Chicken	50 (+50 lasalocid)	42	NS	NS	1 (24 h)	8.57±1.432	3.51±1.12	1.64±0.294	1.95±0.257	EFSA 2021
*Chicken	50 (+50 monensin)	35	NS	0.1	0.25 (6 h)	8.331 (x+2SD)	1.514 (x+2SD)	1.182 (x+2SD)	1.723 (x+2SD)	EFSA 2017

*Chicken	55 (+55 monensin)	10	NS	NS	0.25 (6 h)	6.857±0.920	0.806±0.584	0.761±0.207	1.269±0.326	EFSA 2017; EFSA 2018b
*Chicken	45.4 (+45.4 narasin)	63	NS	2	0	7.6	-	<2	<2	NADA 138-952a
*Chicken	50 (+50 narasin)	35	NS	0.05 liver 0.1 kidney 0.025 muscle 0.025 skin/fat	0	9.19±0.956	4.29±1.034	1.61±0.149	2.04±0.479	NADA 138-952b; EFSA 2010b
*Chicken	70 (+70 narasin)	42	NS	0.02	0 (3h)	8.988±1.965	3.525±1.485	1.813±0.43	2.018±0.66	EFSA 2019
*Chicken	45 (+27 narasin + 4 lincomycin)	NS	NS	NS	0 (6h)	8.27±1.75	-	-	-	NADA 140-947
*Chicken	45 (+27 narasin + 50 bacitracin + 45.4 roxarsone)	21	NS	NS	0 (6 h)	10.4 (2.0-16.5)	-	-	-	NADA 141-112; NADA 141-113
*Chicken	50 (+50 narasin + 200 bacitracin)	49	0.1	NS	0 (6h)	8.5±2.96	-	-	-	NADA 140-926; NADA 141-124; NADA 141-529
*Chicken	113 (+20 bambarmycins, +50 roxarsone)	48	NS	1	0	32.9±6.87	-	4.7±1.86	6.2±1.55	NADA 140-339

*Turkey	50 (+50 monensin) Turkeys	112 (16 wk)	0.01	0.1	0.25 (6 h)	0.276 (5 <LOQ, 1<LOD)	<LOQ	<LOD	<LOQ	EFSA 2017
*Turkey	109	112 (16 wk)	NS	1	0 (1 h)	1.22 (x+2SD)	<LOQ	<LOQ	<LOQ	EFSA 2018a

*Feeding studies with practical zero withdrawal times (less than 12 hours).

Attachment 2


Table: Compilation of residue data for lasalocid measured in edible tissues from poultry fed lasalocid medicated feed.

Species	Feed level (mg/kg)	Dosing period (days)	Marker Residue	LOD (µg/kg)	LOQ (µg/kg)	WHP (days)	Residue level (µg/kg)				Reference
							Liver	Kidney	Muscle	Skin/Fat	
Chicken	125	7	Lasalocid A	20	NS	0	290	130	50	340	EFSA, 2004
						1	90	30	-	70	
						3	20	30	-	40	
						5	40	30	-	40	
						7	-	-	-	-	
Chicken	90	14	Lasalocid	0.01	0.02	0	400	-	10	-	Kennedy et al., 1995
				(Muscle)	(Muscle)	5	30	-	0.5	-	
				0.09 (Liver)	0.15 (Liver)	7	20	-	0.5	-	
Chicken	125 then 132	34 then 21	Total Radioactivity	NS	NS	0	11930	2480	610	1590,	EMA, 2004
						1	2630	360	60	860	
						2	1720	230	30	220,	
						3	1590	< 200	< 200	140	
						4	1370	< 200	< 200	130, 60	
						5	1150	< 200	< 200	< 200	
Chicken	75 (+50 roxarsone)	56	lasalocid	NS	50	0	< LOQ	< LOQ	< LOQ	370 ±	NADA 141-488
						1	< LOQ	< LOQ	< LOQ	40	
						2	< LOQ	< LOQ	< LOQ	50	
						3	< LOQ	< LOQ	< LOQ	< LOQ	
							< LOQ	< LOQ	< LOQ	< LOQ	
Chicken	75 (+50 roxarsone + 2.0 lincomycin)	56	Lasalocid	NS	50	0	< LOQ	< LOQ	< LOQ	360 ±	NADA 141-488
						1	< LOQ	< LOQ	< LOQ	30	
						2	< LOQ	< LOQ	< LOQ	< LOQ	
						3	< LOQ	< LOQ	< LOQ	< LOQ	
Chicken	125 (+2.2 lincomycin + 50 roxarsone)	56	Lasalocid	NS	50	0	-	-	-	270 ±	NADA 141-488
						1	-	-	-	110	
						2	-	-	-	70 ± 20	
						3	-	-	-	< LOQ	
						4	-	-	-	< LOQ	
Chicken	125 (+55 bacitracin zinc)	42	Lasalocid	NS	15	0				428 ±	NADA 107-996
										127	
Turkey	125	14	Total	20	NS	0.33	3380	430 ±	300 ±	300 ±	NADA

			Radioactivity				± 570	50	10	110	096-298
						1	1430				
						2	1490	200	< LOD	160	
						3	1040	170	< LOD	110	
						4	1100	120	< LOD	100	
						5	870	120	< LOD	140	
								80	< LOD	90	
Turkey	125 (+55 bacitracin zinc)	105	Lasalocid	NS	5	0.25	37.3 ± 30.4	-	-	-	NADA 141-109
Quail	90	27	Lasalocid A	NS	NS	0	-	-	-	298.3	EMA, 2004
						3	-	-	-	55	
						6	-	-	-	30.8	
						9	-	-	-	33.7	
Pheasant	132	7	Lasalocid A	NS	NS	0	28.5	-	-	30.7	EMA, 2004
NS – Not Specified.											

PROPOSAL TO DEVELOP CARRY-OVER RISK MANAGEMENT INFORMATION AND A RISK MANAGEMENT DECISION TOOL**ALTERNATIVE PROPOSAL FROM THE UNITED STATES OF AMERICA (USA)****(For comments)****Introduction**

1. To circumvent the challenge of data availability, the United States proposes that the Codex Committee on Residues of Veterinary Drugs in Foods (CCRVDF) develop carry-over risk management information on a case-by- case basis and a Risk Management Decision Tool (RMDT). Unlike action levels, this approach does not require estimates of the carry-over amount and the associated residue concentrations in the food commodity. Also, this approach avoids potentially establishing a residue level that is incompatible with residue amounts detected in human food commodities caused by unavoidable carry-over in animal feed. Instead, this approach provides a tool that competent authorities can choose to employ when they detect a veterinary drug residue in a human food commodity from a non-target animal recognized by Codex to be associated with unavoidable carry-over of a veterinary drug in animal feed.
2. The proposed RMDT is outlined below, followed by descriptions of how the proposed Residue Risk Score (RRS) is calculated, the role of CCRVDF, and how CCRVDF can communicate this information. Following these descriptions, an example is provided using nicarbazin in eggs from laying hens.

Proposed Risk Management Decision Tool (RMDT)**Risk Management Decision Tool (RMDT) for Residues of Veterinary Drugs in Foods Associated with Unavoidable Carry-over in Animal Feed**

3. The RMDT is a decision tree that enables competent authorities to evaluate the human food safety risk associated with detecting the veterinary drug residue in the human food commodity from the non-target animal when unavoidable carry-over of the veterinary drug in the animal feed is the suspected cause. The competent authority can then make a risk management decision based on the specific residue level that is detected and the associated human food safety risk.
4. In general, the RMDT calls for the calculation of a Residue Risk Score (RRS) that places the detected residue value into the context of the established health-based guidance value (HBGV) for the veterinary drug. An RRS less than or equal to one indicates that the detected residue is not a food safety concern. An RRS greater than one indicates that the detected residue represents a food safety concern. As explained in later detail, the RRS equation will be unique for a specific veterinary drug and human food commodity determined by CCRVDF to be affected by unavoidable carry-over. The specific RRS equation will be developed by CCRVDF, as explained later, and can be communicated in the *Maximum Residue Limits and Risk Management Recommendations for Residues of Veterinary Drugs in Foods* (CXM 2).
5. When the RRS is less than or equal to one, indicating no food safety concern, the decision tree then asks whether the detected residue exceeds the Codex MRLs established for the target animal. The answer to this question informs the risk manager whether the detected residue is likely caused by unavoidable carry-over. If the detected residue exceeds the Codex MRLs established for the target animal, then it is unlikely to be caused by unavoidable carry-over because the Codex MRLs for the target animal are based on a pharmacological dose of the veterinary drug, and true unavoidable carry-over amounts are likely to be much lower than a pharmacological dose. Likewise, if the detected residue is less than the Codex MRLs for the target animal, unavoidable carry-over is likely the cause. This information enables the risk manager to determine if additional follow-up actions are necessary.

Calculation of the Residue Risk Score

6. The Residue Risk Score (RRS) is a metric used to determine whether there is a food safety concern associated with veterinary drug residues in human food commodities caused by unavoidable carry-over of veterinary drugs in animal feed. The RRS value quantifies the human food safety risk in relation to the established Codex HBGV for the veterinary drug. The general calculation is as follows:
- $RRS = (\text{Detected Marker Residue Concentration } (\mu\text{g/kg}) \times \text{Risk Score Correction Factor}) + \text{GVP Risk Score}$
 - Where...
 - Risk Score Correction Factor = a numerical value that converts the detected marker residue concentration to a risk score value
 - GVP Risk Score = Fraction of the HBGV utilized when the drug is used in accordance with Good veterinary Practices (GVPs)
7. The information needed to calculate the RRS (*i.e.*, Risk Score Correction Factor and GVP Risk Score) will be established by CCRVDF on a case-by-case basis for a specific veterinary drug and human food commodity determined by CCRVDF to be affected by carry-over. After the RRS is developed, CCRVDF can communicate it within CXM 2. In doing so, the competent authority only needs to apply the RRS equation to the detected residue value.

Determination of the Risk Score Correction Factor and GVP Risk Score

8. For a specific veterinary drug and human food commodity, CCRVDF will determine the following:
- 8.1 Risk Score Correction Factor:** The risk score correction factor is a numerical value that enables rapid conversion of the detected marker residue concentration in the human food commodity to a risk score value. It converts the measured marker residue value into a number that quantifies the risk of the detected residue in relation to the Codex HBGV. Essentially, this risk score value is the percentage of the Codex HBGV that would be utilized by the detected residue, expressed as decimal. Similar to and building upon Step 4 in the Action Level discussion paper, the risk score correction factor is based on the Theoretical Maximum Daily Intake (TMDI) model and consolidates the mathematical steps for the TMDI calculation into a single step for the affected commodity. The risk score correction factor is derived from the following:
- 8.1.1. TMDI consumption factor for the commodity (*i.e.* 0.3 kg muscle, 0.1 kg liver, 0.05 kg kidney, 0.05 kg fat (skin/fat), 1.5 kg milk, 0.1 kg eggs, 0.02 kg honey)
 - 8.1.2 TMDI human body weight (60 kg)
 - 8.1.3 Health-Based Guidance Value (*e.g.*, ADI or ARfD)
 - 8.1.4 An estimated marker residue to total residue ratio (M:T) determined by CCRVDF.
 - 8.1.4.1 The estimated M:T ratio will be based on the JECFA-derived M:T ratios associated with the veterinary drug in the target animal, applying uncertainty factors as deemed necessary if an experimentally derived M:T ratio is not available for the affected commodity.
9. The calculation is as follows:

$$\text{Risk Score Correction Factor} = \frac{\left(\frac{\text{TMDI consumption value (kg)}}{\text{estimated M:T}} \right)}{(\text{HBGV } (\mu\text{g/kg}) \times 60 \text{ kg body weight})}$$

- 8.2 GVP Risk Score:** CCRVDF will identify the GVP Risk Score based on the JECFA evaluation which recommended the Codex MRLs in the target animal. The GVP Risk Score is the percentage of the HBGV utilized when the veterinary drug is used in accordance with GVPs expressed as a decimal. For example, if JECFA determined that use of the veterinary drug in accordance with GVPs results in 23% of the ADI being utilized, then the GVP Risk Score is equal to 0.23.

Communication of the Residue Risk Score Calculation and Risk Management Decision Tool

10. CCRVDF can communicate the information needed to calculate the RRS within CXM2 for each specific veterinary drug residue and human food commodity determined to be affected by unavoidable carry-over. This could be done by providing text describing the issue and the RRS equation to be applied to the detected marker residue concentration in the affected human food commodity.

11. The Codex Committee on Contaminants in Foods (CCCF) has a similar decision tool for contaminants in food where there is no regulatory level, and CCCF has communicated this according to the *Guidelines for Rapid Risk Analysis Following Instances of Detection of Contaminants in Food where there is No Regulatory Level* (CXG 92-2019)¹. Similarly, CCRVDF could communicate the general information on the RMDT as a Codex Guideline. The Codex Guideline would refer to the *Maximum Residue Limits and Risk Management Recommendations for Residues of Veterinary Drugs in Foods* (CXM 2)² for the information needed to calculate the RRS for each specific veterinary drug residue and human food commodity determined to be affected by unavoidable carry-over.
12. Below is an outline of how this proposed process might work, using nicarbazin in eggs as an example.

Nicarbazin in Eggs Example

Step 1: Review of unavoidable carry-over information

13. CCRVDF would review information on the occurrence of unavoidable carry-over of nicarbazin in laying hen feed and its potential to cause detectable residues in eggs from laying hens.

Step 2: Determination of GVP Risk Score and Risk Score Correction Factor

14. If CCRVDF determines that sufficient information demonstrates that detectable residues in eggs from laying hens is associated with unavoidable carry-over of nicarbazin in laying hen feed, then CCRVDF would determine the appropriate GVP Risk Score and Risk Score Correction Factor.
 - **GVP Risk Score:** The 94th meeting of JECFA determined that the maximum estimated ADI utilization is 23% when nicarbazin is used in accordance with GVPs³. Therefore, the GVP Risk Score is 0.23.
 - **TMDI Consumption Factor for Eggs:** 0.1 kg
 - **CCRVDF-estimated M:T for Eggs:** 0.25 (only an example estimate)
 - **Nicarbazin HBGV (ADI):** 900 µg/kg bw
15. Therefore, the Risk Score Correction Factor for nicarbazin in eggs equals 7.41×10^{-6}

$$\text{Risk Score Correction Factor} = \frac{\left(\frac{0.1 \text{ kg}}{0.25}\right)}{(900 \text{ µg/kg} \times 60 \text{ kg body weight})} = 7.41 \times 10^{-6}$$

Step 3: Communication of risk management information

16. CCRVDF would communicate the risk management information for nicarbazin in eggs in CXM 2. The language that follows is provided only as an example.

NICARBAZIN (coccidiostat)

17. Codex has adopted Maximum Residue Levels (MRLs) for nicarbazin in chicken muscle, liver, kidney, and skin with adhering fat. Evidence indicates that unavoidable carry-over of nicarbazin can occur in non-target animal feed intended for laying hens despite adherence to Good Veterinary Practices and Good Manufacturing Practices. This can result in residues of nicarbazin in eggs from laying hens. For this reason, competent authorities should ensure that appropriate mitigation steps are taken to reduce the carry-over of nicarbazin into feed intended for laying hens. Appropriate mitigation steps can be found in the Code of Practice on Good Animal Feeding (CXC 54). In cases where the marker residue for nicarbazin (DNC) is detected in eggs from laying hens, competent authorities can apply the procedures outlined in the Risk Management Decision Tool (RMDT). The equation for calculating the Residue Risk Score (RRS) used in the RMDT for the marker residue for nicarbazin in eggs from laying hens is the following.

$$\text{RRS} = (\text{Marker Residue Concentration (µg/kg)} \times 7.41 \times 10^{-6}) + 0.23$$

¹ <https://www.fao.org/fao-who-codexalimentarius/committees/committee/related-standards/en/?committee=CCCF>

² <https://www.fao.org/fao-who-codexalimentarius/committees/committee/related-standards/en/?committee=CCRVDF>

³ Evaluation of certain veterinary drug residues in food. Report of the 94th Meeting of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series 1041.

Step 4: Hypothetical Implementation of the Residue Risk Score Calculation and Risk Management Decision Tool

18. Hypothetically, if a competent authority detects the marker residue for nicarbazin (DNC) in eggs at a concentration of 300 µg/kg and employs the RMDT, the competent authority would calculate the RRS to be 0.2322.

$$\text{RRS} = (300 \mu\text{g/kg} \times 7.41 \times 10^{-6}) + 0.23 = 0.2322$$

19. In this hypothetical example, the RMDT would indicate that there is no food safety concern associated with the eggs containing the marker residue for nicarbazin (DNC) at a concentration of 300 µg/kg and that unavoidable carry-over likely is the cause.

POTENTIAL HYBRID APPROACH PROPOSAL

PROPOSAL BY THE UNITED STATES OF AMERICA (USA)

(For comments)

GUIDELINES ON RISK MANAGEMENT DECISIONS FOR RESIDUES OF VETERINARY DRUGS DETECTED IN FOOD CAUSED BY UNINTENTIONAL AND UNAVOIDABLE CARRYOVER OF VETERINARY DRUGS IN ANIMAL FEED WHERE THERE IS NO APPLICABLE CODEX MRL

BACKGROUND

- Description of carryover
- Reference to Code of Practice on Good Animal Feeding (CXC 54)
- Other relevant information


SCOPE

- Description of when this guideline applies

APPLICATION OF RISK MANAGEMENT DECISION TOOL (RMDT)

- Description of the risk management decision tool and how it is used
- Description of Carryover Action Levels (CAL) and how they are used
- Description of the Residue Risk Score (RRS) and how it is used
- Reference to appendix that contains risk management information to be used in the RMDT for compounds and commodities known to be affected unavoidable carryover
- Other relevant information

Risk Management Decision Tool (RMDT) for Residues of Veterinary Drugs in Foods Associated with Unavoidable Carryover in Animal Feed



Step 4: Hypothetical Implementation of the Residue Risk Score Calculation and Risk Management Decision Tool

1. Hypothetically, if a competent authority detects the marker residue for nicarbazin (DNC) in eggs at a concentration of 300 µg/kg and employs the RMDT, the competent authority would calculate the RRS to be 0.2322.

$$\text{RRS} = (300 \mu\text{g}/\text{kg} \times 7.41 \times 10^{-6}) + 0.23 = 0.2322$$

2. In this hypothetical example, the RMDT would indicate that there is no food safety concern associated with the eggs containing the marker residue for nicarbazin (DNC) at a concentration of 300 µg/kg and that unavoidable carry-over likely is the cause.

APPENDIX V**LIST OF PARTICIPANTS**

Chair Australia Dugald MacLachlan Department of Agriculture, Fisheries and Forestry	Vice-Chair Canada Cole Enns Health Canada
---	---

MEMBER COUNTRY/ORGANIZATION¹	OBSERVER²
1. Argentina	1. International Feed Industry Federation (IFIF)
2. Australia	
3. Brazil	
4. Canada	
5. Chile	
6. Costa Rica	
7. Denmark	
8. Egypt	
9. France	
10. Germany	
11. Honduras	
12. Hungary	
13. India	
14. Indonesia	
15. Iran	
16. Italy	
17. Japan	
18. Maroc	
19. Panama	
20. Republic of Korea	
21. Saudi Arabia	
22. Thailand	
23. United Arab Emirates	
24. United States	