

Diversity of non-*Apis* insect pollinators in open pollinated flowers (sweep net captures)

COUNTRY: INDIA

SITE : KOSI KATARMAL

FOCUS CROP : MUSTARD (*Brassica campestris*)

YEAR :

Field number & size (ha)	Location	TREATMENT	Date & observer	Recording number	RECORDING CONDITIONS			Transect number in a day	TYPE (SPECIES) & NUMBER OF SPECIMENS PER TYPE				Remarks
					Time at start	Period	Weather conditions		BEES OTHER THAN HONEY BEES		DRONE FLIES (Syrphidae)	OTHER	
								Bumble bees (<i>Bombus</i> spp.)	other wild bees				
	Valley, mid-mountain, hill top	Honey bee colonies nearby present / absent*		1, 2, 3, 4	1000–1200 h or 1400–1600 h	Morning or Afternoon	Sunny, overcast, strong wind (instantaneous temperature if available)	1, 2, ... 6	Flower visiting insects that are likely pollinators caught by sweep net over a 5 minute period along a 25 m transect within or along side the field with a total of 6 transects				

* Indicate the number of colonies of *Apis cerana* and/or *Apis mellifera* nearby the study field & whether these colonies were present or introduced at onset of flowering

				0 ≈ prior to colony introduction (if applicable)				1									
								2									
								3									
								4									
								5									
								6									
				1								1					
												2					
												3					
												4					
												5					
												6					

Diversity of non-*Apis* insect pollinators in open pollinated flowers (sweep net captures)

COUNTRY: INDIA

SITE : KOSI KATARMAL

FOCUS CROP : MUSTARD (*Brassica campestris*)

YEAR :

Field number & size (ha)	Location	TREATMENT	Date & observer	Recording number	RECORDING CONDITIONS			Transect number in a day	TYPE (SPECIES) & NUMBER OF SPECIMENS PER TYPE				Remarks
					Time at start	Period	Weather conditions		BEES OTHER THAN HONEY BEES		DRONE FLIES (Syrphidae)	OTHER	
								Bumble bees (<i>Bombus</i> spp.)	other wild bees				
	Valley, mid-mountain, hill top	Honey bee colonies nearby present / absent*		1, 2, 3, 4	1000–1200 h or 1400–1600 h	Morning or Afternoon	Sunny, overcast, strong wind (instantaneous temperature if available)	1, 2, ... 6	Flower visiting insects that are likely pollinators caught by sweep net over a 5 minute period along a 25 m transect within or along side the field with a total of 6 transects				

* Indicate the number of colonies of *Apis cerana* and/or *Apis mellifera* nearby the study field & whether these colonies were present or introduced at onset of flowering

				2				1								
								2								
								3								
								4								
								5								
								6								
				3							1					
											2					
											3					
											4					
											5					
											6					

Diversity of non-*Apis* insect pollinators in open pollinated flowers (sweep net captures)

COUNTRY: INDIA

SITE : KOSI KATARMAL

FOCUS CROP : MUSTARD (*Brassica campestris*)

YEAR :

Field number & size (ha)	Location	TREATMENT	Date & observer	Recording number	RECORDING CONDITIONS			Transect number in a day	TYPE (SPECIES) & NUMBER OF SPECIMENS PER TYPE				Remarks
					Time at start	Period	Weather conditions		BEES OTHER THAN HONEY BEES		DRONE FLIES (Syrphidae)	OTHER	
								Bumble bees (<i>Bombus</i> spp.)	other wild bees				
	Valley, mid-mountain, hill top	Honey bee colonies nearby present / absent*		1, 2, 3, 4	1000–1200 h or 1400–1600 h	Morning or Afternoon	Sunny, overcast, strong wind (instantaneous temperature if available)	1, 2, ... 6	Flower visiting insects that are likely pollinators caught by sweep net over a 5 minute period along a 25 m transect within or along side the field with a total of 6 transects				

* Indicate the number of colonies of *Apis cerana* and/or *Apis mellifera* nearby the study field & whether these colonies were present or introduced at onset of flowering

				4				1					
								2					
								3					
								4					
								5					
								6					