



Vanuatu STEPS Survey 2011

Fact Sheet

The STEPS survey of chronic disease risk factors in Vanuatu was carried out from end of July 2011 to end of October 2011. Vanuatu carried out Step 1, Step 2 and Step 3. Socio demographic and behavioural information was collected in Step 1. Physical measurements such as height, weight and blood pressure were collected in Step 2. Biochemical measurements were collected to assess blood glucose and cholesterol levels in Step 3. The STEPS survey in Vanuatu was a population-based survey of adults aged 25-64. A multi-stage cluster sampling design was used to produce representative data for that age range in Vanuatu. A total of 4671 adults participated in the Vanuatu STEPS survey.

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females
Step 1 Tobacco Use			
Percentage who currently smoke tobacco	52.5% (49.4 – 55.6)	62.3% (59.3 – 65.3)	20.2% (15.1 – 25.3)
Percentage who currently smoke tobacco daily	27.5% (24.8 – 30.2)	33.5% (30.1 – 36.8)	7.8% (5.0 – 10.6)
<i>For those who smoke tobacco daily</i>			
Average age started smoking (years)	20.2 (19.7 – 20.8)	20.2 (19.7 – 20.7)	*
Percentage of daily smokers smoking manufactured cigarettes	52.0% (45.7 – 58.4)	50.0% (43.5 – 56.5)	*
Mean number of manufactured cigarettes smoked per day (by smokers of manufactured cigarettes)	4.1 (3.0 – 5.3)	4.0 (2.9 – 5.0)	*
Step 1 Alcohol Consumption			
Percentage who are lifetime abstainers	48.8% (46.4 – 51.3)	22.3% (20.0 – 24.6)	71.9% (68.9 – 74.9)
Percentage who are past 12 month abstainers	31.2% (28.9 – 33.4)	43.9% (40.8 – 46.9)	20.2% (17.7 – 22.6)
Percentage who currently drink (drank alcohol in the past 30 days)	6.8% (5.5 – 8.2)	11.9% (9.8 – 14.1)	2.4% (1.3 – 3.5)
Percentage who engage in heavy episodic drinking (men who had 5 or more / women who had 4 or more drinks on any day in the past 30 days)	--	8.6% (6.8 – 10.5)	1.7% (0.9 – 2.6)
Step 1 Fruit and Vegetable Consumption (in a typical week)			
Mean number of days fruit consumed	3.9 (3.7 – 4.0)	4.0 (3.9 – 4.2)	3.7 (3.5 – 3.9)
Mean number of servings of fruit consumed on average per day	1.9 (1.7 – 2.1)	2.0 (1.8 – 2.3)	1.7 (1.6 – 1.9)
Mean number of days vegetables consumed	5.5 (5.4 – 5.6)	5.6 (5.5 – 5.7)	5.5 (5.3 – 5.6)
Mean number of servings of vegetables consumed on average per day	3.1 (2.9 – 3.4)	3.3 (3.0 – 3.6)	3.0 (2.8 – 3.2)
Percentage who ate less than 5 servings of fruit and/or vegetables on average per day	61.8% (57.9 – 65.6)	58.2% (53.4 – 63.0)	65.0% (61.2 – 68.7)
Step 1 Physical Activity			
Percentage with low levels of activity (defined as < 600 MET-minutes per week)*	11.7% (10.3 – 13.1)	10.1% (8.4 – 11.7)	13.2% (11.2 – 15.2)
Percentage with high levels of activity (defined as ≥ 3000 MET-minutes per week)*	73.9% (71.5 – 76.3)	81.7% (79.2 – 84.2)	66.9% (63.8 – 70.0)
Median time spent in physical activity on average per day (minutes) (presented with inter-quartile range)	214.3 (98.6 – 402.9)	278.6 (132.9 – 471.4)	171.4 (81.4 – 325.7)
Percentage not engaging in vigorous activity	32.4% (28.5 – 36.3)	20.6% (17.0 – 24.1)	43.1% (38.4 – 47.7)

* For complete definitions of low and high levels of physical activity, other conditions are specified in the GPAQ Analysis Guide, available at: <http://www.who.int/chp/steps/GPAQ/en/index.html>



Vanuatu STEPS Survey 2011

Fact Sheet

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females
Step 2 Physical Measurements			
Mean body mass index - BMI (kg/m ²)	26.1 (25.9 – 26.4)	25.5 (25.2 – 25.8)	26.7 (26.4 – 27.0)
Percentage who are overweight (BMI ≥ 25 kg/m ²)	50.9% (48.8 – 53.1)	45.5% (42.8 – 48.3)	55.9% (53.2 – 58.5)
Percentage who are obese (BMI ≥ 30 kg/m ²)	18.8% (16.9 – 20.8)	13.9% (11.9 – 15.9)	23.3% (20.7 – 25.8)
Average waist circumference (cm)	--	75.0 (73.6 – 76.4)	79.7 (78.2 – 81.1)
Mean systolic blood pressure - SBP (mmHg), including those currently on medication for raised BP	129.3 (128.3 – 130.3)	131.7 (130.4 – 132.9)	127.2 (126.0 – 128.4)
Mean diastolic blood pressure - DBP (mmHg), including those currently on medication for raised BP	79.5 (78.7 – 80.3)	79.8 (78.9 – 80.7)	79.2 (78.3 – 80.0)
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)	28.6% (26.5 – 30.8)	30.8% (28.0 – 33.5)	26.7% (24.2 – 29.2)
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg) who are not currently on medication for raised BP	94.8% (93.3 – 96.3)	95.2% (93.0 – 97.4)	94.4% (92.4 – 96.4)
Step 3 Biochemical Measurement			
Mean fasting blood glucose, including those currently on medication for raised blood glucose: mmol/L	5.7 (5.6 – 5.8)	5.7 (5.6 – 5.8)	5.7 (5.6 – 5.8)
Percentage with impaired fasting glycaemia as defined below • capillary whole blood value ≥5.6 mmol/L (100 mg/dl) and <6.1 mmol/L (110 mg/dl)	18.8% (17.1 – 20.5)	19.0% (16.6 – 21.5)	18.6% (16.6 – 20.7)
Percentage with raised fasting blood glucose as defined below or currently on medication for raised blood glucose • capillary whole blood value ≥ 6.1 mmol/L (110 mg/dl)	21.2% (19.0 – 23.3)	21.4% (18.8 – 24.0)	21.0% (18.5 – 23.5)
Mean total blood cholesterol, including those currently on medication for raised cholesterol: mmol/L	4.9 (4.9 – 5.0)	5.0 (4.9 – 5.1)	4.9 (4.8 – 4.9)
Percentage with raised total cholesterol (≥ 5.0 mmol/L or ≥ 190 mg/dl or currently on medication for raised cholesterol)	36.5% (34.2 – 38.7)	38.4% (35.3 – 41.6)	34.7% (32.1 – 37.3)
Summary of combined risk factors			
<ul style="list-style-type: none"> • current daily smokers • less than 5 servings of fruits & vegetables per day • low level of activity • overweight (BMI ≥ 25 kg/m²) • raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP) 			
Percentage with none of the above risk factors	9.7% (7.5 – 11.9)	10.8% (8.2 – 13.3)	6.0% (3.6 – 8.5)
Percentage with three or more of the above risk factors, aged 25 to 44 years	18.7% (15.8 – 21.6)	19.7% (16.1 – 23.2)	15.8% (11.3 – 20.2)
Percentage with three or more of the above risk factors, aged 45 to 64 years	30.6% (26.0 – 35.2)	28.0% (23.2 – 32.8)	42.6% (32.2 – 53.1)
Percentage with three or more of the above risk factors, aged 25 to 64 years	22.3% (19.3 – 25.2)	22.3% (18.9 – 25.7)	22.1% (17.2 – 27.1)

* Indicates less than 50 respondents.