

IPRODIONE (111)

EXPLANATION

Iprodione, originally evaluated by the JMPR in 1977 and re-evaluated for residues in 1980, is included in the CCPR periodic review programme. Re-evaluation (ALINORM 89/24A, para 298; Appendix V and VI) was originally scheduled for 1992 but postponed because of the work-load and the late submission of some of the data.

Codex Maximum Residue Limits (CXLs) exist for apple, beans (dry), cucumber, black, red and white currants, garlic, grapes, kiwifruit, head lettuce, bulb onion, peach, pear, sweet peppers, plums, raspberries, husked rice, strawberry, tomato and witloof chicory.

The Meeting reviewed information from the manufacturer in response to the 1977 requests (repeated in 1980), as well as information on GAP and residue trials on other commodities. Additional information was submitted by The Netherlands, Spain, New Zealand, Canada and Germany.

IDENTITY

ISO common name: iprodione

Chemical names

(IUPAC): 3-(3,5-dichlorophenyl)-*N*-isopropyl-2,4-dioxoimidazolidine-1-carboxamide

(C.A.): 3-(3,5-dichlorophenyl)-*N*-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide

CAS No: 36734-19-7

Synonyms: glycophene (non-official: rejected common name proposal)

Structural formula:

Molecular formula: $C_{13}H_{13}Cl_2N_3O_3$

Molecular weight: 330.17

Physical and chemical properties

Vapour pressure: <0.133 mPa (20°C)
 Melting point: 136°C
 Octanol/water
 partition coefficient: $\log P_{ow} = 3.1$
 Solubility (20°C): water 0.013 g/l
 ethanol 25 g/l
 acetone 300 g/l
 dichloromethane 500 g/l

USE PATTERN

Extensive information on GAP was given in the 1977 monograph. The Meeting received additional or revised information on current GAP (Table 1).

Formulations

Rovral WP (50% WP); Rovral Aqua Flo (500 g/l SC); Rovral Flo = Kidan = Verisan (255 g/l SC); Rovral 4F (480 g/l SC); Rovral TS (35% iprodione, 17.5% carbendazim WP); Rovral TS Flo = Calidan = Quintal IC (175 g/l iprodione, 87.5 g/l carbendazim SC); Rovral UFB (350 g/l iprodione, 175 g/l carbendazim FS); Rhapsodie (350 g/l iprodione, 50 g/l imazalil SC); Sumistar (160 g/l iprodione, 24 g/l diniconazole and 80 g/l carbendazim SC); Sirocco (100 g/l iprodione, 150 g/l fenpropimorph).

Table 1. Registered uses of iprodione.

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
Alfalfa	Australia		0.13-0.25	0.012-0.025	7	(1)
(1) PHI before grazing						
Almonds	France		0.75	0.075		
	USA	4	0.56	0.03		(1)
(1) Last application up to 5 weeks after petal fall						
Apple	Australia	1		0.05	-	(1)
	Greece	2		0.05	14	
	Italy	1		0.05-0.075	10	(1) (2)
	Norway	1-2		0.15	14	
	Spain	1		0.05		(1)

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	Switzerland	2	1.0	0.05	14	
(1) Post-harvest treatment (2) PHI is time of storage before marketing						
Apricot	Australia	1		0.05	-	(1)
		2-3		0.025-0.037	-	(2)
	France	2-3		0.075	3	
	Greece	2		0.05	14	
	New Zealand	2-3	up to 1.0	0.037	1	
	Spain	2	0.5-0.75	0.05-0.075	14	
	USA	5	0.56-1.12	0.03-0.06	0	
(1) Post-harvest dipping (2) Pre-harvest spraying						
Asparagus	Belgium	1-4	0.75		-	(1)
	Luxembourg	1-4	0.75	0.075	-	(1)
	Netherlands	3-4	0.75		-	(1)
	New Zealand		0.75		-	(1)
	Switzerland	3	0.75			
(1) Application after harvesting to extend the "green period" of the plant, thus increasing the yield in the following year						
Banana	Spain	1		0.03		(1)
(1) Post-harvest treatment						
Barley	France	1-2	0.4			
	Belgium		0.4-0.47		42	
	Germany	1	0.5-0.7/t seed		-	st ¹
	Luxembourg		0.4-0.47		42	
	UK	1-2	0.5	0.125-0.25	-	(1)
(1) Last application up to growth stage Zadoks 59						
Bean, Common	Canada (4)	1	0.5-0.75	0.17-0.25		(5)
	(6)	2	0.75	0.25		
	Denmark	1-2	0.5-0.75		21	(7)
	France	2-4	0.5-0.75		3	
	Greece	1-2		0.075	7	
	Netherlands	1-2	0.5-0.75	0.025-0.037	14 F ²	(3)
	Norway	2	1.5		14	
	Portugal	4	0.75	0.075	3	
	Sweden	1	0.5	0.07	14	
	Switzerland	2	0.8		-	(8)
	UK	1-2	0.5	0.08-0.25	21	
	USA (1)	2	0.84-1.12	up to 0.3		(2)
(1) Dry snap beans (2) Last application not later than full bloom. Do not allow foraging for 14 days after last application. Do not feed snap bean hay to livestock. Do not feed dry bean hay to livestock until 45 days after last application. Do not feed succulent bean hay to livestock. (3) Dwarf snap beans (4) Dry common beans (5) Apply when 25-75% of blooms have opened (6) Snap beans (7) Field beans (8) Application during flowering						
Beetroot	France	1	2.5 g/kg seed			
	Norway	1-2	1.0		14	
Berries	Australia			0.5	1	(1)
	Denmark	3	0.5-0.75		14	
	Netherlands	2-4	1.13	0.075	14	
	New Zealand	3	up to 1.0	0.05	1	

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	Sweden	2	0.5-0.75	0.06	30	
(1) Post-harvest dipping						
Blackberries	Netherlands	2	1.13	0.075	7	
	New Zealand	3	up to 1.0	0.05	1	
	Switzerland	1-2		0.05	14	
Brassica vegetables, Head cabbages, Flowerhead brassicas	Australia	1		0.075 200 ml/plant	-	(1)
	Denmark	1-3	0.5		14	
	France	1	2.5 g/kg seed		-	st
		2-3	0.5-0.75		15	
	Greece	1-2		0.075	14	
	New Zealand	1	2.5 g/kg seed			st
	Norway	1-2	1.5		14	
	Spain	4	0.75-1.12	0.05-0.075	0-1	F,G ³
	Sweden	1-2	0.75	0.15	14	
	Switzerland	1	0.2			
	UK	1		0.05	60	drench (2)
				27 l/400 cabbages		stored cabbages
		1	2.5 g/kg seed		-	st
(1) Pre-emergence application (2) Soil drench at planting						
Brassica vegetables (seed crops)	UK	1-3	0.5-0.75	0.125-0.375	21	
Broccoli	Netherlands	1	2.0		-F,G	(1)
		1-2	0.5		14F,G	
	Norway	1-2	1.5		14	
	UK	3	0.5	0.08-0.25	21	
	USA	2	1.12	up to 0.3	0	
(1) Pre-emergence application						
Brussels sprouts	Belgium	1-3	0.5		14	
	Luxembourg	1-3	0.5		14	
	Netherlands	1-2	0.5		14F,G	
	Norway	1-2	1.5		14	
	UK	3	0.5	0.08-0.25	21	
Bulb vegetables	Spain	4	0.75-1.12	0.05-0.075	1F,G	
		2-4	0.4-0.6			
Cabbage, Head	Belgium	1	2.5 g/kg seed			st
	Denmark	1-3	0.5		14	
	France	1	2.5 g/kg seed	-		st
		2-3	0.5-0.75		15	
	Greece	1-2		0.075	14	
	Luxembourg	1	2.5 g/kg seed			
	Netherlands	1-2	0.5		14F,G	st
	Norway	1-2	1.5		14	
	Sweden	1-2	0.75	0.15	14	
	UK	3	0.5	0.08-0.25	21	
Canola	Canada	1	0.5-0.75	up to 1.6		(1)
(1) Application when 20-30% of blooms have opened						
Caneberries	USA	4	0.56-1.12	up to 0.12	0	
Caraway	Belgium	2-3	0.75		28	

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	Luxembourg	2-3	0.75		28	
	Netherlands	1-2	0.5		28	
Carrot	Belgium	3-4	0.75		21	
	Denmark	1-2	0.5		14	
	France	1	2.5 g/kg seed			st
		2-3	0.75			
	Luxembourg	3-4	0.75		21	
	Netherlands	1	0.5 g/kg seed			st
		1	0.03 kg/t roots			(1)
		1-4	0.75		28F	
	Norway	3-4	1.5		14	
	Sweden	2-3	0.5	0.06	14	
	Switzerland	2-3	0.3-0.8	0.05	21	
	USA	4	0.56-1.12	up to 1.2	0	
(1) pre-storage treatment						
Cauliflower	Belgium	2-3	0.125		14	
	France	1	2.5 g/kg seed		-	st
		2-3	0.75		15	
	Luxembourg	2-3	0.125		14	
	Norway	1-2	1.5		14	
	Netherlands	1	2.0		-F,G	(1)
		1-2	0.5		14F,G	
	UK	3	0.5	0.08-0.25	21	
(1) Pre-emergence application						
Celeriac	Netherlands	1-2	0.5		14F	
	Norway	1-2	1.5		14	
Celery	Australia	5	0.5	0.05	1	
Cherries	Australia	1		0.05	-	(1)
		2-3		0.025-0.037	-	(2)
	Canada		0.75-0.88	0.025	1	
	Greece	2	1.0	0.05	14	
	Netherlands	2	1.5	0.075	7	
	New Zealand	2-3	up to 1.0	0.037	1	
	Norway	1-5		0.15	14	
	Spain	2	0.5-0.75	0.05-0.075	14	
	USA	5	0.56-1.12	0.03-0.06	-	
		1		0.24		(1)
(1) Post-harvest treatment (2) Pre-harvest spraying						
Chicory	Norway	1-2	0.75		14	
	Switzerland	2	0.3-0.8			(1)
(1) Last application 14 days after planting						
Chinese cabbage	Belgium	2	2.5		14	
	Luxembourg	2	2.5		14	
	Netherlands	2	0.5-0.75		14F,G	
	Norway	1-2	1.5		14	
	Switzerland	2	0.3-0.8	0.05	21	
Chives	Spain	4	0.75-1.12	0.05-0.075	1F,G	
		2-4	0.4-0.6			
Citrus fruits	Italy	1		0.05-0.075	30	(1) (2)
(1) Post-harvest treatment						

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
(2) PHI is time before marketing						
Corn salad	Belgium	2-3	0.75		28	
	France	1	2.5 g/kg seed			st
		3-4	0.75		21	
	Luxembourg	2-3	0.75	0.075	28	
	Netherlands	1	1.0-2.0		-	(1)
(1) Application within 7 days after sowing						
Cress	France	1	2.5 g/kg seed		-	st
Crucifers, oilseed	Finland	1	0.5-0.75		10	
	France	1-2	0.5	0.005	21	
Cucumber	Canada			0.05	2G	
	Denmark	1-3		0.05	3G	
					14F	
	France	2-7	0.75		3	
	Greece	3		0.025	7	
	Netherlands	2-3	0.63-1.25	0.025-0.05	3G	
	Norway	1-6	1.0		4	
	Portugal	3	0.75	0.075	7	
	Sweden	1-3	0.5	0.06	2	
	UK	4		0.05	2	
Cucurbits	Belgium	1-2		0.05	3	
	Greece	3		0.025	7	
	Luxembourg	1-2		0.05	3	
	Spain	3	0.5-0.75	0.05-0.075	7	
Dandelion	France	1	0.05 /m ³ soil			
Dewberries (incl. Boysenberry and Loganberry)	Australia	1	0.5	0.05	1	
Egg plant	Belgium	2-3		0.05	3	
	France	2	0.75-1.0			
	Greece	3		0.05	7	
	Italy	2-3	0.5-0.75	0.05-0.075	21	
	Luxembourg	2-3		0.05	3	
	Netherlands	2-3	0.63-1.25	0.025-0.05	3G	
Endive	Belgium	3	0.75		28	
	France	1		0.06		(2)
		1	0.05 /m ³ soil	0.005		
	Italy	2	0.75	0.075-0.1	21	
	Netherlands	1-2	0.75-2.0		-F,G	(1)
(1) Application within 14 days after planting						
(2) Plant dipping						
Fennel	Netherlands	1-2	0.75		14	
	Spain	4	0.75-1.12	0.05-0.075	1F,G	
		2-4	0.4-0.6			
Fruiting vegetables, cucurbits	Spain	4	0.75-1.12	0.05-0.075	7F,G	
		2-4	0.4-0.6		28	
Garlic	Canada	1	0.4			seed dip
	France	1	1.5 kg/			
			0.75			
	Greece	1-2		0.075	14	
	USA	1	2.2	up to 1.2	-	(1)
(1) Application in the furrow at planting						

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
Gherkin	France	2-3	0.75			
	Netherlands	2	0.63-1.25	0.025-0.05	3G	
		2-3	0.63-1.25	0.025-0.05	3F	
Ginseng	Canada	3	0.55	up to 0.028	30	
	USA	5-7	0.56-1.12	up to 1.2	36	
Grapes	Australia	4	min. 0.5	0.05	7	
	Canada	2	0.75	0.083	-	(3)
	France	1-4	0.75	0.075	15	
	Germany	3	up to 0.75	0.037	28	
	Greece	4	0.5	0.05	28	
	Italy	2-3	0.75	0.075	28	
	Luxembourg			0.037		
	New Zealand	3	up to 1.0	0.037-0.05	1	
	Portugal	4	0.75	0.075	21	
	Spain	3	0.5-0.75	0.05-0.075	14	(1)
		3	0.4-0.6		28	(2)
	Switzerland	2	1.0	0.05	21	
	USA	4	0.56-1.1	up to 0.24	0	
(1) Foliar spray						
(2) Dust						
(3) Last application before complete bunch closure						
Hazelnuts	France		0.75	0.075		
Kale	Greece	1-2		0.075	14	(1)
	UK	3	0.5	0.08-0.25	21	
(1) Leaf brassicas						
Kale (seed crop)	UK	1-3	0.5-0.75	0.13-0.38	21	
Kiwifruit	Australia	5	0.5	0.05	7	
	Italy	1-2	0.5-0.75	0.05-0.075	15	
	New Zealand	2-3	up to 0.75	0.037	1	
Kohlrabi	Belgium		2.0		14	
	Luxembourg		2.0		14	
	Norway	1-6	1.0		14	
	Netherlands	1	2.0		14F,G	
Leafy vegetables	Spain	4	0.75-1.1	0.05-0.075	14F,G	(1)
		2-4	0.4-0.6			
(1) Except chives						
Leek	Spain	4	0.75-1.1	0.05-0.075	16,F	
		2-4	0.4-0.6			
Legume vegetables	Spain	4	0.75-1.1	0.05-0.075	5F,G	
Lentil	France	2	0.52			
Lettuce, Head	Australia		0.5	0.05	7	
	Belgium	3	0.75		28	
	Denmark	1-3	0.25-0.5	0.025-0.05	14G	
	France	3-4	0.75-0.8	0.075	21F,G	
	Germany	3	0.5	0.083	14F, 21G	
	Greece	1-2	0.075		14	
	Italy	2	0.75	0.075	21	
	Luxembourg	3	0.75		28	
	Netherlands	1-2	0.75-2.0		-F,G	(3)
	Norway	1-5	0.75		14	
	Portugal		0.75	0.075	15F,21G	

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	Switzerland	2	0.3-0.8	0.05	-	(4)
	UK	7	0.25	0.025	7F,G	(1)
		3	0.25	0.025	28G	(2)
	USA	3	0.84-1.12	up to 0.3	14	
(1) Treated between March and September						
(2) Treated between October and February						
(3) Application within 14 days after planting (four for iceberg lettuce)						
(4) Application within 14 days after planting						
Lettuce, Leaf	Australia	3	0.5	0.05	7	
	Belgium	3	0.75		28	
	Denmark	1-3	0.25-0.5	0.025-0.05	14G	
	France	3-4	0.75	0.075	21	
	Greece	1-2	0.075		14	
	Italy	2	0.75	0.075	21	
	Luxembourg	3	0.75		28	
	Netherlands	1-2	0.75-2.0		-F,G	(1)
	Norway	1-5	0.75		14	
	Switzerland	2	0.3-0.8	0.05	-	(1)
	USA	3	0.84-1.1	up to 0.3	14	
(1) Application within 14 days after planting						
Lima bean	USA	2	0.84-1.1	up to 0.3		(1)
(1) Do not allow foraging for 14 days after last application. Do not feed dry bean hay to livestock until 45 days after last application. Do not feed succulent bean hay to livestock. Last application not later than full bloom.						
Linseed	UK	1	2.5 g/kg seed		-	st
Lupin	Australia	1	1.0-1.3 kg/t seed		-	st
Melons	Denmark	1-2	0.5		3G	
	Greece	1-2		0.075	7	
	Netherlands	2-3	0.63-1.3	0.025-0.05	3G	
Mushrooms	Spain	1	0.5-0.75	0.05-0.075	45	
Mustard, Chinese	USA	4	0.56	up to 0.12	10	
Mustard (seed crop)	UK	1-3	0.5-0.75	0.13-0.38	21	
Nectarine	Australia	1		0.05	-	(1)
		2-3		0.025-0.037	-	(2)
	New Zealand	2-3	up to 1.0	0.038	1	
	USA	5	0.56-1.1	0.03-0.66	0	
		1		0.24		(1)
(1) Post-harvest treatment						
(2) Pre-harvest spraying						
Oats	Belgium	1	0.4		42	
	Germany	1	0.5-0.7 kg/t seed		-	st
	Luxembourg	1	0.4		42	
Onion, Bulb	Canada	2	0.75	0.15-0.25	15	
			0.38	0.075-0.13	15	
	Denmark	1-3	0.5		14	
	France	1	1.5 kg/t bulbs			
		1	1.5 g/kg seed			st
			0.75			
	Greece	1-2		0.075	14	
	Netherlands	2-4	0.25-0.5		28	
	New Zealand	1	125 g/kg seed		-	
	Sweden	2-3	0.5-0.75	0.25	14	
	Switzerland	1	1-1.25	0.025-0.03	21	

iprodione

709

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	UK	6	0.5	0.08-0.16	7	
	USA	5	0.56-0.84	up to 0.18	7	(1)
				up to 0.9	7	(2)
Onions, Salad	UK	4	0.5	0.08-0.16	7	
Parsley	France	1	2.5 g/kg seed		-	st
			0.75			
Passion fruit	Australia		0.5	0.05	7	
Peach	Australia	1		0.05	-	(3)
		2-3		0.025-0.037	-	(4)
	Canada		0.75-0.88	0.025	1	(4)
	France	2-3	1.5	0.075	3-15	(4)
	Greece	2		0.05	14	
	New Zealand	2-3	up to 1.0	0.037	1	(4)
	Spain	2	0.5-0.75	0.05-0.075	14	
	USA	5	0.56-1.1	0.03-0.06	0	(4)
		1		0.24		(3)
(1) Ground application (2) Aerial application (3) Post-harvest treatment (4) Pre-harvest spraying						
Peanut	Australia			0.11	12	
	USA	3	1.1	up to 0.3	10	
Pear	Australia	1		0.05	-	(1)
	Greece	2		0.05	14	
	Italy	1		0.05-0.075	10	(1) (2)
		3-4	1.1	0.075	21	
	Spain	1		0.05	-	(1)
	Switzerland	2	1.0	0.05	14	
(1) Post-harvest treatment (2) PHI is time of storage before marketing						
Peas	Belgium	2	0.7			
	France	2	0.52-0.75		30	
	Denmark	1-2	0.5-0.75		14	
	Germany	1	0.76	0.19	35	(3)
	Luxembourg	2	0.7			
	Netherlands	1-2	0.5-0.75		14	(1)
					21	(2)
	Norway	1-2	1.5		14	
	Sweden	1	0.75	0.25	-	
	UK	1-2	0.5	0.13-0.25	21	
(1) PHI for dry peas (2) PHI for fresh peas (3) Field peas						
Peppers, Sweet	Denmark	1-3	0.5	0.05	3	
	France	2-4	0.75-1.0		3	(1)
	Greece	1-2		0.075	3	
	Netherlands	2-3	0.63-1.3	0.025-0.05	3G	
	Spain	4	0.75-1.1	0.05-0.075	14F,G	
		2-4	0.4-0.6		7	
(1) Green pepper						
Pimento	Portugal	3	0.75	0.075	7	
Plums	Australia	1		0.05	-	(1)
		2-3		0.025-0.037	-	(2)

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	France	2-3		0.075	3	
	New Zealand	2-3	up to 1.0	0.037	1	
	Spain	2	0.5-0.75	0.05-0.075	14	
	USA	5	0.56-1.1	0.03-0.06	0	
		1		0.24		(1)
(1) Post-harvest treatment (2) Pre-harvest spraying						
Potato	Australia		0.25-0.5	0.025-0.05		
	Denmark	1	0.4-1.3		14	
	USA	4	0.56-1.1	up to 1.2	14	
Potato (seed crop)	Australia	1	0.2 kg/t		-	
	Belgium	1	0.2 kg/t		-	
		1		0.75	-	(1)
	Denmark	1		0.75	-	(1)
		1		0.375	-	(2)
	France	1	0.75-0.1 kg/t		-	(2)
		1	0.3-0.4		-	(1)
	Greece	1	0.6-0.8		-	(1)
	Luxembourg	1	0.2 kg/t		-	
		1		0.75	-	(1)
	Netherlands	1	0.1-0.2 kg/t		-	
	UK	1	0.1 kg/t		-	
(1) Dipping at planting (2) Spraying at planting						
Prunes	USA	5	0.56-1.1	0.03-0.06	0	
Radish	Netherlands	1-2	2.0		14F,G	
Radish, Black	Netherlands	1-2	2.0		14F,G	
Rape, oilseed	Belgium	2	0.43			
	Denmark	1-2	0.5		21	
	France	1	0.5-0.75	0.005		
	Germany	1	0.76	0.13-0.15	56	
	Italy	1	0.5	0.1	45	
	Luxembourg	2	0.43			
	Netherlands	1-2	0.5		-	(1)
	Sweden	1	0.5-0.75	0.25	-	
	UK	1-3	0.5-0.75	0.13-0.38	21	
			2.5 g/kg seed		-	
(1) Application at beginning/end of flowering						
Raspberries	Australia	1		0.05	1	
	Norway	3-4	1.5		14	
	Netherlands	2	1.1	0.075	7	
	New Zealand	3	up to 1.0	0.05	1	
	Switzerland	1-2		0.05	14	
	UK		0.75	0.075	7	
Rice	USA	2	0.56	up to 0.6		(1)
(1) Last application not later than 75% heading						
Root and tuber vegetables	Spain	4	0.75-1.1	0.05-0.075	3F,G	
		2-4	0.4-0.6			
Rye	Germany	1	0.5-0.7/t seed		-	
Shallot	Belgium	1	200g/100kg bulbs		-	
		1		0.75	28	

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	France	1	1.5 g/kg seed			st
			0.75			
	Luxembourg	1	2.0 g/kg seed		-	st
		1		0.75	28	
	Netherlands	2-4	0.25-0.5		28	
Soya bean	Australia	2	0.5	0.13-0.25	51	
Squash, Summer (= courgette, zucchini)	France	2-3	0.75			
	Netherlands	2-3	0.63-1.3	0.025-0.05	3G	
Stone fruits	Belgium	2		0.1	14	
	Greece	2	1.0	0.05		
	Italy	1	1.1	0.075	40	
	Luxembourg	2		0.1		
	New Zealand	2-3	up to 1.0	0.037	1	(1)
		1		0.05	1	
	Spain	2	0.5-0.75	0.05-0.075	14	
	Switzerland	2	1.0	0.05	14	
(1) Dip, post-harvest						
Strawberry	Australia		0.5	0.05	1	
	Belgium	4-5	1.0		14	
	Canada		1.0	up to 0.1	1	
	Denmark	1-3	0.5-0.75		14	
	France	3-4	0.75-1.0	0.1	2-3	
	Germany	3	1.0	0.1	10F	
	Greece	4	0.75	0.075	7	
	Italy	3	0.75-1.0	0.075-0.1	21	
	Luxembourg	4-5	1.0		14	
	Netherlands	2	1.12-1.5	0.1	14G	
		2	1.12-1.5	0.1	4F	
	New Zealand	3	up to 1.0	0.05	1	
	Norway	3-4	1.5		14	
	Portugal	2	0.75	0.075	3	
	Spain	4	0.5-0.75	0.05-0.075	3F,G	(1)
		2-3	0.4-0.6		28F,G	(2)
Sweden	1-3	0.75	0.07	14		
UK	4	0.75	0.037	1G		
	4	0.75	0.037	1F		
USA	4	0.84-1.1	up to 0.12	0		
(1) Foliar spray						
(2) Dust						
Sugar beet	France	1	1.5 kg/t seed			st
	Greece	1	4.0 kg/t seed			st
Sunflower	France	2	0.7		-	(1)
(1) Second treatment at flower bud visible						
Tangelo	New Zealand			0.05	14	
Tomato	Australia	1-6	0.5	0.05	7	
	Belgium			0.05	3	
	Canada			0.05	2G	
	Denmark	1-3	0.5	0.05	3G	
					14F	
	France	2-3	0.75-1.0		3	
	Greece	2-3		0.05-0.075	7	

Crop	Country	Application			PHI, days	Remarks
		No.	Rate, kg ai/ha	Spray conc., kg ai/hl		
	Italy	3-4	0.5-0.75	0.05-0.075	21G	
		1-3			F	
	Luxembourg	1-6		0.05	3	
	Netherlands	2-3	0.63-1.3	0.025-0.5		
	Norway	1-4	1.0		14	
	New Zealand			0.05	3G	
	Portugal	3	0.75	0.075	7	
	Switzerland	2	0.5-0.75	0.05	3	
	UK	6		0.05	1G	
		6		0.05	2F	
Tree nuts	Greece	2		0.05	14	
Turnip, stubble	UK	2	0.5	0.13-0.25	21	
Vegetables	France	1	2.5 g/kg seed	-		st
	Greece	3	0.05-0.075		7	
	Spain	2-4	0.4-0.6		28	
		4	0.75-1.1	0.05-0.075	3-21 F,G	(1)
	Sweden	1-3	0.5-0.75	0.15	14	
(1) For PHIs see individual crops or crop groups						
Wheat	Belgium	1	0.35-0.43		42	
	France	1-2	0.4			
	Germany	1	0.5-0.76	0.12-0.25	35	
		1	0.52-0.70/t seed		-	st
	Luxembourg	1	0.35-0.43		42	
	UK	1-2	0.38-0.5	0.1-0.25		(1)
(1) Last application up to growth stage Zadoks 71						
Witloof chicory (sprouts)	Belgium	1		0.3		(3)
	Luxembourg	1		0.3		(3)
	Netherlands	1	0.065 kg/t tubers			(1)
		1	0.004/m ²			(2)
(1) Tuber pre-storage treatment (2) Tuber treatment at planting (3) Drench before planting						

¹ st = seed treatment

² F = Field (outdoor)

³ G = Glasshouse

RESIDUES RESULTING FROM SUPERVISED TRIALS

The Meeting reviewed supervised trial data for fruits (citrus, pome and stone fruit, berries, grapes, bananas, kiwifruit), vegetables (brassica, bulb, leafy, legume, root and tuber vegetables, stalk and stem vegetables, cucumber, tomato, peppers), cereals, nuts, oilseed, and other seeds.

List of Tables in this section

Table

2	Residues of iprodione in citrus fruits.
3	Residues of iprodione in apples.
4	Residues of iprodione in apples, post-harvest treatment.
5	Residues of iprodione in pears.
6	Residues of iprodione in apricots.
7	Residues of iprodione in cherries.
8	Residues of iprodione in peach and nectarines.
9	Residues of iprodione in plums.
10	Residues of iprodione in currants and gooseberries.
11	Residues of iprodione in blackberries.
12	Residues of iprodione in raspberries.
13	Residues of iprodione in strawberries.
14	Residues of iprodione in grapes.
15	Residues of iprodione in bananas.
16	Residues of iprodione in kiwifruit.
17	Residues of iprodione in fennel.
18	Residues of iprodione in garlic.
19	Residues of iprodione in onions.
20	Residues of iprodione in broccoli.
21	Residues of iprodione in chinese cabbage.
22	Residues of iprodione in red cabbage.
23	Residues of iprodione in white cabbage.
24	Residues of iprodione in kohlrabi.
25	Residues of iprodione in cauliflower.
26	Residues of iprodione in cantaloupes.
27	Residues of iprodione in cucumbers.
28	Residues of iprodione in peppers.
29	Residues of iprodione in tomatoes.
30	Residues of iprodione in witloof chicory.
31	Residues of iprodione in lettuce.
32	Residues of iprodione in dandelion.
33	Residues of iprodione in beans.
34	Residues of iprodione in peas.
35	Residues of iprodione in carrots, post-harvest treatment.
36	Residues of iprodione in carrots, pre-harvest treatment.
37	Residues of iprodione in radishes.
38	Residues of iprodione in potatoes.
39	Residues of iprodione in seed potatoes.
40	Residues of iprodione in sugar beet.
41	Residues of iprodione in swedes and turnips.
42	Residues of iprodione in celery.
43	Residues of iprodione in cumin seed.
44	Residues of iprodione in barley and oats.

- 45 Residues of iprodione in wheat.
 46 Residues of iprodione in rice.
 47 Residues of iprodione in maize.
 48 Residues of iprodione in almonds.
 49 Residues of iprodione in oil-seed rape.
 50 Residues of iprodione in sunflower.
 51 Total residues of iprodione and its metabolites in bovine milk.
 52 Total residues of iprodione and its nonhydroxylated metabolites in bovine tissues.

Underlined residues in Tables 2-50 are from treatments according to GAP.

Crop residues

Citrus fruits (Table 2). Data were available from spraying trials in Japan, New Zealand and Israel.

In post-harvest trials in Italy fruits were dipped once for 20 seconds in a suspension of Rovral WP50. Residues were about 0.7-1 mg/kg and 0.4-0.6 mg/kg after, respectively, 10-60 days and 120-180 days of storage at -12°C.

Table 2. Residues of iprodione in citrus fruits.

New Zealand : Oranges
 Japan and Israel: Mandarins
 Italy : Lemons

Country/Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha			
New Zealand	Rovral Flo	2	1.3	0	0.65	152
		3	1.3	1	1.9	152
1989				3	1.8	
				7	1.4	
				14	1.1	
Japan	Rovral WP 50	3	2.5	7	1) 0.32	3
1981					14	0.34
				30	0.03	
				7	2) 20	
				14	13	
				30	2.6	
					3) 0.04	
	Rovral WP 50	3	2.5	8	1) 0.12	3
					15	0.2
				31	0.11	
				8	2) 23	
				15	23	
				31	15	

Country/Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha			
					3) 0.14	
Israel	Rovral WP 50	4	2.0	145	<0.05-0.33	61
1990						
1992		6	1.5	172	0.19, 0.34	258
		6	1.5	144	0.32, 0.38	
		8	1.5	109	0.34, 0.51	
Italy*	Rovral WP 50	1		10	<u>0.88</u>	11
1985				60	<u>0.72</u>	
				120	0.49	
				180	0.42	
		1		10	<u>1.02</u>	11
				60	<u>0.94</u>	
			120	0.58		
			180	0.46		

1) Pulp 2) Peel 3) Juice
 * Post-harvest dip

Pome fruit

Apple. Residues after spraying are summarized in Table 3 and after post-harvest dipping treatment and cool storage in Table 4.

Table 3. Residues of iprodione in apples.

Country/Year	Application			PHI, days	Residue, mg/kg	Ref.	
	Form	No	kg ai/ha				
Japan	Rovral WP 50	1	5	3	2.9	136	
1975				7	2.6		
				10	2.3		
			10	5	10	6.5	136
			10	5	20	5.8	
			10	5	30	3.8	
Japan	Rovral WP 50	1	5	3	2.2	135	
1975				7	1.8		
			1	5	10	0.38	135
			10	0.5	10	3.4	135
			10	0.5	20	2	
			10	0.5	30	1.7	
France	Calidan**	2	0.52	2	0.6	63	
1986		2	0.57	2	1.2		
					71*	1.2	

* PHI 3 days + 68 days cold storage

** Not registered for this use in France

Apples, post-harvest treatment. Post-harvest dipping trials were carried out in various countries and residues of iprodione were measured by gas chromatography on the day of treatment and after cool storage: see Table 4.

Table 4. Residues of iprodione in apples, post-harvest dip treatment, all single applications.

Country, Year	Application		Storage, days	Residue, mg/kg	Ref.
	Form	kg ai/hl			
France	Rovral WP 50*	0.1	198	2.4	50
1980		0.1	36	0.8	
		0.1	0	1.5	
France	Rovral WP 50*	0.1	0	4.8	112
1979			97	4.5	
			222	3	
				Pulp 1.3	113
				Pulp AC [†] 0.85	
				Peel 17.4	
France	Rovral WP 50*	0.05	181	2.1	144
1979		0.05	222	2.3	
		0.1	181	3.2	
		0.1	222	3.0	
		0.2	181	5.7	
		0.2	222	5.7	
France	Rovral Aqua Flo	0.1	0	0.8	204
1989			28	2.4	

Country, Year	Application		Storage, days	Residue, mg/kg	Ref.
	Form	kg ai/hl			
			153	2.2	
	500 g/l	0.1	0	0.78	
			28	1.7	
			153	1.2	
		0.1	0	0.23	
			32	1.3	
			159	1.0	
		0.1	0	0.84	
			32	0.82	
			159	1.6	
		0.1	0	1.3	
			28	0.74	
			153	1.9	
		0.1	0	1.0	
			32	0.69	
			159	1.1	
Italy	Rovral WP 50	0.05	10-12	<u>2.2</u>	11
1985			60	<u>1.4</u>	
			120	<u>0.67</u>	
			180	<u>0.61</u>	
		0.075	10-12	<u>2.4</u>	
			60	<u>2.0</u>	
			120	<u>1.2</u>	
			180	<u>1.0</u>	

*Not registered for this use

¹AC: pulp after cooking for 60 min. at 100°C

Pears. Residues from the spray treatment of pears in France and Italy are summarized in Table 5.

Table 5. Residues of iprodione in pears.

Country/Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Italy	Rovral WP 50	6	1.35	0.075	6	2.0	142
1979							
1980		7	1.35	0.075	21	3.6	146
					28	3.2	
		4	1.35	0.075	21	<u>2.4</u>	
					28	2.3	
1992		6	1.35	0.075	0	2	221
					10	1.1	
					20	<u>0.51</u>	
					30	0.39	
					40	0.33	
France 1983	Rovral WP 50*	1	0.5	0.25	10**	0.25	150

* Not registered for this use in France
 **10 days PHI plus 3 months cold storage

Stone fruit.

Apricots. Residues of iprodione in apricots were determined at one location in France during 1980 (Table 6).

Table 6. Residues of iprodione in apricots (France, 1980).

Application			PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha			
Rovral WP 50	1	0.75	10	0.3	114
	2	0.75	10	0.95	

Cherries. Residues of iprodione in cherries treated with various formulations of iprodione are shown in Table 7.

Table 7. Residues of iprodione in cherries.

Country/Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Canada	Rovral WP 50	6	0.7	0.025	0	4.0	29
1978					1	3.9	
					5	2.5	
					8	2.6	
Canada	Rovral WP 50	7	0.875		1	1.1, 1.4	170
1982		8	0.875		1	1.4, 6.5	
France	Calidan SC	1	0.146		8	0.11	203
1989		2	0.46		1	0.18	
		1	0.439		13	0.13	
		2	0.439		2	0.26	
France	Kidán SC	1	0.625		8	0.12	202
1989		2	0.625		1	0.30	
		1	0.75		13	0.32	
		2	0.75		2	0.66	

Peaches and nectarines. Residues of iprodione from trials in France, Canada, Australia, Italy, Japan and South Africa on peaches and in Australia and France on nectarines treated with various formulations are shown in Table 8.

Table 8. Residues of iprodione in peaches and nectarines.

Country, Year	Application	PHI, days	Residue, mg/kg	Ref.
---------------	-------------	-----------	----------------	------

iprodone

719

	Form	No	kg ai/ha	kg ai/hl			
Australia	Rovral WP 50	5	1.4	0.05	6	5.9*	237
1976		7	1.4	0.05	3	5.4	239
		3	1.4	0.05	4	1.7	239
Canada	Rovral WP 50	5	1.0		0	<u>2.6</u>	124
1974					3	2.0	
					7	2.2	
					14	2.2	
		5	1.0		0	<u>7.6</u>	
					4	9.0	
					7	10	
					14	8.5	
		5	1.0		0	<u>6.2</u>	
					3	4.1	
					7	3.4	
					14	2.9	
		4	1.0		33	1.7	
		5	1.0		0	<u>6.5</u>	
					3	4.6	
					7	4.9	
		4	1.0		40	1.5	
		7	1.0		0	<u>2.3</u>	
					3	2.0	
					7	1.3	
				14	1.3		
	6	1.0		11	1.8		
				12	1.1		
				15	2.0		
				23	0.9		
Canada	Rovral WP 50	6	0.8	0.75	7	0.82, 1.9	20
1977		5	0.8	0.75	7	0.67, 1.2	
		5	0.8	0.75	7	0.3, 0.86	
		5	0.8	0.75	7	0.59, 1.3	
		5	0.8	0.75	7	1.1, 1.5	
Canada	Rovral WP 50	2	0.75		20	0.76	47
1978		3	0.75		1	<u>3.8</u>	
		3	0.75		7	1.4	
		3	0.75		14	0.8	
		3	0.75		22	0.36	
		3	0.75		6	1.3	
		4	0.75		1	<u>2.2</u>	
					8	1.8	
					15	0.72	
					22	0.65	
		3	0.70		0	<u>2.7</u>	
					1	<u>2.9</u>	
					3	2.3	
				5	1.8, 2.8		
				7	2.5		
France	Calidan SC	2	0.92		7	1.2	63
1985		3	0.92		4	0.62	
France	Calidan SC	3	0.43		15	0.3	45

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
1987		3	0.43		3	0.85	
France	Calidan SC	4	0.74		0	1.5	46
1987		3	0.45		0	0.54	
		4	0.54		0	0.8	
		5	0.52		0	1.7	
		4	0.52		0	2.4	
France	Calidan SC	4	0.43		0	0.5	201
1988		4	0.43		0	0.8	
		4	0.78 x 3		0	1.0	
			0.43 x 1				
France	Calidan SC	3	0.43		18	0.31	205
1989		4	0.43		4	0.37	
		4	0.43		1	0.41	
France	Kidan SC	4	1.2		0	1.8	46
1987		3	0.78		0	1.1	
		4	0.93		0	1.2	
		5	0.9		0	0.91	
		4	0.9		0	2.7	
France	Rovral WP 50	2	0.75		16	0.34*	99
1977		2	1		16	0.53*	
		2	0.9		3	0.73*	
		2	0.9		13	1.7	
France	Rovral WP 50	3	0.75		15	0.69	45
1987		3	0.75		3	1.8	
France	Rovral WP 50	4	1.27		0	4.3	46
1987		3	0.78		0	1.3	
		4	0.93		0	0.81	
		5	0.9		0	1.6	
		4	0.9		0	2.8	
Italy 1990	Rovral WP 50	1	1.1		21	0.05	208
		1	1.1		14	0.15	
Italy	Rovral WP 50	2	1.35		0	3.6	220
1991					10	2.6	
					20	1.1	
					30	0.83	
					40	0.35	
Japan	Rovral WP 50	2	4.0		1	3.7	137
1975					3	2.1	
		3	4.0		1	4.6	
					3	2.9	
		2	3.0		1	6.3	137
					3	4.8	
		3	3.0		1	6.8	
					3	5.8	
South Africa	Rovral WP 50**	1	0.5		3	4.8	21
1978		1	0.5		10	2.6	
		1	0.5		14	5.1	
		2	0.5		3	8.7	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		1	1.0		3	7.5	
		1	1.0		10	7.9	
		1	1.0		14	7.8	
		2	1.0		3	21	

* Nectarines
 ** Not sold in South Africa

Plums. In Australia plums were harvested 7 days after the last of 4 applications of a 0.05% ai suspension of Rovral WP50 to run-off (2800 l/ha; Woods, 1976). Trials were also carried out in France (two treatments with Rovral WP50, Kidan or Rovral Aqua Flo at the rate of 0.75 kg ai/ha) and South Africa. See Table 9.

Table 9. Residues of iprodione in plums.

Country/Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Australia 1975	Rovral WP 50	4	1.4		7	2.2	238
France	Rovral WP 50	2	0.75		10	1.5	81
1979		2	0.75		16	0.45	
France	Rovral WP 50	3	0.5		13	4.2	123
1974		3	1.0		13	6.8	
		2	0.5		52	0.19	
		2	1.0		52	0.64	
France 1979	Rovral Aqua Flo* SC 500	2	0.75		10	1.3	81
		2	0.75		16	0.35	
France	Kidana SC 255	2	0.75		10	1.3	81
1979		2	0.75		16	0.65	
South	Rovral Flo SC 255	2		0.0375	0	1.2	4
Africa					1	1.7	
1982					2	1.2	
					3	1.6	
		2		0.05	0	0.55	4
					1	1.5	
					2	0.51	
					3	1.4	
		2		0.075	0	2.1	4
					1	2.8	
					2	1.4	
					3	3.3	
South	Rovral Flo SC 255	2		0.05	0	0.6	182
Africa					2	0.8	
1985					3	0.7	
		2		0.05	0	1.1	182
					2	1.6	
					3	1.6	

Country/Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
					5	2.3	
		2		0.05	0	1.8	182
					2	1.9	
					3	1.1	
					5	1.5	
		2		0.05	0	1.3	182
					2	1.7	
					3	1.3	
					5	1.6	

* Not yet registered for plums in France

Berries and other small fruits

Currants and gooseberries. Supervised trials with iprodione were carried out on currants in 1981-1984 in The Netherlands and on gooseberries in 1980 in the UK (Table 10).

Table 10. Residues of iprodione in currants and gooseberries.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Netherlands	Rovral	5	0.75	0.075	78	5.4	43
1984*	WP 50	5	0.75	0.075	78	3.21	
		9	0.75	0.075	8	15	
		9	0.75	0.075	8	9.9	
Netherlands	Rovral	2	3.75	0.075	14	20	36
1982*	WP 50						
Netherlands	Rovral	5	0.75	0.075	31	7.1	36
1981*	WP 50	6	0.75	0.075	10	12	
UK 1980**	Rovral WP50	3	0.75	0.035	6	0.34	159

* Currants
** Gooseberries

Blackberries. Iprodione residue trials were carried out 1986 in Oregon (USA) on blackberries with Rovral WP50. The rate of application was 1.1 kg ai/ha. Five applications were made with the first two in the bloom stage and the remainder at bi-weekly intervals thereafter with the last application on the day of harvest. The results are summarized in Table 11.

Table 11. Residues of iprodione in blackberries (USA, 1986).

Form	Application		PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha			
Rovral WP 50	5	1.1	0	<u>5.8</u>	69
	5	1.1	0	<u>11</u>	
	5	1.1	0	<u>15</u>	
	5	1.1	0	<u>22</u>	

Raspberries. Residue data from trials in Canada, France and the UK are given in Table 12.

Table 12. Residues of iprodione in raspberries.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Canada	Rovral WP 50	6	1.0		1	<u>7.9</u>	169
1981					3	5.5	
					7	4.8	
Canada		6	1.0		1	<u>17</u>	169
1981					3	14	
					7	7.1	
Canada	Rovral WP 50	5	1.0		1	<u>4.6, 31</u>	178
1982		5	1.0		1	<u>12, 30</u>	
France	Rovral WP 50	3	0.8	0.05	18	1.1	33
1974		3	0.53	0.033	18	0.85	
UK	Rovral WP 50	4	0.75	0.25	6	<u>1.2</u>	12
1977		5	0.75	0.25	8	<u>2.6</u>	
UK	Rovral WP 50	5	1.12	0.05	17	1.6	125
1974		3	1.12	0.05	15	2.0	
		4	1.12	0.05	0	7.9	
					1	6.0	
					3	5.0	
					7	2.8	
					14	2.1	
UK	Rovral Flo*	1	0.75		1	2.7	163
1981		4	0.75		1	13	
		4	0.75		1	8.3	
UK	Rovral Flo*	4	0.75		1	2.7	173
1982		4	0.75		1	9.6	
		4	0.75		3	1.5	
		4	0.75		3	7.7	
		4	0.75		7	<u>0.92</u>	
		4	0.75		7	<u>5.4</u>	

* No registration for this use in the UK

Strawberries. Trials were conducted in many countries to obtain information on residues in strawberries treated with iprodione. The results are summarized in Table 13.

Table 13. Residues of iprodione in strawberries.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
USA	Rovral WP 50	1	1.1		0	<u>0.5-6.5</u> (5)	245
1985/1986		2	1.1		0	<u>1.9-7.9</u> (5)	
		3	1.1		0	<u>1.9-7.5</u> (5)	
		4	1.1		0	<u>9.1</u>	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		5	1.1		0	0.4-6.7 (7)	
		6	1.1		0	0.5-9.9 (11)	
		7	1.1		0	0.4-8.5 (11)	
		8	1.1		0	0.5-7.2 (8)	
		9	1.1		0	0.5-5.0 (7)	
		10	1.1		0	0.3-6.6 (7)	
		11	1.1		0	0.4-4.8 (2)	
		12	1.1		0	0.6-6.0 (2)	
		17	1.1		0	0.4-1.9 (5)	
Belgium	Rovral WP 50	5	0.75	0.125	13-16	1.9, 4.9	
1974		5	1	0.167	13-16	2.5, 6.0	
Netherlands	Rovral WP 50	4	0.75	0.075	18	1.1	8
1974		5	0.75		0	26	
					7	14	
					14	0.67	
1978		3	1.0 G	0.1	12	3.3	
		3	1.0 G	0.1	14	8.1	
1979		4	1.0	0.1	15	5.3	
France	Rovral WP 50	3	0.75	0.075	10	0.2	245
1981		4	0.5	0.075	19	0.07	
1982		3	0.75	0.075	6	0.15	
1977		3	1.5	0.075	17	2.3	
		3	2.0	0.1	17	4.3	
1974		4	0.5	0.033	13	1.2, 2.5	
		4	0.75	0.05	13	1.9, 3.2	
		4	1.0	0.067	13	2.8, 5.6	
		3	0.75	0.375	33	0.53	
		4	0.75		10	1.75	
		4	0.75		20	0.35	
		3	1.0	0.5	33	1.1	
		4	1.0		10	2.2	
		4	1.0		20	0.44	
Germany	Rovral WP 50	3	0.94	0.0375	0	5.3	
1974					4	2.7	
					7	2.3	
					10	1.1	
					14	1.0	
					21	0.38	

iprodione

725

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		3	1.25	0.05	0	9.1	
					4	3.6	
					7	3.4	
					10	<u>2.0</u>	
					14	1.9	
					21	0.67	
Germany	Rovral WP 50	3	1.0	0.05	0	33	
1978					3	32	
					7	18	
					14	4.7	
		3	1.25	0.05	0	8.4	
					3	5.5	
					5	7.2	
					7	4.5	
					14	2.2	
					21	1.0	
		3	1.25	0.05	0	8.8	
					3	4.3	
					7	2.4	
					10	<u>2.3</u>	
Germany	Rovral WP 50	3	1.0	0.05	12	<u>0.13-0.31</u> (3)	
1976					16	0.14-0.23 (3)	
					22	0.14-0.17 (3)	
Germany	Rovral WP 50	3	1.25	0.05	0	34	
1977					3	13	
					7	7.5	
					10	<u>4.5</u>	
					14	0.75	
					21	0.68	
					30	0.23	
Germany	Rovral WP 50	3	1.25	0.05	0	9.0	
1978					3	5.7	
					5	5.2	
					7	3.8	
					14	1.4	
					21	0.5	
		3	1.25	0.05	0	4.0	
					3	4.5	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
					5	3.3	
					7	2.5	
					14	0.8	
					21	0.4	
Denmark	Rovral WP 50	3	0.37		32	0.21	
1980							
New Zealand	Rovral WP 50	2	0.35		1	2.0	
1979		2	0.65	0.022	3	4.1	
1985		5	0.75	0.05	4	2.5	
		3	0.7	0.05	1	<u>2.9</u>	
		3	0.7	0.05	1	<u>3.9</u>	
Spain	Rovral WP 50		1.5 ²	0.075	0	12	10
1986					3	6.9	
					7	4.9	
					14	2.9	
					21	1.3	
Spain	Rovral WP 50		0.75 ³	0.025	0	2.2	
1987					3	<u>1.6</u>	
					7	1.0	
					14	0.8	
					21	0.4	
Spain	Rovral WP50		0.75 ⁴		0	3.4	
1988					3	<u>2.0</u>	
					7	1.6	
					14	0.6	
					21	0.6	
UK	Rovral WP 50	3	1.1	0.1	36	1.7	245
1974		4	1.1	0.1	7	4.5	
		4	1.1	0.1	15	4.7	
		2	0.98	0.1	28	0.3	
		3	0.98	0.1	20	0.7	
1976		3	1.0		8	1.6	
		3	1.0		16	0.6	
UK		4	0.75		1	<u>1.9</u>	245
1980		4	0.75		1	<u>2.5</u>	
UK	Rovral Flo	4	0.74	0.035	2	1.5, 2.1	
1979		4	0.74	0.035	2	3.1, 5.3	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		4	0.75	0.035	1	2.3,4.2	
1980		4	0.75	0.036	1	<u>2.0,3.8</u>	
1981		4	0.75		1	<u>2.1,4.8</u>	
Canada	Rovral WP 50	4	1.1	0.1	0	3.1-3.5 (3)	6
1974					1	<u>2.8-3.0</u> (3)	
					3	1.8-2.4 (3)	
					7	1.4-1.6 (3)	
					10	0.75-0.95 (3)	
					14	0.85-1.1 (3)	
		3	1.1	0.1	16	0.9	
					18	0.83	
					30	0.23	
1982		6	1.0		1	<u>3.5,6.0</u>	
1983		1	1.0	0.15	0	0.94	
					1	<u>0.97</u>	
					3	0.99	
					7	0.56	
Italy	Rovral Flo	3	1.5	0.1	45	3.1	245
1982			3	1.5	0.1	51	4.6

G = Glasshouse

¹ Nos. of samples in ranges are in parentheses

² Application indoors, no. and time not reported

³ Application outdoors, March-April 1987, no. not reported

⁴ Application indoors, April-May 1988, no. not reported

Grapes. Trials were in France (48), Chile (2), Portugal (10), Spain (7), Italy (14), Germany (6), Morocco (1), Canada (7) and the USA (6). Results are summarized in Table 14.

Table 14. Residues of iprodione in grapes.

Country, Year	Application				PHI, days	Residue, mg/kg (no. of samples in ranges)	Ref.
	Form	No	kg ai/ha	kg ai/hl			
France 1979	Kidán	4	0.75		14	<u>4.4</u>	111
		4	0.75		15	<u>1.6</u>	
		4	0.75		22	2.4	
		4	0.75		27	2.4	
		4	0.75		29	2.8	
France	Kidán	4	0.75	0.5	17	<u>2.0</u>	49
1980		4	0.75	0.3	20	0.9	
		4	0.75	0.63	23	0.4	
		4	0.75	0.75	23	0.9	
		4	0.75	0.38	30	0.3	
		4	0.75	0.5	31	2.0	
		3	0.75	0.5	40	0.7	
France	Kidán	4	0.75		13	<u>4.9</u>	42
1984		4	0.75		23	3.8	
France 1977	Rovral WP 50	4	0.75	0.5	18	5.1	100
		4	0.75	0.75	22	4.7	
		4	0.75	0.75	28	4.7	
		4	0.75	0.25	36	3.3	
France 1979	Rovral WP 50	5	0.75	0.75	10	5.4	49
		4	0.75	0.38	14	<u>1.5</u>	139
		4	0.75	0.35	27	1.7	
		4	0.75		29	2.1	
		4	0.75	0.69	15	<u>1.9</u>	
		5	0.75	0.38	7	5.7	
		5	0.75	0.75	12	5.4	
5	0.75	0.75	15	3.6			
France 1975	Rovral WP 50	4	0.5	0.25	25	1.5	139
		4	0.5	0.25	27	3.4	
		4	0.75	0.38	25	3.1	
		4	0.75	0.38	27	6.3	
France 1974	Rovral WP 50	4	0.5-1		20	3.1-9.7 (3)	126
		4	0.5-1		22	3.2-7.8 (3)	
		4	0.5-1		27	2.2-6.3 (3)	
		4	0.5-1		30	1.9-5.1 (3)	
		4	0.75	0.375	0	8.4	
	Rovral WP 50				7	6.4	
					15	<u>4.8</u>	
					21	4.2	
	Rovral WP 50	4	0.5	0.05	29	0.45-0.75 (3)	126

iprodione

729

Country, Year	Application				PHI, days	Residue, mg/kg (no. of samples in ranges)	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		4	0.75	0.075	29	0.52-1.1 (3)	
		4	1.0	0.1	29	1.0 -1.8 (3)	
		3	0.75	0.38	13	<u>2.7</u>	
France	Rovral WP 80	2	0.75		14	<u>0.64*</u>	207
1989		2	0.75		14	<u>1.1*</u>	
France	Rovral Aqua Flo	4	0.75		14	<u>1.2</u>	111
1979		4	0.75		15	<u>1.9</u>	
		4	0.75		22	1.8	
		4	0.75		28	1.8	
		4	0.75		28	1.9	
France	Rovral Aqua Flo	4	0.75		8	0.46	219
1991		4	0.75		14	<u>1.2</u>	
Chile	Rovral WP 50	4	0.75	0.063	7	5.4	148
1982		5	0.67	0.063	55	3.0	
Portugal	Rovral WP 50	4	0.6	0.05	35	6.6	245
1974		4	0.6	0.05	28	2.3	
		4	0.9	0.075	28	5.0	
		4	0.9	0.075	35	10	
		4	1.2	0.1	28	8.0	
		4	1.2	0.1	35	10	
Portugal	Rovral WP 50	4	0.25	0.05	23	2.6	89
1975		4	0.25	0.05	30	1.7	
		4	0.5	0.1	23	3.4	
		4	0.5	0.1	30	5.9	
Spain	Rovral WP 50	4	0.5	0.17	11	2.4	127
1974		4	0.5	0.05	47	6.1	
		4	0.75	0.25	15	<u>4.2</u>	
		4	0.75	0.075	47	10.0	
Spain	Rovral WP 50	4	0.5	0.05	38	2.4	88
1975		4	0.75	0.075	38	4.6	
Spain	Rovral WP 50	1	0.9	0.075	0	7.5	10
1986					2	4.7	
					5	3.8	
					7	3.0	
					14	1.1	
					21	0.8	
Italy	Rovral WP 50	4	0.5	0.05	28	3.6	84
1974		4	0.5	0.033	38	0.65	
		4	0.75	0.075	28	8.2	
		4	0.75	0.05	38	1.9	
Italy	Rovral WP 50	4	0.5	0.033	8	3.5	90

iprodione

Country, Year	Application				PHI, days	Residue, mg/kg (no. of samples in ranges)	Ref.
	Form	No	kg ai/ha	kg ai/hl			
1975		4	0.75	0.05	8	5.5	
		4	0.5	0.033	35	0.8	
		4	0.75	0.05	35	2.0	
Italy	Rovral Flo	4	0.75	0.05	21	12	115
1980					28	6.9	
Italy	Rovral Flo	4	0.5	0.03	21	3.6	120
1981					28	2.9	
		2	0.75	0.05	21	3.9	120
					28	3.7	
Italy	Rovral Flo	4	0.75	0.075	21	6.6	38
1982		4	0.75	0.075	28	5.8	
Italy	Rovral WP 50	3	0.75		0	4.3*	206
1990					9	3.7*	
					20	3.6*	
					30	2.7*	
					40	2.1*	
Germany	Rovral WP 50	5	0.4	0.1	8	2.7	85
1974					31	3.4	
					62	1.4	
					71	0.55	
		4	0.5	0.1	0	3.9	85
					28	2.1	
					42	1.9	
					84	1.3	
		5	0.6	0.125	8	4.9	85
					31	2.9	
					62	2.4	
					71	2.1	
		4	0.75	0.15	0	5.8	85
					28	3.9	
					42	2.8	
					84	2.2	
Germany	Rovral WP 50	5	0.75	0.0375	0	2.1-7.3 (5)	101
1977					7	2.2-6.8 (5)	
					14	2.0-6.7 (5)	
					21	1.5-5.1 (5)	
					28	1.5-5.9 (5)	
Germany	Rovral WP 50	4	0.75		14	0.71-5.2 (6)	91
1975					42	0.37-2.7(6)	
					53	0.42-2.8(6)	
					70	0.6 -1.2(6)	
					75	2.8	

Country, Year	Application				PHI, days	Residue, mg/kg (no. of samples in ranges)	Ref.	
	Form	No	kg ai/ha	kg ai/hl				
Morocco 1984	Rovral WP 50	2	0.75		7	1.2	226	
Canada	Rovral WP 50	4	0.75		42	2.5	156	
1979		1	1.0		0	2.5		
						2	2.4	
						7	1.2	
						15	1.0	
						19	1.1	
Canada	Rovral WP 50	4	0.75		41	1.5	179	
1980/81		5	0.75		34	1.7, 3.0		
		5	0.75		48	2.3		
		5	0.75		39	1.4		
		5	0.75		45	2.2		
USA	Rovral WP 50	4	1.1		0	0.88, 4.7	55	
1988		4	1.1		0	2.7, 4.4		
		4	1.1		0	0.9, 4.1		
		4	1.1		0	1.5, 2.1		
		4	1.1		0	2.7, 4.6		
		4	1.1		0	0.12, 0.66		

* Table grapes

Tropical and sub-tropical fruit

Bananas. Results of supervised trials from the Philippines (foliar spray) and Spain (post-harvest treatment, one location) are summarized in Table 15.

Table 15. Residues of iprodione in bananas.

Country, Year	Application				Post-treatment period, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Philippines	Rovral WP 50	41	0.5	0.37	9	Peel: 0.15, 1.5 Pulp: <0.03-0.06	145
1980		9	0.5	1.7	21	Peel: <0.1 Pulp: <0.1	121
1981	Rovral Aqua Flo*	9	0.5	1.7	21	Peel: <0.1 Pulp: <0.1	121
Spain 1991	EC 100 g/l	1 ¹		0.05	0	whole fruit: 3.4 Peel: 7 Pulp: <0.1	10
						3	Peel: 6.8 Pulp: <0.1
	EC 100 g/l	1 ¹		0.03	0	whole fruit: 2.5 Peel: 5.1	10

iprodione

Country, Year	Application				Post-treatment period, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
						Pulp: <0.1	
					3	Peel: 5.1 Pulp: <0.1	
	EC 100 g/l	1 ¹		0.025	0	whole fruit: <u>1.7</u> Peel: 4 Pulp: <0.1	10

* Not registered for this use in the Philippines

¹ Post-harvest treatment

Kiwifruit. Residues of iprodione have been determined in kiwifruit treated one to eight times with Rovral at various sites in France and New Zealand (Table 16).

Table 16. Residues of iprodione in kiwifruit (pre-harvest treatment).

Country, Year	Application				PHI, days	Residue, mg/kg ¹	Ref.	
	Form	No	kg ai/ha	kg ai/hl				
New Zealand 1979/80	Rovral WP 50	3	0.73	0.038	74	0.1-0.29 (4)	154,	
		5	0.75		0	3.5		
					1	<u>3.9</u>	157,	
					4	3.4	231	
					8	2.2	18	
					15	0.12		
			5	0.75		1	<u>3.7</u>	
					3	4.5		
					14	0.23		
					22	1.8		
			5	0.75		0	3.2	18
					1	<u>2.6</u>		
					4	2.4		
					15	1.7		
				23	0.92			
1981/82	Rovral WP 50	8	0.75		0	4.7	18	
					3	3.4		
					7	4.2		

Country, Year	Application				PHI, days	Residue, mg/kg ¹	Ref.
	Form	No	kg ai/ha	kg ai/hl			
	Rovral Flo	8	0.75		0	4.7	18
					3	3.8	
					7	2.5	
	Rovral Flo	8	0.75		0	9.3	18
					3	3.7	
					7	2.9	
1991	Rovral Flo	3	0.75		1	0.28-4.5 (10)	41
1979/80	Rovral WP 50	5	0.75	0.038	1	Pulp: 0.3	213
					3	0.42	
					1	0.16	
					8	0.19	
					1	0.91	
					4	0.14	
France 1986/87	Kidan*	1	0.75		1 + a ²	2.3 [1.3] ²	193
					15 + a	2.4 [1.5]	
		1	0.75		2 + b	2.5 [1.4]	
					12 + b	0.55 [1.1]	
		1	0.75		2 + c	2.1 [1.6]	
					15 + c	1.9 [1.2]	
		2	0.75		1 + a	3.8 [2.0]	
		2	1.5		2 + b	4.7 [2.6]	
		2	0.75		2 + c	2.0 [3.0]	
		2	0.75		3 + d	1.4 [1.5]	
		2	0.75		3 + d	1.0 [1.1]	
					17 + d	0.55 [1.2]	

* Kidan = Rovral Flo

¹ Nos. of samples in ranges are in parentheses

² [] after storage at 0°C for a = 131 days; b = 175 days; c = 161 days; d = 112 days

Kiwifruit. Treatment by post-harvest dip or spray. In a post-harvest residue trial in Australia (1989), mature fruits were dipped in Rovral WP 50 at 500 and 1000 mg/l for 30 seconds before being packed in commercial kiwifruit trays with polyethylene liners and cool-stored at 20°C. Residue analyses on pulp and whole fruit did not show a consistent decrease with increased withholding period (0 to 14 days). The residue of iprodione at 500 mg/l in the whole fruit was less than 10 mg/kg, but at 1000 mg/l was up to 13 mg/kg. At both rates the residue in the pulp was 1 mg/kg (Anon., 1990).

In New Zealand (1988), post-harvest application of Rovral Flo by means of a "Singulator" spray apparatus at a rate of 1.5 ml/l resulted in a mean iprodione residue level of 1.36 mg/kg on a whole-fruit basis and 0.05 mg/kg in the pulp. Residue levels in the whole fruit were significantly lower, and those in the pulp somewhat lower, than those from post-harvest dipping (Heffernan, 1990).

Bulb vegetables

Fennel. Results of one supervised trial (three replicates) on fennel (bulbs) were reported to the Meeting by The Netherlands (see Table 17).

Table 17. Residues of iprodione in fennel bulbs (Netherlands, 1983).

Application			PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha			
WP 50	3	0.75	16	0.05	8
			24	<0.05	

Garlic. Residues were about 5 mg/kg in garlic harvested approximately nine months after pre-planting treatment with Rovral WP 50. They were almost wholly in the outer sheaths of the bulbs (Table 18).

Table 18. Residues of iprodione in garlic (France, 1975-6).

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/t	kg ai/hl			
Rovral WP 50	1	1.5		268	4.6 *	96
	1	1.5		275	≤ 0.1 **	
					15.3 ***	

* Whole bulb ** Cloves *** Sheaths

Onions. Analyses were carried out by GLC to determine residues of iprodione in onions after spray applications of Rovral formulations. Results from several countries are summarized in Table 19.

Table 19. Residues of iprodione in onions.

Country, Year	Application	PHI, days	Residue, mg/kg (no. of samples)	Ref.
------------------	-------------	--------------	------------------------------------	------

	Form	No	kg ai/ha	kg ai/hl			
France 1991	Rovral Aqua Flo	5	0.75		21	<0.02	215
Netherlands	Rovral WP 50	2	0.25	0.17	36	≤0.05	107
1979							
Denmark	Rovral WP 50	1	until run-off	0.05	0	3.0	14
1980					7	0.14	
						14	0.25
Canada	Rovral WP 50	2	1.0	0.06	13	<u>0.05</u>	17
1978, 1979		3	0.75	0.23	19	<u>0.05</u>	31
		3	0.75	0.23	19	<u>0.07</u>	
1980		5	0.75		15	<u>0.18</u>	160
		5	0.75		15	< <u>0.1</u>	
		5	0.75		19	< <u>0.1</u>	
USA 1985	Rovral WP 50	4	1.1		4	<0.05	74
1989	Rovral 4F	5	0.75		7	< <u>0.05</u> (3)	53
		5	0.75		7	<u>0.06, 0.11</u>	

Brassica vegetables, head cabbage, flowerhead brassicas, broccoli. In the USA, Rovral WP50 was applied once or twice (at thinning and on the day of harvest) to broccoli. See Table 20.

Table 20. Residues of iprodione in broccoli (USA).

Year	Application				PHI, days	Residue, mg/kg (no of samples)	Ref.
	Form	No	kg ai/ha	kg ai/hl			
1983	Rovral WP 50	1	1.1		49	0.05	70
1984		2	1.1		0	<u>4.1-9.1</u> (5)	58
		2	1.1		0	<u>14</u>	
		2	1.1		0	<u>14</u>	
		2	1.1		0	<u>22</u>	

Chinese cabbage. Table 21. Residue trials on Chinese cabbage were conducted in the USA in 1984 and 1986. From two to five post-emergence applications were made with Rovral WP50, at rates of 0.5 and 1.1 kg ai/ha/application.

In Denmark, at one site during 1980, Chinese cabbage was sprayed once to run-off with an aqueous suspension of Rovral WP50 containing 0.5 g ai/l. A trial was carried out in Germany (1978) with four treatments at 0.5 kg ai/ha.

Table 21. Residues of iprodione in Chinese cabbage.

Country, Year	Application				PHI, days	Residue, mg/kg (No. of samples)	Ref.
	Form	No	kg ai/ha	kg ai/hl			

Country, Year	Application				PHI, days	Residue, mg/kg (No. of samples)	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Canada	Rovral	3	0.75	0.14	0	9.8	6
1990	WP 50				1	17	
					3	12	
					7	2.5	
					10	3.0	
					15	0.27	
					21	0.16	
Germany	Rovral	4	0.5	0.083	0	0.18-8.4 (6)	106
1978	WP 50*				3	0.23-3.8 (6)	
					5	≤0.1-0.5 (6)	
					7	≤0.1-3.4 (6)	
					14	0.04-2.8 (6)	
					21	≤0.03-4.9 (6)	
Denmark	Rovral	1	until	0.5	0	22	13
1980	WP 50		Run-off		7	1.1	
					14	0.23	
USA	Rovral	4	0.5		0	0.42	75
1984	WP 50**	4	0.5		0	1.83	
		2	1.1		18	0.3	
		4	1.1		0	1.1	
		4	1.1		0	5.7	
1986		5	1.1		0	5.0, 12	67
		5	1.1		0	9.7	
		5	1.1		0	11	
		5	1.1		0	12	

* Not registered for this use ** Registration expected for this use in the USA

Red cabbage. In trials in the UK at one site (1978-1979) a single post-harvest treatment with Rovral WP50 as a 0.1% w/v ai drench was followed by cold storage for 90 days. See Table 22.

Table 22. Residues of iprodione in red cabbage (UK, 1978-9).

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	ai concn.				
Rovral WP 50	1*	0.1% w/v (drench)		90	0.94, 3.1	28

White cabbage. Iprodione residues from Canadian and German spraying trials and UK post-harvest treatments are shown in Table 23.

Table 23. Residues of iprodione in white cabbage.

Country, Year	Application	PHI or storage period, days	Residue, mg/kg	Ref.
------------------	-------------	--------------------------------	----------------	------

	Form	No	kg ai/ha	kg ai/hl			
Canada	Rovral WP 50	3	0.75	0.094	0	0.30	6
1990					2	0.37	
					7	0.03	
					9	0.05	
Germany	Rovral WP 50	2		0.15	0	0.9	147
1979/80					7	1.1	
					10	0.95	
					21	1.3	
					50	1.2	
UK	Rovral WP 50	1*	5% dust		188	0.85	240
1976		1*	50% dust		188	2.0	
		1*	0.1% dip		188	5.0	
1977	Rovral WP 50	1*	10% w/v spray		103	0.47	242
					118	0.24	
		1*	0.1% w/v spray		103	0.1	
					118	0.06	
		1*	0.05% w/v spray		214	0.05	
1977/78	Rovral WP 50	1*	0.05% drench		220	5.9	16
		1*	0.05% dip		217	0.86	

* Post-harvest treatment

Kohlrabi. The results of five supervised trials (two locations) in The Netherlands in 1978 using Rovral WP 50 as a pre-harvest treatment are summarized in Table 24.

Table 24. Residues of iprodione in kohlrabi (Netherlands, 1978).

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha	kg ai/hl			
Rovral WP 50	1	2.0	0.33	14	0.15	104
	1	2.0	0.4	60	0.1	
	2	0.75	0.075	103	0.05	
	1	1.0	0.1	117	0.1	
	1	2.0	0.2	117	0.1	

Cauliflower. The results of residue trials in France and Canada during 1986 to 1991 are summarized in Table 25.

Table 25. Residues of iprodione in cauliflower.

Country, Year	Application	PHI, days	Residue, mg/kg	Ref.
---------------	-------------	-----------	----------------	------

iprodione

	Form	No	kg ai/ha	kg ai/hl			
Canada	Rovral WP 50	3	0.75	0.094	1	4.15	6
1991					3	4.55	
					7	1.8	
					10	1.7	
					15	1.0	
1990		3	0.75	0.094	0	2.3	6
					2	1.7	
					7	0.68	
					9	0.56	
France	Kidan*	1	0.75		57	0.02	227
1986		2	1.5		14	0.04	
		2	1.25		55	0.16	
1987		1	0.75		85	<0.05	195
		2	0.75		42	<0.05	
1991	Rovral Aqua Flo	2	0.75		26	0.02	224

* Not registered for this use in France

Fruiting vegetables, cucurbits

Cantaloupes. Residues of iprodione in cantaloupes treated with a 500 g/l suspension concentrate formulation of Rovral Aqua Flo are shown in Table 26.

Table 26. Residues of iprodione in cantaloupes (France, 1991).

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha	kg ai/hl			
Rovral	3	0.75		10	0.14	216
Aqua	2	0.75		21	<0.02	
Flo	3	0.75		0	0.04	218
				4	<0.02	
				7	0.03	
				13	0.02	

Cucumbers. The results of trials in France, the UK, Denmark, Canada and Japan are shown in Table 27.

Table 27. Residues of iprodione in cucumbers.

Country, Year	Application	PHI, days	Residue, mg/kg	Ref.
---------------	-------------	-----------	----------------	------

	Form	No	kg ai/ha	kg ai/hl			
France	Rovral WP 50	1	0.6	0.75	5	0.15	105
1978							
1991	Kidan	1	0.93		3	1.8	214
					8	1.8	
UK	Rovral WP 50	3	1.1		37	0.02, 0.07	22
1977		4	1.1		2	0.06, 0.23	
		11	1.1		9	<0.02	
		12	1.1		2	0.49	
Denmark	Rovral WP 50	1	until	0.05	0	0.68	19
1980			run-off		4	0.2	
					7	0.27	
1981		1	until	0.075	1	0.50	168
			run-off		2	0.51	
					4	0.44	
					7	0.22	
Canada	Rovral WP 50	3		0.05	1	<0.1	171
1982		9		0.05	1	0.18, 0.28	
		3		0.05	1	0.58	
Japan	Rovral WP 50	1	3.0	0.1	1	1.7	92
1975					3	1.0	
					7	0.36	
		3	3.0	0.1	1	1.6	
					3	0.93	
					7	0.67	
		4	3.0	0.1	1	2.2	
					3	1.5	
					7	1.2	
		1	2.5	0.1	1	1.2	
					3	1.0	
					7	0.25	
		3	2.5	0.1	1	2.0	
					3	1.4	
					7	0.29	
		4	2.5	0.1	1	1.8	
					3	0.98	
					7	1.0	

Peppers. Peppers in California, Florida, New Jersey, North Carolina and Texas were treated 8 times with Rovral 4F (480 g/l SC) at 0.85 or 1.7 kg ai/ha. See Table 28.

Table 28. Residues of iprodione in peppers from ground foliar treatments (USA, 1990).

Application			PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha			

iprodione

Application			PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha			
Rovral	8	0.85	0	0.55-2.4*	56
4 F	8	1.7	0	0.62-3.9*	
	8	0.85	0	0.49-4.9**	
	8	1.7	0	0.47-4.8**	

* Sweet pepper (7 trials) ** Hot pepper (6 trials)

Tomatoes. Residue trials on tomatoes were conducted in several countries with Rovral WP 50, Rovral Flo or Rovral Aqua Flo. Results are summarized in Table 29.

Table 29. Residues of iprodione in tomatoes.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
UK 1974	Rovral WP 50	5*	until run-off	0.05	0	3.1	83
					2	3.4	
					5	3.0	
					7	2.5	
					9	2.1	
					14	2.7	
					14	1.7	
					14	2.3	
					14	2.9	
					14	3.7	
					14	4.2	
					28	5.8	
					14	2.7	
					14	3.8	
	14	5.0					
	14	4.2					
			27	2.3			
			41	3.3			
1977	Rovral WP 50	2*	1.1		2	0.24, 1.0	243
		4*	1.1		2	1.4	
		2*	0.5		2	0.1	
		4*	0.5		3	0.23	
		4*	0.5		2	1.4	
		5*	0.5		3	0.28	
France	Rovral WP 50	1*	0.82	0.75	9	0.19	102

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
1978		4*	0.75	0.15	7	0.85	
France	Rovral Aqua Flo	3	2.2		0	0.1	212
1991					4	<0.05	
					7	<0.05	
					14	<0.05	
		2	0.75		19	<0.05	213
Japan	Rovral	3	3.0	0.1	1	2.1	134
1975	WP 50				3	3.4	
					7	1.8	
					14	3.0	
		4	3.0	0.1	1	4.6	
					3	3.6	
					7	4.1	
					14	2.8	
		5	3.0	0.1	1	4.5	
					3	4.1	
					7	3.9	
					14	3.8	
		1*	2.5	0.1	1	1.3	132
					3	1.4	
					7	1.2	
					14	0.79	
		3*	2.5	0.1	1	5.3	
					3	3.5	
					7	3.0	
					14	2.4	
		4*	2.5	0.1	1	5.6	
					3	5.4	
					7	4.3	
					14	3.5	
Italy	Rovral Flo	3	0.75	0.075	15	0.06	38
1982		3	0.75	0.075	28	0.03	
Denmark	Rovral WP 50*	1*	until run-off	0.05	0	1.3	15
1980					4	0.74	
						7	0.49
1981		7*	1.35		1	1.9	166
					3	1.7	
					5	0.9	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
					7	1.0	
					14	1.0	
Canada	Rovral WP 50	3*		0.05	2	0.15	6
1981/82					7	0.41	
		3*		0.1	2	0.54	
					7	0.47	
		2*	until run-off	0.05	3	<0.1, 0.39	
USA	Rovral WP 50	5	1.1		0	0.22-1.6	68
1986		5	2.2		0	0.46-2.8	
		3	1.1		0	<0.05-0.6	52
		3	1.1		0	2.2	

* Greenhouse

Leafy vegetables

Trials on witloof chicory, lettuce and dandelions are summarized in Tables 30-32.

Table 30. Residues of iprodione in witloof chicory.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Belgium 1975	Rovral WP 50	1	30 ***		48	leaf: 0.41 root: 1.5	153
					59	leaf: 0.09 root: 2.7	
					93	leaf: 0.77 root: 6.2	
					104	leaf: 0.55 root: 2.7	
Belgium 1975	Rovral WP 50	1	60 ***		48	leaf: 0.36 root: 3.1	153
					59	leaf: 0.21 root: 4.4	
					93	leaf: 0.59 root: 10.0	
					104	leaf: 1.0 root: 3.7	
France 1987	Rovral WP 50	2	40 ***		24	leaf: 1.0 root: 3.0	194
		2	40 ***		24	leaf: 0.8 root: 5.1	
1987		3	0.6 *	0.12	21	leaf: <0.05 root: 2.2	196
			0.3 **	0.06			
			40 ***	8.0			
		3	1.25 *	0.25	21	leaf: <0.05 root: 1.9	
			0.3 **	0.06			
			40 ***	8.0			

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
1992	Rovral Aqua Flo	1	40 ***		20	leaf: 0.03 root: 0.73	225
1992		1	40 ***		21	leaf: 0.08 root: 3.8	
1992		1	40 ***		21	leaf: 0.03 root: 0.89	
		1		0.06 **	28	leaf: 0.03 root: 1.0	
		2	0.06 **		21	leaf: <0.02 root: 2.2	
			40 ***				
Netherlands	Rovral WP 50	1	40 ****		25	leaf: <0.1	140
1977					25	root: 13 not washed	
						2.8 washed	
1977		1	40 ****		62	leaf: <0.1 root: <0.1 not washed & washed	

* Foliar spray ** Dipping
 *** Stem base drenching **** Spray after roots arranged for forcing

Table 31. Residues of iprodione in lettuce. All trials with Rovral WP 50.

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
Head lettuce						
UK	2	0.25		7	4.0, 4.3	26
1978/79	2	0.25		7	0.59, 0.94	
	3	0.25		13	1.0, 2.0	27
	4	0.25		7	7.2, 9.1	
	5	0.25		7	3.8	
1977	5	0.25		7	<0.2	241
	5	0.25		14	2.7, 3.3	
	6	0.25		6	7.7	
1974/75	5*	0.5	0.05	37	H 0.4	86
					OL 20	
					WL 13	
	3*	0.56	0.05	19	4.9	34
	3*	0.56	0.05	22	8.8	
	4*	0.56	0.05	0	19, 59	
				1	16, 40	
				7	14, 41	
				14	21	
Japan	2*	3.0	0.1	7	4.6	130
1975				21	0.05	
	3*	3.0	0.1	7	11	
				21	0.1	
	4*	3.0	0.1	7	7.4	
				21	0.11	
	2*	1.5	0.1	7	0.48	133
				21	0.39	
	3*	1.5	0.1	7	1.9	

iprodione

Country, Year				PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
				21	1.2	
	4*	1.5	0.1	7	2.3	
				21	1.7	
Netherlands	1*	7.5 (soil)		54	0.09	8
	1*	7.5 (soil)		67	0.08	
1975	2	0.75		10	<u>9.2</u>	
				17	<u>3.3</u>	
				24	<u>0.05</u>	
				32	<u><0.05</u>	
	2	0.75		21	<u>8.2</u>	93
				28	<u>4.9</u>	
				35	<u>1.8</u>	
				42	<u>0.65</u>	
France	4*	0.75	0.075	30	0.6	86
1977/78	4	0.75	0.075	36	0.06	
	4	0.9	0.075	13	4.3	250
	3*	0.75	0.075	23	<u>4.7</u>	
	4	0.75	0.094	26	0.07	
	3	0.75	0.15	26	≤0.1	
	3*	0.75	0.075	52	0.67	
1976	3*	0.75	0.075	39	6.7	95
1975	7	0.5	0.05	45	0.16	87
	7	0.75	0.075	45	0.29	
1974/75	4	0.75	0.05	53	WL 0.75	251
					OL 2.5	
					H 0.08	
1974	6	0.5	0.05	31	0.03	35
	6	0.75	0.075	31	0.03	
	6	1.0	0.1	31	0.03	
Germany	3*	0.3	0.05	0	28	98
1977				7	8.0	
				14	2.4	
				21	<u>1.4</u>	
				28	0.8	
	3*	0.5	0.05	0	48	98
				7	18	
				14	8.6	
				21	<u>3.7</u>	
				28	1.6	
1978	1*	1.5	0.25	0	96	103
				14	15	
				21	6.8	
				28	3.6	
				35	2.0	
				42	0.3	
	3*	0.5	0.083	0	32	103
				7	22	
				14	21	
				21	<u>6.9</u>	
				28	3.2	
1976	2*	0.5	0.05	32	0.17	97

iprodione

745

Country, Year				PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
1977	3*	0.3	0.05	0	28, 32	97
				7	8.0, 8.7	
				14	1.9, 2.4	
				21	<u>0.47, 1.4</u>	
				28	0.08, 0.78	
1977	3*	0.5	0.083	0	38, 48	97
				7	5.9, 18	
				14	2.3, 8.6	
				21	<u>0.75, 3.7</u>	
				28	0.55, 1.6	
1976	3*	0.75	0.15	0	22, 46	97
				5	14, 6.3	
				7	4.8, 11	
				10	2.2, 8.0	
				14	3.6, 7.2	
1976	3	0.75	0.13	0	16	97
				7	1.8	
				14	0.6	
				21	≤0.05	
1980	1*	1.0	0.17	0	88	117
				7	60	
				14	43	
				21	36	
				28	4.5	
1976	3	0.75	0.13	0	16	97
				7	1.8	
				14	0.6	
				21	≤0.05	
				28	≤0.05	
1976	3	0.75	0.15	0	13, 18	97
				5	4.3, 5.5	
				7	3.0, 4.2	
				10	2.3, 3.0	
1977	3	0.5	0.083	0	6.9	97
				7	0.66	
				14	0.25	
				21	<u><0.03</u>	
1979	1*	1.0	0.17	0	141	109
				14	11	
				28	0.55	
				35	0.3	
				42	0.05	
1979	1*	1.0	0.25	18	1.8	109
				25	4.1	
				32	3.7	
				39	2.4	
				46	2.1	
1978	1*	1.5	0.25	0	79	103
				14	8.4	
				21	3.2	

iprodione

Country, Year				PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
				28	0.9	
1978	1*	1.5	0.25	0	122	103
				14	25	
				21	6.4	
				28	≤0.1	
1979	1	1.0	0.17	0	70	109
				14	30	
				21	23	
				28	3.2	
Germany	1*	1.0	0.17	7	32	7
1980		autumn -		14	15	
		winter		21	<0.1	
				28	3	
				63	<0.1	
	1*	1.0	0.17	7	81	
		autumn -		14	60	
		winter		21	23	
				29	17	
				48	6	
	1*	1.0	0.17	7	0.8	
		autumn		14	0.2	
				21	<0.1	
				28	<0.1	
	1*	1.0	0.17	14	1.5	
		autumn		21	<0.1	
				28	0.1	
				48	<0.1	
	1*	1.0	0.17	14	41	
		spring		21	8	
				28	1.2	
	1*	1.0	0.17	14	2.1	
		summer		22	<0.1	
				28	<0.1	
				35	<0.1	
	1*	1.0	0.25	14	2.5	
		autumn		21	0.5	
				28	0.1	
				35	<0.1	
Spain		1.5 [†]	0.075	0	51 (3)	10
1986				3	42 (3)	
				7	35 (3)	
				14	25 (3)	
				21	10 (3)	
Spain	1	0.75		0	5.4 (3)	
1989				7	0.5 (3)	
				22	0.1 (3)	
Leaf lettuce						
Canada	3	0.75	0.136	0	14	6
1987				1	5.8	
				3	4.1	
				7	0.86	

iprodione

747

Country, Year				PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
				10	0.39	
				14	0.19	
				21	0.13	
Canada 1982	3	0.75	0.136	0	31	
Cos lettuce				1	11	
				3	9.4	
				7	4.7	
				10	2.2	
				14	1.0	
Canada 1982	3	0.75	0.136	0	42	
Leaf Lettuce				1	16	
				3	17	
				7	4.2	
				10	1.7	
				14	1.2	
Canada	3	0.75	0.136	0	25	
1982				1	12	
				3	5.9	
				7	4.6	
				10	2.8	
				14	1.5	
Spain		7.0 ⁺	0.25	0	7.4 (3)	10
1987				3	4.9 (3)	
				7	3.8 (3)	
				14	1.9 (3)	
				21	1.7 (3)	
				28	0.9 (3)	
Spain	1	0.7	0.035	0	5.0 (3)	
1988				3	3.4 (3)	
				7	0.6 (3)	
				14	0.5 (3)	
				21	0.1 (3)	
Denmark	1	run-off	0.05	0	19	13
1980				7	1.7	
				14	<u>0.44</u>	
1981	1*	0.25		1	4.8	167
				3	4.3	
				5	4.9	
				7	3.6	
				14	<u>1.1</u>	
USA	3	1.1		5	11	77
1985/86	3	1.1		10	7.9	
Bibb Lettuce	3	1.1		10	1.1	
	3	1.1		14	<u>4.7</u>	
	3	1.1		14	<u>14</u>	
	3	1.1		14	<u>9.0</u>	
	3	1.1		14	<u>4.9</u>	
USA	3	1.1		10	3.8	77
1984/ 85/86	3	1.1		10	1.7	
	3	1.1		14	<u>20</u>	
Romaine Lettuce	3	1.1		14	<u>0.18</u>	

iprodione

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
	3	1.1		14	<u>7.4</u>	
	3	1.1		14	<u>0.07</u>	
	3	1.1		14	<u>5.4</u>	
	3	1.1		14	<u>5.3</u>	
	3	1.1		14	<u>1.8</u>	
USA	3	1.1		10	1.4	77
1985/86	3	1.1		13	17	
Escarole Lettuce	3	1.1		14	<u>0.16</u>	
	3	1.1		14	<u>22</u>	
	3	1.1		14	<u>0.17</u>	
	3	1.1		14	<u>17</u>	
	3	1.1		14	<u>16</u>	
	3	1.1		14	<u>2.6</u>	
	3	1.1		14	<u>0.32</u>	

* Greenhouse lettuce; H Hearts; OL Outer leaves; WL Whole lettuce

¹ Application outdoor March-April, no. not reported

(3) Mean of 3 replicates

Table 32. Residues of iprodione in dandelions (France, 1991).

Application			PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha			
Rovral	1	5	15	2.1	209

Legume vegetables

Beans. Data on residues in succulent and dried beans are summarized in Table 33.

Table 33. Residues of iprodione in beans.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Succulent beans							
Canada	Rovral WP 50	1	0.75		27	SB <u><0.1</u>	177
1982		2	0.75		14	SB <u>0.15</u>	
		2	0.75		15	SB <u>0.21</u>	
		2	0.75		15	SB <u>0.33</u>	
Netherlands	Rovral WP 50	2	1.1	0.075	2	1.5	108
1979					6	1.5	
					9	0.6	
		2	1.1	0.075	8	0.85	108
					12	0.85	
					15	0.8	

iprodione

749

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
France	Rovral WP 50	2	0.75	0.075	6	FB 0.71	141
1978		2	0.75	0.075	11	FB 0.49	
1978	Rovral WP 50	1	0.75	0.075	7	FB 0.1	143
		1	0.75	0.075	12	FB 0.3	
		2	0.75	0.075	11	FB 0.6	
1988		1	0.75		8	FBP <0.25	199
						W <0.1	
1979	Kidán*	1	0.75	0.075	7	FB 0.25	143
		1	0.75	0.075	12	FB 0.25	
		2	0.75	0.075	11	FB 0.6	
1981		3	0.75	0.075	6	FB 0.25	252
1982	Kidán*	3	0.75	0.075	4	FB 0.83	39
		2	0.75	0.15	17	FB 0.13	
		1	0.75	0.15	18	FB 0.22	
		2	0.75	0.15	14	FB 0.31	
1983	Kidán*	2	0.75	0.075	8	FB 0.74	149
1979	Kidán*	1	0.75	0.075	12	FB 0.25	143
1991	Rovral Aqua Flo	4	0.75		5	FB 0.49	217
		2	0.75		4	FB 0.28	
Germany	Rovral WP 50*	2	0.5	0.083	0	FB 1.8	103
1978					7	FB 0.6	
					14	FB 0.6	
		2	0.75	0.125	0	FB 1.9	
					7	FB 0.9	
					14	FB 0.6	
Germany	Rovral WP 50*	2	0.5	0.083	0	FB 2.3	253
1979					7	FB 0.2	
					14	FB 0.1	
					21	FB 0.05	
		2	0.5	0.125	0	FB 1.8	
					14	FB 0.4	
				21	FB 0.1		
Belgium	Quintal IC*	2	0.7		20	0.05	254
Spain	Rovral WP 50	1	1.125		0	FB 1.4	10
1990					3	1.1, 1.7, 1.3, 1.4	
					7	FB 1.4 ¹	
					10	FB 0.5 ¹	
					14	FB 0.4 ¹	
Portugal	Rovral WP 50	6		0.075	6	1.1	
1983 (G)		1		0.075	1	2.7	
					4	1.9	
					7	1.0	
		1		0.075	4	2.9	
		1		0.075	3	2.4	
		1		0.075	2	3.9	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.	
	Form	No	kg ai/ha	kg ai/hl				
USA	Rovral WP 50	2	1.1		21	LB 0.08	57	
1983							BF 0.89	
		2	1.1		3	SB <u>0.84</u>	57	
						BF 11		
		2	1.1**		9	SB <u>0.1</u>		
						BF 1.4		
		2	1.1		9	SB <u>0.32</u>		
						BF 13		
		2	1.1**		10	SB <u>0.07</u>		
						BF 2.9		
		2	1.1		15	SB <u>0.33</u>		
						BF 11		
		2	1.1**		15	SB <u>0.06</u>		
						BF 2.9		
		2	1.1		18	SB <u>0.11</u>		
						BF 11		
1984		2	1.1**		17	SB <u><0.05</u>	71	
1985		2	1.1		24	SB <u>0.58</u>	78	
						BF 75		
		2	1.1		14	SB <u>0.82</u>		
						BF 66		
		2	1.1		14	SB <u>0.7</u>		
						BF 48		
1985		2	1.1		14	LB <u>1.3</u>	78	
						BF 27		
		2	1.1		14	LB <u>1.2</u>		
						BF 28		
1992		2	1.1**		15	SB <u><0.05</u>	256	
Dried beans								
Canada	Rovral WP 50	1	0.85		60	WB <u><0.02</u>	23	
1977						76	WB <u><0.02</u>	
		1	1.1		60	WB <u>0.06</u>		
					76	WB <u><0.02</u>		
1979		1	0.5		39	WB ≤ 0.1	32	
					41	WB ≤ 0.1		
1982		2	0.75		50	WB <u><0.1</u>	175	
					53	WB <u><0.1</u>		
		2	0.75		50	KB <u><0.1</u>	176	
					68	KB <u><0.1</u>		
Japan	Rovral WP 50	1	1.0	0.1	14	KB <u><0.05</u>	138	
1975						28	KB <u><0.05</u>	
		3	1.0	0.1	21	KB <u><0.05</u>	131	
					28	KB <u><0.05</u>		
UK 1983	Rovral Flo	2	0.5		28	<0.1	186	
1984			2	0.5		66	<0.1	188
			2	0.5		66	<0.1	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		2	0.5		73	<0.1	

BF Bean forage; LB Lima beans
 FB French beans; SB Snap beans
 FBP Processed French beans; WB White beans
 KB Kidney beans; W Water
 (G) Glasshouse application
 * Not registered for this use; ** Aerial application
¹ Mean of four replicates

Peas. Residues in fresh and dried peas are shown in Table 34. Information on the portion of the commodity analysed (peas or peas plus pods) was available only from The Netherlands.

Table 34. Residues of iprodione in peas.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Fresh peas							
France	Kidan*	1	0.5	0.1	25	0.1	39
1982		1	0.75	0.15	25	0.13	
		2	0.5	0.1	18	0.16	
		2	0.75	0.15	18	0.48	
		2	0.5	0.1	10	0.28	
		2	0.75	0.15	10	0.39	
		2	0.75		14	0.2	190
1987		2	0.52		30	0.1	198
		2	0.52		32	0.1	
1991	Rovral Aqua Flo	3	1.5		7	<0.05	211
Belgium 1988	Quintal IC	2	0.7		63	2.2	228
Netherlands	Rovral Aqua Flo	3	0.75		3	0.53**	197
1987		3	0.75		3	0.71**	
Dried peas							
France 1986	Calidan	2	0.52		33	0.05	192

Calidan = Quintal IC : Iprodione 175 g/l + carbendazim 87.5 g/l
 * Not registered for this use in France
 ** Peas + pods

Root and tuber vegetables

Carrots, post-harvest treatment. Residues of iprodione were determined in stored carrots treated at three sites in The Netherlands (Table 35).

Table 35. Residues of iprodione in carrots, post-harvest treatment (Netherlands, 1983 and 1985).

Year	Application				Storage period, days	Residue, mg/kg	Ref.
	Form	No	g ai/t	kg ai/hl			
1983	Rovral Flo	1*	8		126	7.4	8
1985	Rovral Flo	1*	15		0	2.8	
					63	3.5	
					130	3.0	
		1*	30		0	4.5	
					63	7.9	
					130	6.7	
		1*	61		0	13	
					63	14	
					130	8.7	
1985	Rovral	1*	15		0	3.4	
	Flo				53	3.3	
					120	3.6	
		1*	30		0	4.5	
					53	4.2	
					120	8.0	
		1*	61		0	9.3	
					53	8.0	
					120	7.6	

* Post-harvest spray treatment

Carrots, pre-harvest treatment. Rovral WP50 was applied to carrot plots in the USA (1985) at two rates, 1.13 and 2.25 kg ai/ha. Eight or thirteen applications were made, starting with the onset of early leaf and continuing weekly until maturity. The last foliar spray was made on the day of harvest. For this study, a total of ten trials were conducted in Arizona (1), California (3), Florida (1), Michigan (1), New Jersey (1), Oregon (1) and Texas (2). The trials covered the various market seasons and agricultural practices. Additional data from one trial (Netherlands, 1993) were reported by The Netherlands (PHI 30-37 days). Spain reported two additional trials from New Zealand. Table 36 shows the results from all the trials.

Table 36. Residues of iprodione in carrots. Pre-harvest treatment.

Country, Year	Application	PHI, days	Residue, mg/kg (No. of samples)	Ref.
---------------	-------------	-----------	---------------------------------	------

	Form	No	kg ai/ha	kg ai/hl			
USA	Rovral WP 50	8	1.3		0	0.49-3.1 (9)	60
1984		8	2.3		0	0.77-7.2 (8)	
		13	1.3		0	1.3	
		13	2.3		0	2.2	
Netherlands	Rovral WP 50	7	0.75	0.075	30	1.4	8
1981					37	1.2	
New Zealand	EC	3	0.5		19	0.16, 0.23	9
1983	250 g/l*	3	0.75		19	0.30, 0.33	

* Not registered for this use in New Zealand

Radishes. Results of two supervised trials at two locations in The Netherlands using Rovral WP 50 as a pre-harvest treatment are summarized in Table 37.

Table 37. Residues of iprodione in radishes (Netherlands, 1978).

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha	kg ai/hl			
Rovral WP 50	1	2.0	0.33	21	0.2	8
					0.25	
						0.35
	1	2.0	0.2	12	2.0	
					1.8	
					2.2	

Potatoes. Residues of iprodione applied as either Rovral WP50 or Calidan have been measured in potatoes. After treatment of tubers at planting or spraying of plants, residues in the tubers at harvest were all less than 0.05 mg/kg (Table 38).

Seed potatoes. Residues of iprodione in potato tubers following post-harvest treatment with Rhapsodie at three sites in The Netherlands and of seed potatoes in the UK and France are shown in Table 39.

Table 38. Residues of iprodione in potatoes (USA).

Year	Application			PHI, days	Residue, mg/kg, in tubers	Ref.
	Form	No	kg ai/ha			
1985	Rovral WP 50	4F	1.1	6	<0.05	76
				12	<0.05	
				14	<0.05	
				30	<0.05	
		4F	2.2	6	<0.05	76
				12	<0.05	
				14	<0.05	
				30	<0.05	
1986		4F	5.5	14	0.17	64
1986		4F	11	14	0.32	64
1984		2	0.5	48	<0.05	73
				89	<0.05	

F Foliar spray

Table 39. Residues of iprodione in seed potatoes.

Country, Year	Application				Storage period or PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/t	kg ai/hl			
UK 1978	Rovral WP 50	1P	0.04-0.1		91-123	pulp <0.02 whole <0.02	25
France 1982	Calidan	1D		0.21	136	<0.05	40
1982		1D		0.21	169	0.05	
		1M		0.7	136	<0.05	
1976	Rovral WP 50	1M		0.7	169	0.1	
		1D*		0.2-0.3	162	culls <0.01 pulp <0.01	40
		1M*	0.05-0.1		162	culls <0.01 pulp <0.01	40
		1D**		0.2	34-70	peel 27 pulp 0.4 tuber 3.0	40
		1D**	0.2		34-70	peel 27 pulp 0.27 tuber 5.0	40
		1S**	0.05		34-70	peel 160 pulp 0.5 tuber 25	229
		1S**	0.05		34-70	peel 100 pulp 0.5 tuber 25	229
		1S**	0.1		34-70	peel 340 pulp 1.0 tuber 58	229
		1S**	0.1		34-70	peel 205 pulp 2.2 tuber 67	229
		1S**	0.15		34-70	peel 320 pulp 1.5 tuber 120	229

Country, Year	Application				Storage period or PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/t	kg ai/hl			
		1S**	0.15		34-70	peel 330 pulp 3.2 tuber 140	229
Netherlands	Rhapsodie 350 g/l	1	0.07		5	28	48
1987***		1	0.07		56	27	
		1	0.09		56	35	
		1	0.1		105	3.2	232
1990***		1	0.1		0	0.55, 16	210
					91	2.0	
					92	9.6	
					97	2.9, 8.4	

D* Dipping before planting P Pre-planting treatment
D** Dipping during storage S** Spraying during storage
D Dipping at planting SI Sprinkler irrigation system
M* Misting before planting *** treatment for storage
M Misting at planting

Sugar beet. Sugar beet was grown from seed treated with Rovral WP50 at a rate of 1.5 g iprodione/kg seed at three sites in the UK. Roots and tops from sugar beet harvested 22 or 26 weeks after drilling contained less than 0.1 mg/kg iprodione.

In France residues 42 days after the last of two foliar treatments with Sumistar were less than 0.05 mg/kg in the roots. The tops contained 0.8 mg/kg (Table 40).

Table 40. Residues of iprodione in sugar beet.

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	g ai/kg seed			
UK 1986	Rovral WP	1	1.5	154	R <0.1 T <0.1	185
		1	1.5	154	R <0.1 T <0.1	
		1	1.5	182	R <0.1 T <0.1	
France 1988	Sumistar*	2	0.4 kg ai/ha	42	R <0.05 T 0.81	200

R Roots * Not registered for this use in France
T Tops

Swedes and turnips. The data are from trials in the UK at 3 sites (Table 41).

Table 41. Residues of iprodione in swedes and turnips (UK).

Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha			
1985*	Rovral Flo	3	0.5	28	roots \leq 0.1 leaves 2.6	183
				28	roots \leq 0.1 leaves 2.9	
1986*		3	0.5	28	roots \leq 0.1 leaves 1.2	184
1984**	Rovral Flo	3	0.5	28	\leq 0.1	187
1985**		3	0.5	21	roots 0.7 leaves 32	183
		3	0.5	26	roots $<$ 0.1 leaves 1.8	

* Swedes ** Turnips

Stem vegetables

Celery. Trials in the USA with Rovral WP50 applied as a directed spray at weekly intervals from planting to harvest yielded the residues shown in Table 42.

Table 42. Residues of iprodione in celery (USA, 1984).

Form	Application		PHI, days	Residue, mg/kg (No. of samples)	Ref.
	No	kg ai/ha			
Rovral WP 50	11	1.1	0	8.7	65
	11	1.1	0	8.7	
	12	1.1	0	17	
	13	1.1	0	6.2-48 (4)	
	11	2.2	0	20, 25	
	12	2.2	0	31	
	13	2.2	0	7.5-69 (4)	

Cumin seed. Results of one trial reported by The Netherlands are summarized in Table 43.

Table 43. Residues of iprodione in cumin seed (Netherlands, 1985).

Form	Application			PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha	kg ai/hl			
Rovral Flo	2	0.5	0.125	55	1.5	8

Application				PHI, days	Residue, mg/kg	Ref.
Form	No	kg ai/ha	kg ai/hl			
				55	1.9	

Cereal grains

Barley and oats. Results from Germany and the UK are summarized in Table 44. In some trials iprodione was used as a seed dressing (Rovral TS).

Table 44. Residues of iprodione in barley and oats.

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha			
Barley						
UK 1981	Rovral Flo	2	0.5	65	G 0.19, 0.3	181
1981/82		3	0.5	52	G 0.56, 0.6	180
		3	0.5	64	G <0.1, 0.2 S 0.63, 0.8	
1987		3	0.5	41	G 1.3, 1.5 S 3.0	2
		3	0.5	49	G 0.6, 0.8 S 1.5, 1.8	
		3	0.5	51	G 0.4 S 1.4, 2.3	
1986	Sirocco	3	0.5	51	G <0.1 S 1.5, 2.2	1
		3	0.5	44	G 0.14	
					S 3.8, 5.5	
Germany 1980	Rovral TS *, **	1	53 g/100 kg seed	310	G ≤0.05 S ≤0.05	118
1979		1	53 g/100 kg seed	121	G ≤0.05 S ≤0.05	110
		1	70 g/100 kg seed	121	G ≤0.05 S ≤0.05	110
		1	53 g/100 kg seed	125	E ≤0.1 S ≤0.1	
		1	70 g/100 kg seed	302	G <0.1 S 0.12	110
Oats						
Germany 1979	Rovral TS *, **	1	70 g/100 kg seed	150	G 0.1 S ≤0.05	110

G Grain E Ear * Seed dressing
S Straw ** Not registered in Germany.

Wheat. Results of trials in The Netherlands (1986) after one foliar treatment of wheat at one site and in the UK after 3 treatments with Rovral Flo or Sirocco are summarized in Table 45. In Germany

residues in grain and associated straw samples from wheat grown after seed dressing with Rovral TS did not exceed 0.1 mg/kg.

Table 45. Residues of iprodione in wheat.

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha			
Netherlands 1986	Sumidione*	1	0.4		G 0.09 S 4.6	236
UK 1981	Rovral Flo	3	0.5	57	G <0.1, 0.1	180
1981/82		3	0.5	74	G <0.1 S 3.3, 3.9	180
		3	0.5	67	G <0.1 S 1.8, 4.7	180
		3	0.5	68	G 0.1	180
1987		3	0.5	42	G <0.1	2
				59	G <0.1	
				81	G 0.1	
1986	Sirocco*	3	0.5	50	G <0.1	1
				53	G <0.1	
Germany 1980	Rovral TS**	1	70 g/100 kg seed	307	G ≤0.01 S <0.05	118
		1	70 g/100 kg seed	336	G <0.1 S <0.05	
1979		1	53 g/100 kg seed	305	G <0.1 S <0.1	110
		1	70 g/100 kg seed	305	G <0.1 S <0.1	

G Grain * Not registered
S Straw ** Seed dressing

Rice. Residues of iprodione in grain and straw from rice treated with Rovral WP50 are shown in Table 46.

Table 46. Residues of iprodione in rice.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Japan 1975	Rovral WP 50	1	1.2	0.1	14	G* 0.1 S 16	94, 255
					21	G* 0.1 S 15	
		1	1.2	0.1	28	G* 0.1 S 10	
		3	1.2	0.1	21	G* 2.2 S 45	
		3	1.2	0.1	22	G+ 0.79 S 49	
		3	1.2	0.1	30	G* 1.5 S 32	
		4	1.2	0.1	22	G+ 1.9	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
						S 44	
		1	1.2	0.1	15	G+ 0.31 S 32	
					22	G+ 0.37 S 13	
						G* S	59
USA	Rovral WP 50	2	0.5		32	1.2 3.6	
1985		2	0.5		32	0.74 1.0	
Arkansas		2	0.5		33	7.1 7.5	
Louisiana		2	0.5		34	1.4 2.8	
Mississippi		2	0.5		35	2.0 2.5	
Texas		2	0.5		36	2.5 9.7	
		2	0.5		38	0.55 1.6	
		2	0.5		42	0.2 2.3	
		2	0.5		43	0.2 0.74	
		2	0.5		43	2.5 4.6	
		2	0.5		58	0.66 0.5	
		2	1.1		35	3.6 4.1	
		2	0.5		28	0.51	51
		2	0.5		28	0.59	
		2	0.5		28	0.54	
					47	0.09	
		2	0.5		28	0.92	
					47	0.89	
1985		2	0.5		35	G* 2.0 PR 0.14 H 8.9 B 5.2 S 2.5	59
		2	1.1		35	G* 3.6 PR 0.42 H 18 B 5.8 S 4.1	59
1982		2	1.1		40	H/S 9.9 S 36 H 6.2 M 8.3 B 4.5 BR 0.78 PR 0.26	230
Italy 1986	Rovral WP 50	1	100 g/100 kg		107	<0.05	234

B Bran-polish. Bran with a small amount of kernel
 BR Brown rice (kernel with bran). There is 1-4% bran in rice grain
 G+ Grain decorticated, not polished rice
 G* Grain, unpolished rice
 H/S Head/stalks
 H Hulls
 PR Polished rice
 M Millfeed: Hull with a small amount of kernel
 S Straw

Maize. In Italy maize grain was treated before sowing by dusting with Rovral WP50, at a rate of 100 g/100 kg seed. At harvest, 236 days after treatment, grain grown from the seeds contained less than 0.05 mg iprodione/kg (limit of detection).

In the USA maize grain was treated with Rovral WP50 at rates of 5, 10 and 20 mg/kg before

being placed in a storage silo. Table 47 shows the residues in grain during 84 days of storage.

Table 47. Residues of iprodione in maize.

Country, Year	Application			PHI, days	Residue, mg/kg	Ref.
	Form	No	g ai/100 kg seed			
Italy 1986	Rovral WP 50	1+	100	236	<0.05	235
USA 1986	Rovral WP 50*	1 [^]	0.5	0	6.2	66
				28	4.8	
				56	4.3	
				84	3.6	
		1 [^]	1.0	0	8.9	66
				28	8.6	
				56	4.6	
				84	4.9	
		1 [^]	2.0	0	17	66
				28	19	
				56	10	
				84	14	

- [^] Before storage
⁺ Before sowing
^{*} Not yet registered for this use in the USA.

Tree nuts

Almonds. Almond orchards in California were treated four times with Rovral WP50 at 0.5 kg ai/ha. In one trial the last spray was at 1.1 instead of 0.5 kg ai/ha. The interval between the last application and harvest ranged from 132 to 144 days. The results are summarized in Table 48.

Table 48. Residues of iprodione in almonds (USA, 1984).

Form	Application		PHI, days	Residue, mg/kg	Ref.
	No	kg ai/ha			
Rovral WP 50	4	0.5	139	Hulls 1.1	72
	4	0.5	144	Hulls 1.2	
	4	0.5	144	Hulls 0.6	
	4	0.5+1.1	132	Hulls 1.3	
	4	0.5	139	Nut <0.05	
	4	0.5	144	Nut <0.05	
	4	0.5	144	Nut <0.05	
	4	0.5+1.1	132	Nut 0.18	

Oilseed

Rape. Residues of iprodione in oil-seed rape treated with iprodione as either a 255 g/l SC or a 50% WP are shown in Table 49.

Table 49. Residues of iprodione, and carbendazim where applicable, in rape seed and straw.

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
Canada 1977	Rovral WP 50	1	1.0		59	seed <0.1	24
1980		1	0.5		40	seed <0.1	
		1	0.5		50	seed <0.1	161
France 1980	Rovral WP 50	2	0.75	0.15	48	seed <0.02	116
1981		1	0.75	0.15	77	seed <0.1	119
		1	0.75	0.16	87	seed <0.1	
		1	0.75	2.5	45	seed 0.15	
		1	0.38	0.075	64	seed <0.1	
1981	Kidan SC	1	0.75	0.15	77	seed <0.1	
		1	0.75	0.16	87	seed <0.1	
		1	0.38	0.075	64	seed <0.1	
1982		1	0.5	0.13	43	seed 0.13	37
		1	0.75	0.18	43	seed 0.13	37
1985	Calidan SC	1	0.61i 0.30c	0.20i 0.1c	61	seed 0.16i <0.05c	62
		1	0.54i 0.27c	0.11i 0.05c	60	seed 0.09i 0.09c	
		1	0.43i 0.21c	0.13i 0.06c	77	seed <0.02i <0.05c	
		1	0.52i 0.26c	0.1i 0.05c	63	seed 0.03i <0.05c	
Germany 1980	Rovral WP 50	3	0.75		43	seed <0.02	116
		3	0.75		55	seed <0.02 straw <0.05	
		2	0.75		77	seed <0.1 straw <0.1	
		2	0.75		95	seed <0.1 straw <0.1	
		1	0.75		298	seed <0.1 straw <0.1	
		2	0.75		67	seed <0.02 straw 0.55	
		2	0.75		70	seed <0.02 straw <0.05	
		3	0.75		72	seed 0.04 straw 0.1	
1981		2	0.5 + 0.75	0.083 + 0.13	73	seed <0.01	122
		1	0.75	0.13	73	seed 0.025	
		1	0.75	0.13	80	seed ≤0.02	
		1	0.75	0.13	66	seed ≤0.02	
		1	0.75	0.13	69	seed ≤0.02	
1981	Rovral Flo	1	0.75	0.13	73	seed 0.025	122
	SC	1	0.75	0.13	80	seed ≤0.02	

Country, Year	Application				PHI, days	Residue, mg/kg	Ref.
	Form	No	kg ai/ha	kg ai/hl			
		1	0.75	0.13	92	seed \leq 0.02 straw 0.2	
		1	0.75	0.13	66	seed \leq 0.02 straw 0.2	
Germany	SC 255 g/l	1	0.75	0.13	48	pods <0.05	7
1983					55	pods <0.05	
					62	pods <0.05	
					82	seed <0.01	
	SC 255 g/l	1	0.75	0.19	78	seed <0.01	
	SC 255 g/l	1	0.75	0.13	56	pods <0.05	
					74	seed <0.01	
	SC 255 g/l	1	0.75	0.19	75	seed <0.01	
Netherlands 1983	Rovral Flo SC	1	0.5	0.17	17	seed 0.27	8
UK 1981	Rovral Flo	1	0.5		39	seed 0.43	162
	SC	1	0.5		41	seed <0.1, 0.13	
		1	0.5		40	seed 0.19, 0.24	
		2	0.5		20	seed 0.20, 0.41	
1982		2	0.5		22	seed 0.12, 0.16	174
		2	0.5		22	seed 0.12, 0.14	
USA 1989	Rovral 2F SC	2	1.1		57	seed <0.05, 0.32	54
	or 4F SC	2	1.1		50	seed 0.25, 0.32	
		2	1.1		52	seed 0.13, 0.21	
		2	1.1		59	seed <0.05	
		2	1.1		56	seed <0.05, 0.25	
		2	1.1		48	seed 0.34, 0.77	

i iprodione c carbendazim

Sunflower. Table 50 shows the results of trials in France (registered) and Italy (not registered).

Table 50. Residues of iprodione in sunflower.

Country, Year	Application	PHI, days	Residue, mg/kg	Ref.
---------------	-------------	-----------	----------------	------

	Form	No	kg ai/ha	kg ai/hl		
France	Calidan	2	0.7		75	<0.04
1986		3	0.7		70	0.06
		2	0.7		85	<0.04
1990	Sumistar	2	0.4		61	Seed 0.11 Oil <0.05 Cake <0.05
1990		2	0.4		61	S 0.36 O <0.05 C <0.05
1991		2	0.4		98	S <0.05
		2	0.4		90	S <0.05
		2	0.4		114	S <0.05
		2	0.4		101	S <0.05
Italy 1986	Rovral	2	0.75		41	S 0.07
	WP 50*	1	150 g/100 kg seed		133	S <0.05

Coffee. In two trials during 1981/82 at one site in Brazil plants were treated four times with Rovral WP50 at 0.5 or 1 kg ai/ha. The seed contained 0.8 and 0.9 mg/kg iprodione after 35 days (ref. 191).

Animal transfer studies

Cows. Dairy cattle were treated once daily for 29 days with iprodione at levels corresponding to 5, 15, 50 and 200 ppm in the feed; the cows (3 per group) were slaughtered on the morning of the 29th day. Milk samples collected on treatment days 8, 17 and 28 were analyzed for iprodione and its non-hydroxylated and hydroxylated metabolites. Tissue samples (muscle, kidney, liver and fat) collected at slaughter were analyzed for iprodione and its non-hydroxylated metabolites.

The total residue of iprodione and its metabolites in the milk and tissues of the cattle increased in proportion to the level fed, reaching a plateau level by day 7. At 5 ppm there were no detectable levels of iprodione or its metabolites in the milk (<0.01 mg/kg) or tissues (<0.05 mg/kg) except a single value of 0.05 mg/kg in kidney (Craig *et al.*, 1982 (ref. 265); see Tables 51 and 52). The kidneys contained the highest residue levels. The residues in the liver, kidney and fat were roughly proportional to the feeding level; the muscle values were too low for reliable comparison.

Table 51. Total residues of iprodione and its metabolites in bovine milk.

Feeding level, Cow ppm		Total residues (mg/kg expressed as iprodione) at treatment day and time					
		8 am	8 pm	17 am	17 pm	28 am	28 pm
200	1	0.34	0.31	0.39	0.29	0.3	0.33
	2	0.31	0.38	0.29	0.32	0.25	0.26
	3	0.28	0.32	0.25	0.33	0.27	0.29
50	4	0.096	0.2	0.089	0.17	0.073	0.14
	5	0.041	0.064	0.026	0.043	0.042	0.104
	6	0.52	0.044	0.031	0.104	0.068	0.11
15	7	0.022	0.099	0.017	0.019	0.023	0.02
	8	0.018	0.018	0.016	0.025	0.016	0.019
	9	0.024	0.029	0.018	0.027	0.022	0.035
5	10	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
	11	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
	12	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01

Table 52. Total residues of iprodione and its non-hydroxylated metabolites in bovine tissues.

Feeding level, Cow ppm		Total residues (mg/kg expressed as iprodione) in			
		Muscle	Kidney	Fat	Liver

200	1	0.13, 0.06	2.2, 1.7	0.5, 0.52	1.9, 1.95
	2	0.1, <0.05	2.4, 2.1	0.44, 0.45	1.2, 1.2
	3	0.07, 0.06	2.2, 2.9	0.4, 0.46	1.2, 1.4
50	4	0.07, 0.06	0.8, 0.76	0.21, 0.18	0.66, 0.58
	5	<0.05, <0.05	0.58, 0.8	0.13, 0.13	0.42, 0.44
15	6	<0.05, <0.05	0.36, 0.45	0.09, 0.12	0.43, 0.34
	7	<0.05	0.08	<0.05	0.13
	8	<0.05	0.06	<0.05	0.08
5	9	<0.05	0.16	0.05	0.13
	10	<0.05	<0.05	<0.05	<0.05
	11	<0.05	<0.05	<0.05	<0.05
	12	<0.05	0.05	<0.05	<0.05

Hens. A study by Wargo and Guardigli (1983, ref. 266) was conducted to determine the residue levels of iprodione and its metabolites in the tissues and eggs of laying hens fed with three levels of iprodione.

Three groups of hens were dosed by capsule for 28 days at levels corresponding to 2, 20 and 100 ppm in the diet or 0.15, 1.5 and 7.5 mg/kg bw. Tissues were collected at slaughter and eggs were collected every third day and on the last three days during the treatment period and on every third day during withdrawal.

The mean residues in eggs at the plateau levels reached after day 7 of treatment were 0.1 mg/kg, 0.64 mg/kg and 1.9 mg/kg and had decreased to non-detectable levels (<0.01 mg/kg) by day 9 of withdrawal at the 2 and 20 ppm feeding levels and day 12 in the 100 ppm group. The residues in liver, muscle, kidney and fat at the 2 ppm feeding level on the 28th day of feeding were 0.53, <0.05, 0.29 and 0.15 mg/kg respectively and had decreased to non-detectable levels (<0.05 mg/kg) by day 14 of withdrawal in all three groups. At the highest dosage level of 100 ppm in the diet, the maximum residues in liver, muscle, kidney, fat and eggs were 9.13 mg/kg, 1.3 mg/kg, 6.46 mg/kg, 5.66 mg/kg and 1.9 mg/kg, after 28 days feeding.

FATE OF RESIDUES

Nomenclature of metabolites

Code	Chemical name
RP25040	3-(3,5-dichlorophenyl)hydantoin
RP30228	<i>N</i> -(3,5-dichlorophenyl)-3-isopropyl-2,4-dioximidazolidine-1-carboxamide
RP32490	3-(3,5-dichlorophenyl)-2,4-dioximidazolidine-1-carboxamide
RP32596	3,5-dichloroaniline
RP36112	<i>N</i> -(3,5-dichlorophenyl)-2,4-dioximidazolidine-1-carboxamide
RP36114	1-(3,5-dichloro-4-hydroxyphenyl)biuret
RP36115	1-(3,5-dichlorophenyl)biuret
RP36221	1-(3,5-dichlorophenyl)-5-isopropylbiuret
RP44160	3-(3,5-dichlorophenylcarbamoyl)ureidoacetic acid
RP44247	3,5-dichlorophenylurea

In animals

The metabolism of iprodione in goats, dairy cows and chickens is extensive and follows a similar qualitative pathway. Excretion into urine and faeces is extensive and rapid.

The major pathway from iprodione leads through RP30228 via hydrolysis and rearrangement to a major metabolite RP36112. Other hydrolysis reactions lead to 3,5-dichlorophenylurea (RP44247) as one of the major metabolites.

Iprodione does not accumulate in the milk or tissues of these animals.

In general, residues of iprodione and its metabolites increase linearly in the tissues and milk of dairy cattle and in the tissues and eggs of chickens and bobwhite quail fed increasing amounts of iprodione in their diets. However, withdrawal of the iprodione from the diet leads to the decrease of the residues in milk, eggs and tissues, indicating that neither iprodione nor its metabolites accumulate in dairy cattle or avian species fed the parent compound.

Dairy cows. Wilkes and Herrera (1981, ref. 267) showed that iprodione is excreted rapidly and extensively in cows. Two cows were dosed orally with [U-¹⁴C]phenyl-labelled iprodione at a daily rate of approximately 2 mg/kg bw, which corresponds to approximately 60 ppm in the daily feed. One cow received a single dose; the other received multiple doses over a period of 5 days. Excretion was monitored daily and became insignificant 7 days after the last dose, at which time the cow was slaughtered. Liver, kidney, muscle, fat, heart and blood were collected from each carcass. Urine, faeces and milk were collected throughout the treatment and withdrawal periods.

The fate of iprodione in the lactating cow was characterized by extensive urinary and faecal excretion. From 41.6 to 47% and from 27.4 to 45.9% of the total dose was found in the urine and faeces, respectively, over the entire dosing period.

A maximum of 0.3% of the dose was recovered in milk, where the total residue expressed as iprodione peaked at 0.17 mg/kg (single dose) and at 0.43 mg/kg (multiple doses). The depletion of the milk residue level during withdrawal was rapid; the residue decreased to half of the maximum within 36 hours.

Of the total dose, only an estimated 0.7% (single dose) and 0.3% (multiple doses) were retained in tissues. With the exception of the liver, none of the analyzed tissues contained more than 0.05 mg/kg total residue expressed as iprodione; 0.003 mg/kg was found in muscle. The liver contained higher residue levels (0.13 mg/kg from the single dose; 0.45 mg/kg from multiple doses).

It can be concluded that there is no significant accumulation of residues of iprodione or its metabolites in the tissues or milk. The major metabolites (> 10% of the total ¹⁴C residue) in urine and milk were RP36114 and RP32490. In faeces, iprodione and RP36114 were the predominant compounds.

A supplementary study was conducted to demonstrate the stability of residues of iprodione in milk and animal tissue samples stored at <0°C for 13 to 22 months. Iprodione and its

hydroxylated and non-hydroxylated metabolites remained stable in milk, and iprodione and its non-hydroxylated metabolites remained stable in liver, during prolonged freezer storage.

Lactating goats. In this study (Wilkes and Herrera, 1982, ref. 268; Piznik and Wargo, 1983b, ref. 259) a goat was dosed orally for five days with [U-¹⁴C]phenyl-labelled iprodione at a daily rate of approximately 2 mg/kg body weight, which corresponded to approximately 200 ppm in the daily feed. Slaughter was 4 hours after the final dose. Liver, kidney, muscle, fat and blood were collected from the carcass. Urine, faeces and milk were collected throughout the treatment period. Total ¹⁴C residue levels were determined in urine, faeces, milk and tissues. Samples of tissue and urine were extracted and the extracts analyzed by thin-layer chromatography to identify the metabolites.

Iprodione is excreted rapidly and metabolized extensively in lactating goats, with no significant accumulation of residues of iprodione or its metabolites in tissues or milk. The major compounds (>10% of the total ¹⁴C residue) found in the goat were RP36114, RP32490, RP36115, 3,5-dichloro-4-hydroxyphenylurea and iprodione itself. The metabolism is similar to that in other ruminant and avian species.

The fate of iprodione in the lactating goat was primarily characterized by extensive urinary and faecal excretion within 7 hours of dosing. The urine and faeces contained 50.7% and 8.7% of the total dose of [¹⁴C]iprodione respectively.

Only 0.3% of the total dose was recovered in the milk, where the total residue expressed as iprodione reached a maximum of 1.5 mg/kg after 4 days of dosing. Of the total dose, an estimated 4% was retained in the tissues. The liver contained the highest level of ¹⁴C (7 mg/kg as iprodione).

Laying hens. Fifteen hens were dosed orally with [¹⁴C]iprodione for 15 consecutive days at a daily rate equivalent to approximately 10 ppm in the feed which corresponds to approximately 0.7 mg/kg body weight. Five of the hens were slaughtered at each of the following intervals after the last dose: 2 hours, 3 days, and 7 days. Liver, kidney, heart, gizzard, breast muscle, thigh muscle, fat, skin, and blood were collected from each carcass. Eggs and excreta were collected throughout the treatment and withdrawal periods. Total ¹⁴C residues were determined in all samples. Samples of liver, kidney, muscle, fat, egg and excreta were extracted and the extracts analyzed by thin-layer chromatography to identify the metabolites (Wilkes *et al.*, 1982, ref. 264).

Iprodione is excreted rapidly and extensively in laying hens, with no significant build-up of residues of iprodione or its metabolites in chicken tissues or eggs. The major metabolites of iprodione in chickens were RP32490, RP36112, RP36115, RP36114, unknown Y and unknown Z.

Most of the iprodione was excreted: 85% of the administered dose was recovered from the excreta. The ¹⁴C residue levels in the eggs and tissues were very low. When a plateau was reached after 10 days of dosing, the eggs contained approximately 0.6 mg/kg expressed as iprodione. Of the tissues, the liver contained the highest ¹⁴C residue (3 mg/kg iprodione equivalents 2 hours after the last dose). The residue levels in the eggs and tissues decreased rapidly during the 7-day withdrawal period.

Figure 1: Metabolic pathways of iprodione in plants.

In plants

Several studies carried out on plants with ^{14}C -labelled iprodione showed that the chemical is not systemic by foliar application, which is the normal method of treatment, and that root uptake is a minor route of absorption (Prost, 1992b). The parent compound is generally the main component of the total residue resulting from foliar application. Metabolites (and especially RP30228) account for a significant proportion of the total residue in some plants such as lettuce and rice (straw).

Wheat and strawberries. From a study by Gouot *et al.* (1977, ref. 269), after foliar applications of [^{14}C]phenyl-labelled iprodione in a greenhouse the half-life of the parent compound ranged from 30 to 60 days. The principal compounds isolated were iprodione itself, its isomer (RP30228) and a desisopropylated metabolite (RP32490).

Peaches. Peaches were obtained from a field experiment in which [^{14}C]phenyl-labelled iprodione was sprayed at a rate equivalent to 1.1 kg ai/ha. Three applications were made: at pink bud, petal fall and pre-harvest stages (i.e. 93, 72 and 8 days before harvest, respectively).

Almost all (95%) of the total radioactive residue found in or on mature peaches was identified as unchanged iprodione. Again, the isomer of iprodione (RP30228) and the desisopropyl compound (RP32490) were found in small but detectable amounts (Gereck *et al.*, 1981, ref. 270).

Lettuce. Iprodione was applied to young lettuce under glasshouse conditions six weeks after sowing by foliar spraying, at a dose of 750 mg ai/l. Thirty-eight days after treatment, the ^{14}C compounds in the aerial portion of the plant consisted mainly of the parent compound (81% of the total radioactivity), with some RP30228 (9%) and a small amount of RP32490 (Gouot *et al.*, 1981, ref. 271).

All these metabolism studies demonstrate the same metabolic pathway.

Rice. Outdoor rice plots were treated twice with iprodione in a wettable powder formulation at a rate equivalent to 1.1 kg ai/ha, once at the "booting" stage and once at the "heading" stage (Piznik and Wargo, 1983, ref. 230). The results from this study were similar to those reported above and the residues in all the samples analyzed consisted mainly of the parent compound iprodione and its isomer RP30228. Small amounts of the desisopropyl analogue RP32490 and its isomer RP36112 were also found, as well as minor amounts of a product of further metabolism RP25040 (3-(3,5-dichlorophenyl)hydantoin).

Figure 1 above shows the metabolic pathways of iprodione in plants.

In soil

Iprodione is not persistent in the soil. It is converted rapidly into its isomer RP30228, which is the most abundant degradation compound, and several minor products including RP36221, 3,5-dichloroaniline (RP32596), bound residues, and CO_2 which was found to be the ultimate product of degradation.

3,5-Dichloroaniline has been identified in traces (3% maximum) with a very short half-life (15 days maximum), so no build-up would occur in soil.

The half-life of iprodione depends on several factors including the temperature and pH of the soil. Values at 25°C under aerobic or anaerobic conditions range from 20 to 80 days in most cases (20 to 60% of the applied compound is bound to the soil within 3 to 4 months), and 90% degradation times are in the range 100 to 320 days under the same conditions. In some cases light has been shown to accelerate the degradation process.

Accumulation does not occur after intensive use of iprodione, as demonstrated by many field trials: they showed that there was a progressive increase in the rate of degradation of iprodione with successive treatments (Prost, 1992b).

In storage and processing

Blackcurrant juice. In the UK blackcurrant bushes were treated five times with Rovral WP50 at four sites during 1981 at a rate of 0.75 kg ai/ha. Blackcurrants harvested between 6 and 9 days after the last application contained between 2.6 and 5.4 mg iprodione/kg. Juice obtained from the treated fruit contained between 0.25 and 0.37 mg iprodione/l.

At one site in 1979, blackcurrants treated five times with Rovral Flo at a rate of 1.5 kg ai/ha and harvested 7 days after the last application contained 2 mg/kg iprodione, whilst the associated juice contained 0.74 mg/l.

In 1981, blackcurrants treated 4 times at a rate of 0.75 kg ai/ha at two sites and harvested 7 days after the final application contained between 8.7 and 16 mg/kg iprodione. Juice prepared from the treated fruit contained between 1.1 and 1.4 mg/l (see Table 53).

Table 53. Residues of iprodione in black currants and their juice in the UK.

Year	Form	Application			PHI, days	Residue mg/kg:		Ref.
		No	kg ai/ha	kg ai/hl		Fruit	Juice	
1974	Rovral	4	1.1	0.05	62	3.9		125
	WP 50*	4	1.1	0.05	62	4.6		
1979	Rovral	5	0.75		7	6.1		30
	WP 50*	5	0.75		7	5.9		
		5	0.75		7	1.9		
		5	0.75		7	3.7		
		5	1.5		7	1.5	0.61	
1981	Rovral	5	0.75		6	2.9	0.37	165
	WP 50*	5	0.75		7	4.8	0.27	
		5	0.75		7	2.6	0.25	
		5	0.75		9	3.2	0.36	
1979	Rovral Flo*	5	1.5		7	2.0	0.74	155
1981	Rovral	4	0.75		7	14	1.4	164
	Flo*	4	0.75		7	8.7	1.1	

* Not registered for this use in the UK.

Hops and beer. In Germany, hops were treated with Rovral WP50 at the rate of 0.75 to 1.3 kg ai/ha and residues of iprodione were determined in the hops and beer brewed with them. Data summarized in Table 54 show that the highest residue in beer was 0.05 mg/kg (Laurent and Chabassol, 1980, 1981, refs. 260, 261).

Table 54. Residues of iprodione in hops and beer in Germany.

Year	Form	Application			PHI, days	Residue, mg/kg	Ref.
		No	kg ai/ha	kg ai/hl			
1979	Rovral WP 50**	2	0.75 + 1.2	0.037 0.037	7	hops 41	248
					14	23	
					21	26	
					28	25	
					28	beer 0.05	
1980		2	1.3	0.037	21	hops* 11	248
						beer 0.025	
		2	1.1	0.037	22	hops* 12	
						beer 0.025	

* Dried cones

** Not registered for this use in Germany.

Grape fractions: juice, pomace and raisins. (Residues in wine and spirits are discussed in the 1977 monograph). In the USA (Chow *et al.*, 1982, ref. 262), Rovral WP50 was applied to grape vines at eleven test sites located in California (6), New York (3), Ohio (1) and Pennsylvania (1) at a rate of 1.1 kg ai/ha. Generally, each site received 4 treatments: two before bunch closing and two more before harvest. All tests were small-scale except two in California where commercial ground equipment was used. Grapes, pomace, juice and raisins were analysed for iprodione and its metabolites RP 30228 and RP 32490. Residues in juice were generally lower than in grapes. Results are summarized in Table 55.

Grapes. There is a striking difference in the residue picture between the large- and small-plot trials: about one week after the last application, the iprodione residues ranged from 16 to 49 mg/kg (mean 32 mg/kg) in the small plots but the corresponding residues from the two large plots were 1.02 and 1.7 mg/kg.

Iprodione decline data from the three New York trials show only a slight decrease in iprodione during the first week after the last application, followed by a more pronounced decline towards the end of the test period.

Pomace. As with grapes, iprodione constituted the greatest proportion of the residue. In the small-scale trials, the iprodione content in or on pomace ranged from 6.8 to 39 mg/kg in those samples taken 7 or 8 days after the last application. The iprodione residues from the large plots were 0.74 and 1.5 mg/kg. Generally, the iprodione residues in pomace were higher than in the corresponding fresh grape samples.

The detection of iprodione residues in the untreated control samples indicates that there was a contamination problem, possibly occurring during processing.

Juice. The iprodione residues in juice made from the grapes taken one week after the last application ranged from 3.9 to 19 mg/kg in the small-scale trials but were lower in juice from the two large-scale sites (*c.* 2 mg/kg). As with the pomace, contamination of the untreated samples was noted.

Raisins. In raisins prepared from the fresh grapes, excluding the large trial plots, iprodione residues ranged from 43 to 130 mg/kg. The residues in or on samples from the large plots were 5.3

and 9.6 mg/kg. The iprodione content in raisins was higher than in the corresponding grapes owing to the drying of the grapes.

Table 55. Residues of iprodione, RP 32490 and RP 30228 in grapes, pomace, juice and raisins (USA, 1981). Rovral WP 50, 4 applications of 1.1 kg ai/ha, 7/8 day PHI (reference 248).

Commodity	Residue, mg/kg		
	iprodione	RP 32490	RP 30228
grapes	16-49 (mean 32)* 1.02, 1.7**	<0.05-1.9	0.06-0.47
pomace	6.8-39* 0.74, 1.48**	2.9 (maximum)	0.51 (maximum)
juice	3.9-19* 2**	0.58 (maximum)	0.07 (maximum)
raisins	43-130* 5.3, 3.7**	1.8 (maximum)	3.7 (maximum)

* small plots ** large plots

Tomato fractions. Iprodione residue trials on tomatoes were conducted in the USA on small plots sprayed with Rovral WP50. The rates were 1.1 kg ai/ha, the maximum use rate, and 2.2 kg ai/ha. Five applications were made at weekly or bi-weekly intervals, the last being on the day of harvest. Random representative fruits were picked from each treatment of each plot for residue analysis, and a processing study was conducted to obtain samples of wet pomace, dry pomace, juice and ketchup. Table 56 shows the results.

Residues in the various fractions were in the order dry pomace > wet pomace > ketchup > puree = juice. Residue levels in wet pomace and dry pomace were respectively about 5 times and 21 times those in the whole tomatoes. In the other fractions (juice, puree and ketchup) residue levels were lower than in the whole fruit.

Table 56. Residues of iprodione in tomato processed fractions (USA, 1986).

Form	Application		Residue, PHI, days	mg/kg	Refer- ence
	No	kg ai/hl			
Rovral WP 50	5	1.1	0	WP 1.5 DP 5.1 J 0.12 P 0.08 K 0.16 F 0.22,1.6	68
Rovral WP 50	5	2.2	0	WP 1.4 DP 8.7 J 0.17 P 0.29 K 0.59 F 0.46,2.8	

WP Wet pomace J Juice F Fruit
DP Dry pomace P Puree K Ketchup

Processed potatoes: chips, flakes, granules. In the USA, plants were sprayed initially with Rovral WP50 at approximately row closing at the rates of 1.1 and 2.2 kg ai/ha. Three additional sprays were made at approximately two-week intervals with the last spray 14 days before harvest. In some cases the pre-harvest interval was shorter or longer than the targeted 14 days. The trials were on autumn, spring and summer crops. At harvest representative, random samples of commercial tubers

and culls were collected from each treatment for residue analysis. In addition to the tubers and culls, potatoes from two trials were used to prepare potato flakes and chips (residues were <0.05 mg/kg).

No finite residues were found in potato chips or granules (<0.05 mg/kg; USA, 1986, reference 64) after treating potatoes with 4 x 11.0 kg ai/ha. In potato flakes 0.16 mg/kg was found in the sample treated at the high rate while no residues were detected in the low-rate sample. No residues (<0.05 mg/kg) were found in the stock feed (potato peels) from the granule processing after treatment at either rate. The stock feed from the flake and the chip processing contained respectively 0.32 and 0.27 mg/kg iprodione after treatments at 5.5 kg ai/ha and 0.1 and 0.69 mg/kg from those at 11 kg ai/ha. The concentration factor for the residues from the whole potato to the peel varied from about 0.3 to 2.

Analyses have also been carried out in the UK to show the effects of cooking on residues of iprodione in potatoes treated post-harvest with an aqueous flowable formulation containing 500 g/l iprodione, at a rate of 100 g ai/tonne. Samples were removed from the farm store approximately one month after treatment.

Raw, washed potatoes contained 13 mg/kg iprodione. It was found that in boiled and chipped potatoes iprodione levels were reduced to 0.57 and 1.7 mg/kg respectively, while in oven-baked and microwaved potatoes the residues were 12 and 15 mg/kg respectively. Results are summarized in Table 57.

Table 57. Residues of iprodione in processed potatoes.

Country year	Form	Application No kg ai/ha		PHI, days	Residue mg/kg	Ref.	
USA 1985	Rovral WP 50	4	1.1	6	T <0.05	76	
				12	<0.05		
				14	<0.05 (11)		
				14	<0.05 (2)		
				30	<0.05 (2)		
				6	C <0.05		
				12	0.08		
		14	<0.05 (11)				
		14	0.09, 0.11				
		30	<0.05 (2)				
		1985	4	2.2	6	T <0.05	76
					12	<0.05	
					14	<0.05 (9)	
					14	<0.05 (4)	
30	<0.05 (2)						
6	C <0.05						
12	<0.05						
14	<0.05 (9)						
14	0.05-0.1 (4)						
30	<0.05 (2)						
1985	4	1.1	14	T	<0.05	76	
				FL	<0.05		
		2.2	15	T	<0.05		
				FL	<0.05		
				CH	<0.05		
				CH	<0.05		
1986	4	5.5	14	T	0.17	64	
				FL	<0.05		
				CH	<0.05		
				SC	0.27		
				SF	0.32		
		11.0	14	SG	<0.05		
				GR	<0.05		
				T	0.32		
				FL	0.16		
				CH	<0.05		

					SC	0.69
					SF	0.10
					SG	<0.05
					GR	<0.05
UK	Rovral	1	0.1	30	tuber,raw	257
1990/91	Aqua		kgai/t		13.0	
	Flo				tuber, boiled	
					0.5	
					tuber, baked	
					12.0	
					tuber, fried	
					1.7	
					tuber,microwaved	
					15.0	
T	Tubers	SC	Stock feed from chip processing			
C	Culls	SF	Stock feed from flake processing			
FL	Flakes	SG	Stock feed from granule processing			
CH	Chips	GR	Granules			

Cereal fractions, flour and bread. In the USA, maize grain was treated with Rovral WP50 at rates of 2 and 4 kg ai/t before being placed in a storage silo.

Table 58 shows the residues in the grain after storage up to 84 days and in the grits, meal, flour, crude oil and refined oil from dry and wet milling operations. The residues in grits, flour, meal and oil (crude and refined) were lower than in the grain, demonstrating that the residues were not concentrated by processing. A reduction in the residue of over 100-fold was observed from crude to refined oil.

Table 58. Residues of iprodione in processed fractions of maize (USA, 1986).

Form	Application		Residue, mg/kg	Ref.
	No	g ai/ 100 kg		
Rovral WP 50	1	2	grain 18	66
			grits 1.9	
			flour 14	
			meal 9.9	
			crude oil 10.4	
			refined oil 0.06	
			starch 10.4	
	1	4	grain 44	66
grits 4.0				
flour 35				
meal 24				
crude oil 11				
refined oil 0.06				
starch 24				

In the UK, flour from wheat treated three times with iprodione at a rate of 0.5 kg ai/ha and harvested between 5 and 47 days after the last application contained 0.14-0.73 mg/kg iprodione, whilst the corresponding bread samples contained <0.1-0.45 mg/kg. Flour from wheat treated four times with iprodione at a rate of 0.5 kg ai/ha and harvested on the day of the last application contained 0.63-1.1 mg/kg iprodione and the corresponding bread 0.30-0.48 mg/kg (Brockelsby *et al.*, 1987, ref. 263; see Table 59).

Table 59. Residues of iprodione in wheat flour and bread (UK, 1986).

Form	Application No kg ai/ha	PHI, days	Residue, mg/kg	Ref.
Sirocco	3	0.5	47 flour 0.21,0.22 bread 0.13,0.14	248
	3	0.5	9 flour 0.14,0.20 bread <0.1,0.12	
	3	0.5	5 flour 0.39,0.73 bread 0.29,0.45	
Sirocco Rovral Flo	4	0.5	0 flour 0.66,0.85 bread 0.33,0.37	248
	4	0.5	0 flour 0.63,1 bread 0.30,0.38	
	4	0.5	0 flour 0.88,1.1 bread 0.47,0.48	

Peanut processed fractions. Rovral WP50 was applied to peanut trial plots in the major peanut-producing areas of the USA in 1983 and 1984. Three foliar applications were made to the plants at 1.1 or 2.2 kg ai/ha per application. Residues in hay, hulls, nut-meat and processed fractions are shown in Table 60.

Hay. Iprodione accounted for the greatest part of the residues. The highest residue of iprodione after 11 days was 65 mg/kg.

Hulls contained residues up to 5.2 mg/kg of iprodione.

Nut-meat. No residues (<0.05 mg/kg) were present in any of the 1983 samples, but residues of 0.21 and 0.27 mg/kg iprodione from the low and high application rates respectively were found in the 1984 samples owing to contamination from the hulls during commercial shelling of the peanuts.

Processed fractions. In the 1984 trials 0.54 and 0.67 mg/kg of iprodione were found in the screw-press crude oil but none in the solvent-extracted crude oil, refined oil or peanut meal. The residues in the screw-press oil were caused by surface contamination from the hulls during their commercial separation from the nut-meat. Soapstock contained residues of 1.3 and 2.6 mg/kg iprodione.

Table 60. Residues of iprodione in processed fractions of peanuts (USA, ref. 248).

Year	Form	Application		PHI, days	Residue, mg/kg (No. of samples)
		No	kg ai/ha		
1983	Rovral WP 50	3	1.1		hay
				0	28-147 (3)
				3	57
				4	16- 77 (3)
				5	56
				9	90
				11	25 -65 (3)
					hulls
				0	0.57-5.2 (3)
				3	0.39
				4	1.2-5.2 (3)
				5	0.67
				9	0.61
				11	0.26-1.1 (3)
	nut-meat				
0-11	<0.05 (12)				
1984		3	1.1	4	hulls 2.3
				4	nut-meat 0.21
				4	SPCO 0.54
				4	SECE <0.05
				4	refined oil<0.05
				4	soapstock 1.3
		3	2.2	4	hulls 4.3
				4	nut-meat 0.27
				4	SPCO 0.67
				4	SECE <0.05
				4	R. oil <0.05
				4	soapstock 2.6
				4	peanut meal<0.05
SPCO screw-press crude oil		SECE solvent-extracted crude oil			
R.oil refined oil					

RESIDUES IN FOOD IN COMMERCE OR AT CONSUMPTION

The results of enforcement analyses by the Governmental Food Inspection Services in The Netherlands in 1987-1991 were brought to the attention of the Meeting.

In 1991 65 commodities were analysed (1058 samples). Residues above the MRLs were found in only a few samples in eight commodities (see Table 61).

Table 61. Enforcement analyses for iprodione of the Governmental Food Inspection Services in The Netherlands (1991).

Commodity	No. of samples		
	Analysed	<MRL	>MRL
scarole	144	142	2
kiwifruit	29	28	1
lettuce, head			
incl. cos lettuce	246	244	2
nectarine	8	6	2
sugar pea			
(young pods)	2	1	1
leek	8	7	1
turnip tops,			
incl. rucola	19	15	4
corn salad	27	26	1

The results of enforcement analyses in 1987-1990 are summarized in Table 62.

Table 62. Results of enforcement analyses for iprodione in The Netherlands, 1987-1990.

Commodity	No.*	Mean	Std. dev.	Median	Min.	Max.	95% max.
Currants, Black, Red, White	32	1.171	1.793	0.5	0.05	8.1	5.6
Strawberry	398	1.163	1.984	0.5	0.01	22	4
Scarole	493	0.841	2.875	0.22	0.01	41.5	3
Apple	33	0.602	0.671	0.5	0.02	3.1	1.6
Egg plant	35	0.419	0.653	0.07	0.02	2.95	2.2
Celery, blanched	57	0.368	0.772	0.1	0.01	3.54	2.56
Kale	5	0.338	0.472	0.14	0.08	1.18	1.18
Beans (runner/snap/slicing)	79	0.159	0.422	0.07	0.01	3.7	0.56
Blackberries	22	1.073	0.982	0.905	0.05	4.4	2.5
Broccoli	23	0.24	0.264	0.1	0.03	1	0.75
Chinese cabbage, amsoi, choisum and paksoi	33	0.328	0.424	0.14	0.02	1.7	1.32
Grapes	41	0.463	0.564	0.28	0.02	2.5	1.7
Raspberries	12	1.086	1.315	0.55	0.05	4.6	4.6
Crisphead lettuce	48	0.185	0.235	0.1	0.02	1.3	0.5
Cherries	7	0.911	0.928	0.683	0.11	2.72	2.72
Kiwifruit	27	0.805	1.056	0.43	0.01	5	2.4
Celeriac	5	0.614	0.512	0.62	0.1	1.35	1.35
Fennel, bulb	27	1.979	3.592	0.5	0.03	15.5	11.8
Cucumber	23	0.095	0.201	0.04	0.01	1	0.14
Lettuce, Head incl. cos lettuce	1351	0.72	1.982	0.27	0.01	42	2.9
Lettuce, Leaf	43	0.663	1.363	0.21	0.03	8.1	2.1
Melons, except watermelon	10	0.063	0.076	0.04	0.003	0.27	0.27
Other agricultural crops	6	0.772	1.258	0.21	0.09	3.3	3.3
Other vegetables	40	1.179	1.887	0.33	0.02	9	4.5
Other agricultural and horticultural crops	33	0.241	0.516	0.07	0.01	2.6	1.54
Sweet and chilli peppers	148	0.346	0.762	0.08	0.01	4.99	2.1
Peach	14	0.241	0.222	0.22	0.01	0.64	0.64
Parsley, root parsley	23	0.832	1.155	0.3	0.1	4.23	3.35
Winter squash, courgette and patisson	9	0.201	0.113	0.23	0.04	0.4	0.4
Purslane	6	1.303	1.512	0.785	0.21	4.18	4.18
Leek	24	0.521	0.496	0.38	0.02	1.8	1.8
Plums, inc. prunes	10	0.197	0.153	0.195	0.03	0.57	0.57
Turnip, tops inc. rucola	60	5.308	7.594	2.24	0.05	30.6	29
Rhubarb	8	0.34	0.28	0.33	0.04	0.75	0.75
Radish	226	0.289	0.629	0.1	0.01	5	1.3
Radish, Black	10	0.274	0.608	0.075	0.04	2	2
Celery leaves	59	1.544	4.613	0.2	0.03	27.8	7
Leaf lettuce	17	0.355	0.652	0.11	0.05	2.6	2.6
Spinach	65	0.74	1.433	0.2	0.02	9.5	2.6
Cabbage, pointed	5	0.258	0.183	0.19	0.1	0.5	0.5
Brussels sprouts	49	0.788	1.467	0.1	0.02	6	4
Tomato	21	0.172	0.16	0.14	0.01	0.55	0.52
Corn salad	208	2.77	3.916	1.425	0.02	28	8.54
Fennel	7	0.624	1.144	0.16	0.1	3.2	3.2

Commodity	No.*	Mean	Std. dev.	Median	Min.	Max.	95% max.
Winter carrot	16	0.368	0.325	0.215	0.03	1.2	1.2
Witloof chicory (sprouts)	73	0.394	0.861	0.09	0.02	4.8	2.6
Carrot	65	0.915	1.34	0.3	0.01	5.28	3.63

* Only the numbers of samples with residues were reported

METHODS OF RESIDUE ANALYSIS

General method for plant substrates. Analytical methods for determining residues in plant substrates are specific for the parent compound iprodione and when needed (as in the USA) for its metabolites RP30288 (isomer of iprodione) and RP32490 (desisopropylated metabolite). Determination is by GLC with an EC detector after solvent extraction of the substrate and clean-up by Florisil column chromatography. The limit of determination in most crop and animal samples is between 0.01 and 0.1 mg/kg (Prost, 1992d).

NATIONAL MAXIMUM RESIDUE LIMITS

The following national MRLs were reported to the Meeting.

Country	Commodity	MRL, mg/kg
Argentina	Grapes	1
	Lettuce	0.5
	Strawberries	0.5
Australia	Beans	0.2
	Berry fruit	12
	Celery	2
	Grapes	20
	Kiwifruit	10
	Lettuce	5
	Lupin seed	0.1
	Meat	0.1
	Milk and milk products	0.1
	Passion fruit	10
	Peanut	0.05
	Peanut forage (green)	20
	Pome fruits	3
	Potato	0.05
	Soya bean (dry)	0.05
	Soya bean forage (green)	5
Belgium	Stone fruits	10
	Tomato	2
	Banana	5
	Carrot	2
	Citrus	5
	Corn salad	10
	Grapes	10
	Oilseed rape	0.2
	Other commodities of plant origin	0.05
	Potatoes	0.5
Brazil	Shallot	5
	Stone fruit	5
	Strawberries	10
	Vegetables	5
	Witloof (leaves)	1
	Carrots	0.02
	Coffee	0.8
	Cotton	0.1
	Garlic	0.1
	Grapes	1
	Lettuce	1
	Onion	0.01
	Peaches	5
	Potatoes	0.02
	Strawberries	1
	Sugar cane	2.5
	Sweet pepper	4

iprodione

779

Country	Commodity	MRL, mg/kg
	Tomatoes	4
	Wheat	0.3
Canada	Beans	0.3
	Cherries	5
	Cucumber	0.5
	Ginseng	0.1*
	Grapes	10
	Kiwifruit	0.5
	Onion	0.1*
	Peach	5
	Strawberry	5
	Tomato	0.5
Denmark	Berries and small fruits	10
	Cabbage	10
	Leafy vegetables	10
	Other vegetables	5
	Pome fruits	10
	Potato	5
	Stone fruits	10
Finland	Boysenberry	10
	Kiwifruit	10
	Other small fruits and berries	5
	Vegetables	5
France	Beans, French	5
	Berries, other	7
	Carrots	2
	Cereals, grain	0.05
	Cucumbers	5
	Grapes	10
	Kiwifruit	7
	Lettuce	10
	Onions	2
	Pome fruit	10
	Potato	0.02
	Stone fruit	10
	Strawberry	10
	Tomato	5
	Rape seed	0.5
	Witloof	1
Germany	Cereal grains	0.5
	Cucumber	5
	Grapes	5
	Kiwifruit	5
	Kiwifruit, peeled	0.5
	Lettuce	10
	Oilseed	0.2
	Other commodities of plant origin	0.02
	Peas, field	0.1
	Peppers, Sweet	5
	Plums	3
	Strawberry	10
	Tomato	5

iprodione

Country	Commodity	MRL,mg/kg
	Witloof chicory (sprouts)	1
Hungary	Apples	3
	Cabbage	0.2
	Grapes	5
	Nuts	0.2
	Pears	3
	Soft fruit	3
	Stone fruit	5
	Strawberries	3
	Sugar beet	0.2
	Sunflower	3
Italy	Apples	3
	Cabbage	0.5
	Egg plant	0.5
	Endive	0.5
	Grapes	5
	Kiwifruit	3
	Lemon	1
	Lettuce	0.5
	Oilseed rape	0.05
	Pears	5
	Pineapple (whole fruit)	5
	Rice	0.5
	Stone fruits	5
	Strawberry	0.5
	Tomato	0.5
Japan	Banana	10
	Beans	1
	Beans (Soya, Broad)	0.2
	Cereals (wheat and barley)	10
	Cherries	10
	Chicory	1
	Citrus	10
	Mango	10
	Melon	10
	Papaya	10
	Pineapple	10
	Potatoes	0.1
	Rice	3
	Sugar beet	1
	Vegetables (incl. Cucumber)	5
	Watermelon	10
Netherlands	Beans (fresh)	1
	Beans (dry)	0.2
	Berries, other	5
	Blackberries	5
	Caraway, seed	2
	Cherries	5
	Corn salad	10
	Carrots (post-harvest)	10
	Currant, black	5
	Currant, red	15

iprodione

781

Country	Commodity	MRL,mg/kg
	Garlic	0.1
	Grapes	10
	Kiwifruit	5
	Kohlrabi	5
	Lettuce, leaf	5
	Meat	0.05*
	Milk	0.05*
	Onions	0.1
	Other food commodities	0.05*
	Peaches	10
	Pome fruits	10
	Plums	10
	Prunes	10
	Pulses (dry)	0.2
	Raspberries	5
	Rice	3
	Squash	1
	Strawberries	10
	Vegetables, other	5
New Zealand	Berries	10
	Grapes	10
	Kiwifruit	7
	Stone fruits	10
	Tangelo	2
	Tomato	5
South Africa	Grapes (table, wine)	5
	Onions	0.5
	Peaches	5
	Peaches (canned)	0.05
	Pears	0.05
	Tomato	2
Spain	Banana	5
	Bulb vegetables	1
	Citrus fruits	5
	Grapes	10
	Kiwifruit	5
	Leafy vegetables	10
	Fruiting vegetables	5
	Pineapple	5
	Pome fruits	10
	Root and bulb vegetables	1
	Stone fruits	10
	Strawberry	10
Sweden	Strawberry	10
	Vegetables	5
Switzerland	Asparagus	0.05
	Beans	0.5
	Blackberries	2
	Carrot	2
	Chinese cabbage	1
	Grapes	7
	Kiwifruit	5

iprodione

Country	Commodity	MRL,mg/kg
	Kiwifruit (peeled)	0.5
	Lettuce	6
	Onions	0.1
	Pome fruits	0.05
	Raspberries	2
	Stone fruits	0.05
	Tomato	6
	Wine	2
United Kingdom	Apples	10
	Bulb vegetables	0.1
	Cabbage	3
	Cucumbers	5
	Common bean	3
	Currants (red, black, white)	5
	Grapes	10
	Lettuce	10
	Nectarines	10
	Onions	0.1
	Peas	3
	Peaches	10
	Pears	10
	Plums	10
	Raspberries	5
	Strawberry	10
	Tomato	5
USA	Almond hulls	2
	Almond nut-meat	0.3
	Apricots	20
	Bean hay, forage	90
	Beans (dry, succulent)	2
	Blackberries	25
	Blueberries	15
	Boysenberries	15
	Broccoli	25
	Carrots	5
	Cherries, sour	20
	Cherries, sweet (pre- and post-harvest)	20
	Currants (import MRL)	15
	Field corn grain (stored)	20
	Garlic	0.1
	Ginseng (dry)	4
	Ginseng (fresh)	2
	Grape pomace	225
	Grapes	60
	Lettuce (head)	25
	Loganberries	25
	Nectarines, peaches	20
	Onions, Bulb	0.5
	Peanut	0.5
	Peanut forage, dry and hay	150
	Peanut hulls	7
	Plums	20

Country	Commodity	MRL, mg/kg
	Potatoes	0.5
	Prunes (fresh)	20
	Raisin waste	300
	Raisins	300
	Raspberries	15
	Rice bran	30
	Rice grain	10
	Rice hulls	50
	Rice straw	20
	Strawberries	15
	Tomatoes	3
	Youngberries	25
Yugoslavia	Apples	3
	Grapes	2
	Strawberries	2

APPRAISAL

Iprodione, used as a fungicide for a variety of crops, was reviewed in the CCPR periodic review programme. It is applied by pre- or post-harvest foliar spray, dipping of plants or roots, and as a seed treatment. It is formulated as WP or SC, and for foliar spraying it is applied between one and five times at a rate of 0.25-1.5 kg ai/ha. In the post-harvest treatment of fruits and vegetables it is used at spray concentrations of 0.075-0.5 kg ai/hl and for seed treatment at rates up to 2.5 g/kg seed. Residue data have been received from supervised trials on citrus, pome and stone fruits, berries, grapes, bananas, kiwifruit, vegetables (brassica, bulb, leafy, legume, root and tuber, stalk and stem vegetables, cucumbers, tomatoes and peppers), cereals, nuts and seeds.

Metabolism studies in plants have shown that iprodione is degraded to *N*-(3,5-dichlorophenyl)-3-isopropyl-2,4-dioximidazolidine-1-carboxamide (RP30228), which is the only significant metabolite in certain crops, and also to the desisopropyl derivative (RP32490), a relatively minor metabolite.

The parent compound is generally the main component of the total residue resulting from foliar application. Except in potatoes, certain rice fractions (especially straw) and peanut hay, residues of metabolites are undetectable.

Iprodione is metabolized extensively in goats, chickens and dairy cows. Excretion in urine and faeces is extensive and rapid. Iprodione does not accumulate in the milk, eggs or tissues. Residues of iprodione and its metabolites decrease rapidly when the dose is withdrawn. Metabolism is similar in goats, chickens and dairy cattle.

Iprodione has low water solubility, is not volatile and is expected to dissipate rapidly in water under natural field conditions. Under laboratory conditions it shows no significant degradation at pH 3, a half-life of 20 days at pH 6 and complete degradation in less than 24 hours at pH 9.

Iprodione has low mobility in various soil types and does not accumulate in soil: its half-life generally varies, in the laboratory, between 20 and 80 days. Light has been shown to accelerate degradation under aerobic conditions. Field trials showed that there was a progressive increase in the

rate of degradation with successive treatments.

Processing studies on black currants, grapes, tomatoes, potatoes, maize, wheat, peanuts and hops showed a reduction of the residues in juice, beer, flour and potato flakes and chips, but a concentration in dry pomace and raisins.

Methods of analysis are based on GLC with an EC detector after solvent extraction of the substrate and clean-up by Florisil column chromatography. The limit of determination in most crop and animal samples is between 0.01 and 0.1 mg/kg. The methods determine the parent compound, and the two metabolites if required.

The residue data from supervised trials are on the parent compound only and were evaluated as follows.

Citrus fruits. Data were available from New Zealand, Japan and Israel, but there was no information on GAP. As there were only two trials with dip treatments (0.05-0.075 kg ai/hl, in Italy), the data were also rather limited. No MRL could be recommended.

Pome fruits. Trials on apples with spray treatments in Japan were not in line with GAP. Only three results could be evaluated for pears from spray treatments (Italy, 4-7 treatments, application rate 1.35 kg ai/ha, PHI 20 to 21 days). The residues were 0.51-3.6 mg/kg. These results are not sufficient to recommend an MRL, particularly as according to GAP there should be only 2-3 treatments.

Post-harvest dipping trials (one treatment, 0.05-0.075 kg ai/hl) carried out in France and Italy on apples could be used to estimate a maximum residue level. As there was no decrease in residues during storage, ten results from storage periods between 0 and 222 days could be evaluated (minimum residue 0.61 mg/kg, maximum 2.4 mg/kg). The Meeting agreed to replace the recommendation for apples and pears (10 mg/kg Po), by a recommendation for pome fruits of 5 mg/kg Po.

Apricots. There were only two trials from France, and samples were not taken within the recommended PHI. An MRL could not be recommended.

Cherries. There were five values from Canadian spray trials. There were 6-8 treatments at rates of 0.7-0.875 kg ai/ha with a one-day PHI (residues 1.1-6.5 mg/kg). The Meeting considered the data from cherries and from other stone fruits to be mutually supportive and estimated a maximum residue level of 10 mg/kg for cherries.

Peaches. There were nine values from Canadian trials, based on 2-5 pre-harvest spray treatments at rates of 0.75-1.0 kg ai/ha with a PHI of 0-1 day, similar to Canadian GAP. Residues were 2.2-7.6 mg/kg. In one trial the residues on days 4, 7 and 14 after treatment (9, 10 and 8.5 mg/kg respectively) were higher than on day 0 (7.6 mg/kg). There were no new data on post-harvest treatments. The Meeting estimated a maximum residue level of 10 mg/kg for peaches to replace the previous recommendation (10 mg/kg Po).

Data from France and Australia on nectarines were made available to the Meeting, but there is no GAP in France. A single value from Australia was not sufficient to recommend an MRL.

Plums. Iprodione is registered in France, New Zealand and the USA. For pre-harvest spraying there are 2-3 treatments, a rate of 0.037-0.075 kg ai/hl and a PHI of 0-3 days. There were no residue data from these countries that could be evaluated. Data from South Africa were made available, but there was no GAP. The Meeting agreed to withdraw the recommendation for an MRL for plums (including prunes) of 10 mg/kg.

Gooseberry; Currants, Black, Red, White. The trials on currants and gooseberries carried out in the UK and The Netherlands were not in conformity with GAP, with one exception. The residues considerably exceeded the MRL recommended in 1977, the maximum being 20 mg/kg. The Meeting agreed to recommend withdrawal of the MRL for currants (5 mg/kg).

Blackberries and raspberries. Residue data from four trials on blackberries in the USA in 1986 were provided. The US GAP is included in the GAP for caneberries, which specifies a maximum of 4 applications. Five applications were used in the trials, but this was not regarded as significantly different from 4 in its effect on residues. The residues covered a wide range, from 5.8 to 22 mg/kg, reflecting the typical variation of residues in berry fruits. The Meeting noted that trials in Canada on raspberries with a similar use pattern gave similar residues. The Meeting estimated a maximum residue level of 30 mg/kg for iprodione in blackberries.

In raspberry trials there were four results from the UK which corresponded to GAP: 4-5 treatments, 0.75 kg ai/ha and a PHI of 7 to 8 days. The residues were 0.92-5.4 mg/kg. The Meeting also evaluated Canadian raspberry data (residues 4.6 to 31 mg/kg) according to US GAP and used US blackberry data in support. The Meeting estimated a maximum residue level of 30 mg/kg for raspberries to replace the previous recommendation (5 mg/kg).

Strawberries. There were many results from world-wide strawberry trials. The five US trials with 22 results (minimum residue 0.5 mg/kg, maximum 9.1 mg/kg) were based on 1-5 treatments at 1.1 kg ai/ha with a 0-day PHI. From Belgium there were 4 residues from trials according to GAP (1.9-6.0 mg/kg). In Germany (6 results), residues 10-12 days after the last of 3 treatments at 0.94-1.25 kg ai/ha were lower at 0.13-4.5 mg/kg. The UK provided 7 results from trials according to GAP (0.75 kg ai/ha, 1-day PHI) with residues of 1.9-4.8 mg/kg. There were 6 results from the similar Canadian GAP (1.1 kg ai/ha, 1-day PHI) (minimum residue 0.97 mg/kg, maximum 6 mg/kg) and two results each from New Zealand (3 treatments, 0.7 kg ai/ha, 1-day PHI) and Spain (2-3 treatments, 0.75 kg ai/ha, 3-day PHI) with values from 1.6 to 3.9 mg/kg. The Meeting agreed to maintain the current recommendation of 10 mg/kg for strawberries.

Grapes. Extensive data on grapes were available from France, Chile, Portugal, Spain, Italy, Germany, Morocco, Canada and the USA. The official application rates (mostly 0.75-1.0 kg ai/ha) and number of applications (mostly 3 or 4) were similar from country to country, but the official PHI varied from 0 days (USA) to 28 days (Germany and Italy). Iprodione residues were quite persistent, and sometimes higher at longer intervals than at the official PHI. Numerous residues were in the 1-5 mg/kg range, with three in the 6-10 mg/kg range and three at 10-11 mg/kg. The Meeting agreed to maintain the current recommendation of 10 mg/kg for grapes.

Bananas. In Spain, iprodione is registered as a post-harvest treatment (1 application, 0.03 kg ai/hl, no PHI). Three residue values from trials according to GAP were available (pulp <0.1 mg/kg, peel 4-7 mg/kg, whole fruit 1.7-3.4 mg/kg). Only three values from one country were not sufficient to

recommend an MRL.

Kiwifruit. The suggested MRL is based on 13 values from New Zealand 3-5 applications at 0.75 kg ai/ha and a 1-day PHI (minimum residue 0.28 mg/kg, maximum 4.5 mg/kg). The residues determined in six samples of peeled fruit were between 0.14 and 0.91 mg/kg. The Meeting agreed to maintain the current recommendation of 5 mg/kg for kiwifruit. No GAP was available for post-harvest treatment.

Fennel. The data from one supervised trial were insufficient to recommend an MRL.

Garlic. Data from two trials were available, but their validity could not be confirmed. The Meeting agreed to withdraw the recommendation for garlic (0.1 mg/kg).

Bulb onions. There were six results from Canada (2-5 spray treatments, 0.75-1 kg ai/ha, a 13 to 19-day PHI) which approximated GAP, giving residues of 0.05-0.18 mg/kg. The 5 values from two US trials according to GAP ranged from <0.05 mg/kg to 0.11 mg/kg. The Meeting estimated a maximum residue level of 0.2 mg/kg for onions to replace the previous recommendation (0.1 mg/kg).

Broccoli. Eight values from 4 trials in the USA (1.1 kg ai/ha, 2 treatments) showed iprodione residues between 4.1 and 22 mg/kg. The Meeting estimated a maximum residue level of 25 mg/kg for iprodione in broccoli.

Cauliflower. Trials from Canada could not be evaluated because no Canadian GAP was available. Some French trials were within the application conditions of GAP, but the intervals after treatment were much longer than the official PHI of 15 days. The data could not be used to estimate a maximum residue level.

Head cabbages. Trials from Canada, Germany and the UK were not in conformity with GAP. An MRL could not be recommended.

Chinese cabbage. Trials from Canada, Germany, Denmark and the USA were generally not in accord with GAP. An MRL could not be recommended.

Kohlrabi. There was only one residue value from a treatment according to GAP, from The Netherlands. An MRL could not be recommended.

Cantaloupe. Although there were two trials from France, there was no information on French GAP. An MRL could not be recommended.

Cucumbers. Residue data were available from 7 trials which did not closely reflect GAP. An evaluation of a total of 10 results from France (4 treatments, 0.075 kg ai/hl, a 5-day PHI), the UK (4 treatments, 1.1 kg ai/hl, a 2-day PHI), Denmark (2 to 4 treatments, 0.05 to 0.075 kg ai/hl, PHI 2 to 4 days) and Canada (3 to 9 treatments, 0.05 kg ai/hl, 1-day PHI) showed residues of <0.1-1.8 mg/kg). The Meeting therefore estimated a maximum residue level of 2 mg/kg to replace the previous recommendation (5 mg/kg).

Sweet peppers. Seven US trials based on 8 applications and a 0-day PHI were not in line with GAP (up to 4 treatments, PHI of 3-14 days). The Meeting agreed to withdraw the recommendation for

sweet peppers (5 mg/kg).

Tomatoes. Outdoor trials from the USA could not be evaluated because no US GAP was available. The conditions of application in indoor trials in Denmark, Canada and the UK (1 to 7 treatments, 0.05 kg ai/hl, a 0- or 1-day PHI) approximated GAP, but a total of 6 results was not sufficient to support an MRL for a major crop. The Meeting agreed to withdraw the previous recommendation for tomato (5 mg/kg).

Witloof chicory (sprouts). The MRL of 1 mg/kg for witloof chicory recommended in 1977 has been supported by trials in France (1987-1992). There was a total of 20 results which could be evaluated, with residues of 0.03-1 mg/kg. The Meeting agreed to maintain the current recommendation of 1 mg/kg.

Lettuce, Head and Leaf. Numerous trial results were available from many countries. Data on head lettuce were from six German greenhouse trials based on 3 treatments, 0.5 kg ai/ha and a 21-day PHI, two outdoor UK trials (2-6 treatments, 0.25 kg ai/ha, a 7-day PHI), two trials from The Netherlands (2 applications, 0.75 kg ai/ha, 10-42-day PHI, indoor) and one French trial (3 treatments, 0.75 kg ai/ha, 23-day PHI, indoor). The total of 24 residues from <0.02 to 9.2 mg/kg were within the range of the previous MRL. The Meeting agreed to maintain the current recommendation of 10 mg/kg for head lettuce.

Data from US trials on leaf lettuce (3 treatments, 1.1 kg ai/ha, 14-day PHI) were made available to the Meeting. The 18 residue values ranged from 0.16 to 22 mg/kg. The Meeting estimated a maximum residue level for leaf lettuce of 25 mg/kg.

Dandelions. Iprodione is authorized for use in France, but there was only one residue trial which was not in line with GAP. An MRL could not be recommended.

Common beans (pods and/or immature seeds). Trials on succulent beans (snap and lima beans) from Canada and the USA approximated GAP. Residue data from 4 Canadian trials (1-2 treatments, 0.75 kg ai/ha) and 15 US trials (2 treatments, 1.1 kg ai/ha) were provided. In a total of 19 values residues were from <0.05 to 1.3 mg/kg. The Meeting estimated a maximum residue level of 2 mg/kg for common bean (pods and/or immature seeds).

Beans (dry). An evaluation of 14 results from trials in Canada and Japan on white and kidney beans showed that after 14-76 days the residues were lower than 0.1 mg/kg. The Meeting estimated a maximum residue level for dry beans of 0.1 mg/kg to replace the previous recommendation (0.2 mg/kg).

Peas. The description of the analyzed commodity in the submitted trials could not be verified and the data were considered inadequate for the estimation of a maximum residue level.

Carrots. There were two residue trials according to GAP from The Netherlands covering pre-storage treatment (1 post-harvest spray, 30.4 g ai/t). As the residues do not decrease during storage but even increase because of the water loss, residues at days 0, 63 and 130 were considered for evaluation. Residues were 4.1-7.9 mg/kg.

Results from pre-harvest trials using foliar sprays were available from the USA with rates

of application (1.3 to 2.25 kg ai/ha) and numbers of treatments (8 to 13) which were somewhat higher than GAP (4 treatments, 0.56 to 1.12 kg ai/ha). The residues, 0.49-3.1 mg/kg, were substantially below those from the post-harvest treatments. The data were consistent among the various post-harvest trials and the Meeting recommended an MRL of 10 mg/kg. The residues from pre-harvest treatments were lower and would be covered by the proposal for post-harvest treatment.

Radishes. Only three residues from GAP treatments were available, from The Netherlands (1 treatment, 2 kg ai/ha, a 12-day PHI). The residues were 1.8-2.2 mg/kg from a single trial. An MRL could not be recommended.

Potatoes. Four results from US trials after foliar spraying (4 treatments, 1.1 to 2.2 kg ai/ha, 14-day PHI) were in conformity with GAP. The residues in the tubers were below the limit of detection (<0.05 mg/kg). As there were only two trials and potatoes are a major crop no MRL is recommended. Post-harvest uses are registered only for seed potatoes; there is no GAP for the post-harvest treatment of potatoes for consumption.

Sugar beet. Iprodione is an authorized product for seed treatment in France and Greece. After seed treatment in UK trials (1.5 kg ai/tonne of seed), there were no residues above the limit of detection (<0.1 mg/kg). An MRL of 0.1* mg/kg is recommended.

Swedes and turnips. Only one trial from the UK corresponded to GAP. The data are not sufficient to estimate a maximum residue level.

Celery. Iprodione is authorized in Australia, but there were only residue values up to 69 mg/kg from US trials which did not correspond to GAP. An MRL could not be recommended.

Cumin Seed. A single trial was inadequate to recommend an MRL.

Barley. Thirteen residue values from UK trials in accordance with GAP (2-3 treatments, 0.5 kg ai/ha, PHI 41-65 days) were <0.1-1.5 mg/kg. After seed treatment according to German GAP, the residues in the grain were less than 0.05 mg/kg. The Meeting estimated a maximum residue level of 2 mg/kg for barley.

Oats. A single trial of a seed treatment was inadequate to recommend an MRL.

Wheat. As with barley, the trials were in the UK (3 treatments, 0.5 kg ai/ha, a PHI of 42 to 74 days). The residues in grain (5 trials, 8 results) were all \leq 0.1 mg/kg. The data were considered inadequate to estimate a maximum residue level for a major crop.

Maize. There were trials in Italy and the USA, but as there was no information on GAP an MRL could not be recommended.

Rice. USA GAP for the use of iprodione on rice permits 2 treatments and a rate of 0.56 kg ai/ha, with the last application not later than when the heading is 75% complete. There were 18 results on husked rice after PHIs of 28 to 58 days. Residues were 0.09-7.1 mg/kg. The Meeting estimated a maximum residue level of 10 mg/kg for rice, husked, to replace the previous recommendation (3 mg/kg).

Almonds. USA GAP for the use of iprodione on almonds permits 4 treatments and a rate of 0.56 mg/kg ai/ha, with the last application within 5 weeks from petal fall. Four trials could be evaluated, which showed that the residues were mainly on the hulls. Those in the nuts were <0.1-0.18 mg/kg. The Meeting estimated a maximum residue level of 0.2 mg/kg for almonds. This would be compatible with residues of 2 mg/kg in almond hulls.

Rape, oilseed. Data from UK trials (residues 0.12-0.43 mg/kg from 1 to 2 treatments at 0.5 kg ai/ha with a 21-day PHI) could be used to recommend an MRL. The Meeting estimated a maximum residue level of 0.5 mg/kg for rape seed.

Sunflower seed. The use of iprodione on sunflower is registered in France. There were five values from France (2 to 3 treatments, 0.4 to 0.7 kg ai/ha, PHI 61-85 days) and 1 value from Italy (2 treatments, 0.75 kg ai/ha, 41-day PHI) that could be used. Residues were <0.04-0.36 mg/kg. The Meeting estimated a maximum residue level of 0.5 mg/kg for sunflower seed.

Coffee. Although iprodione was applied to coffee in two trials, there was no information on GAP. The data were insufficient to recommend an MRL.

Milk, cow tissues. Dairy cattle were treated once daily for 29 days with iprodione at levels corresponding to 5, 15, 50 and 200 ppm in the feed. Milk samples were collected on treatment days 8, 17 and 28 and tissue samples at slaughter. At the 5 ppm feeding level there were no detectable residues of iprodione or its metabolites in milk (<0.01 mg/kg) or tissues (<0.05 mg/kg). At the 15 and 50 ppm feeding levels the total residues in milk were 0.026-0.052 mg/l and in tissues <0.05-0.76 mg/kg. In the absence of detailed information on metabolism the Meeting was unable to recommend MRLs for milk or tissues.

Poultry. Three groups of hens were treated for 28 days by capsule with iprodione at nominal levels of 2, 20 and 100 ppm in the diet or 0.15, 1.5 and 7.5 mg/kg body weight. The total residues in eggs reached a plateau after day 7 of the treatment at 0.1 mg/kg, 0.64 mg/kg and 1.9 mg/kg at the three feeding levels. They had decreased to undetectable levels (<0.01 mg/kg) by day 9 for the 2 and 20 ppm feeding levels and day 12 in the 100 ppm group.

The total residues in liver, muscle, kidney and fat from the 2 ppm feeding level at day 28 were 0.53, <0.05, 0.23 and 0.15 mg/kg respectively, and had decreased to undetectable levels (<0.05 mg/kg) by day 14 of withdrawal at all three feeding levels. In the absence of detailed information on metabolism the Meeting was unable to recommend MRLs for poultry meat or eggs.

Animal feeds. Results of trials on bean fodder and the straw and fodder of cereal grains were available.

Bean forage. Residue data from 13 supervised US trials on snap and Lima bean forage covered a wide range. On the basis of the highest value of 75 mg/kg the Meeting could support a maximum residue level of 100 mg/kg for bean forage (green).

Straw and fodder (dry) of cereal grains. Residue data from supervised trials in the UK on the straw of barley (11 values) and wheat (4 values) were provided. Residues ranged from 0.63 to 5.5 mg/kg. Residues in the grains were of the order of 0.1 mg/kg (wheat) to 1.5 mg/kg (barley). The Meeting agreed the data could support a maximum residue level of 5 mg/kg for the straw and fodder (dry) of

cereal grains.

RECOMMENDATIONS

On the basis of the residue data from supervised trials the Meeting concluded that the residue levels listed in Annex I are suitable for establishing MRLs.

Definition of the residue: iprodione

Commodity		Recommended MRL (mg/kg)		PHI on which based, days
CCN	Name	New	Previous	
TN 0660	Almonds	0.2	-	132-144
FP 0226	Apple	W	10 Po	
GC 0640	Barley	2	-	41-65
VD 0071	Beans (dry)	0.1	0.2	41-76
FB 0264	Blackberries	30	-	0
VB 0400	Broccoli	25	-	0
VR 0577	Carrot	10Po	-	0-130
VP 0562	Common bean (pods &/or immature seeds)	2	-	3-27
FS 0013	Cherries	10	-	1
VC 0424	Cucumber	2	5	1-5
FB 0021	Currants, black, red, white	W	5	
VA 0381	Garlic	W	0.1	
FB 0269	Grapes	10	10	0-30
FI 0341	Kiwifruit	5	5	1
VL 0482	Lettuce, head	10	10	7-42
VL 0483	Lettuce, leaf	25	-	10-14
VA 0385	Onion, bulb	0.2	0.1	7-19
FS 0247	Peach	10	10 Po	0-1
FP 0230	Pear	W	10	
VO 0445	Peppers, sweet	W	5	
FS 0014	Plums (including Prunes)	W	10	

Commodity		Recommended MRL (mg/kg)		PHI on which based, days
CCN	Name	New	Previous	
FP 0009	Pome fruits	5 Po		0-222
SO 0495	Rape seed	0.5	-	21
FB 0272	Raspberries,	30	5	7-8
VR 0596	Sugar beet	0.1*	-	154-182
SO 0702	Sunflower seed	0.5	-	41-85
FB 0275	Strawberry	10	10	0-1
CM 0649	Rice, husked	10	3	28-58
VO 0448	Tomato	W	5	
VS 0469	Witloof chicory (sprouts)	1	1	21-104

Po Post-harvest treatment
 * Limit of determination
 W Withdrawal

REFERENCES

1. Adams, A.M. *et al.* 1988a. Report May and Baker D.Ag.868 of April 1988. Fungicides: Iprodione. Residue studies on cereals (UK 1986). Unpublished.
2. Adams, A.M. *et al.* 1988b. Report May and Baker D.Ag. no 880 of May 1988. Fungicides: Iprodione. Residue studies on cereals (UK 1987). Unpublished.
3. Anon. 1981. Report from Chemical Analysis Consultant, Japan, of 10.05.82. Results of iprodione residue on citrus. Unpublished.
4. Anon. 1986. Report SABS no 17/36/8 of 22.03.82. Iprodione residues in plums. Unpublished.
5. Anon. 1990. Report Department of Agriculture and Rural Affairs no F182-RPT BEZ-PB of 11.01.90. Rovral post-harvest dip for Botrytis control on kiwifruit. Unpublished.
6. Anon. 1993a. Information on GAP, MRLs and residue data on lettuce, endivie, cauliflower, chines cabbage and strawberry by the Delagation of Canada to be considered by the JMPR 1993 (Canada, 1993). Unpublished.
7. Anon. 1993b. Information on GAP, MRLs and residue data on lettuce and rape, oilseed by the Federal Biological Research Centre of Agriculture and Forestry, Braunschweig (Germany, 1993). Unpublished.
8. Anon. 1993c. Information on GAP, MRLs and residue data on commonbeans, carrots, cumin, currants, endivie, fennel, lettuce, kohlrabi, onions, potato, radish, rape seed, witloof and strawberry by the Delagation of The Netherlands to be considered by the JMPR 1993 (Netherlands, 1993). Unpublished.
9. Anon. 1993d. Information on GAP, MRLs by the Ministry of Agriculture, Fisheries to be considered by the JMPR

1993 (New Zealand, 1993). Unpublished.

10. Anon. 1993e. Information on GAP, MRLs and residue data on banana, grapes, strawberry, beans, carrots, celeriac and lettuce by the Delagation of Spain to be considered by the JMPR 1993 (Spain, 1993). Unpublished.

11. Bertolini, P. *et al.* 1985. *Informatore fitopatologico* 12/85 51-54. Verifica dei residui di iprodione nel trattamento post-raccolta delle pomaceae e degli agrumi. Unpublished.

12. Brockelsby, C.H. 1978. Report May and Baker no AR/1379 of March 1978. Iprodione - Residue studies on sprayed raspberries (UK 1977). Unpublished.

13. Brockelsby, C.H. 1981a. Report May and Baker AR/1842 of March 1981. Fungicides: Iprodione. Residue studies in field lettuces (Denmark 1980). Unpublished.

14. Brockelsby, C.H. and Cooper, I. 1981a. Report May and Baker AR/1841 of March 1981. Fungicide: Iprodione. Residue studies on onions (Canada,1979). Unpublished.

15. Brockelsby, C. and Cooper, I. 1981b. Report May and Baker AR/1843 of March 1981. Fungicides: Iprodione. Residue studies in tomatoes (Denmark 1980). Unpublished.

16. Brockelsby, C.H. and Heijbroek, W.M. 1978). Report May and Baker AR/1428 of September 1977. Fungicides: Iprodione Residue studies on Dutch White cabbage (UK 1977 - 1978). Unpublished.

17. Brockelsby, C.H. and Maycey, P.A. 1979). Report May and Baker AR/1507 of March 1979. Fungicide: Iprodione Residue studies on onions (Canada,1978). Unpublished.

18. Brockelsby C.H. and Maycey, P.A. 1982. Report May and Baker no AG Tech 293 of 10.12.82. Fungicides: Iprodione-Residue studies in kiwi fruit (New Zealand 1981/82). Unpublished.

19. Brockelsby, C.H. and Sharpe, J.P. 1981. Report May and Baker AG/Tech/2 of April 1981. Iprodione residue studies on cucumbers (Denmark 1980). Unpublished.

20. Brockelsby, C.H. and Woods, L.S. 1978a. Report May and Baker AR/1378 of March 1978. Fungicides: Iprodione Residue studies on sprayed peaches (Canada 1977). Unpublished.

21. Brockelsby, C.H. and Woods, L.S. 1978b. Report May and Baker AR/1418 of July 1978. Fungicides: Iprodione Residue studies on fresh peaches (South Africa 1978). Unpublished.

22. Brockelsby, C.H. and Woods, L.S. 1978c. Report May and Baker AR/1370 of March 1978. Iprodione residue studies on cucumbers (UK 1977). Unpublished.

23. Brockelsby, C.H. *et al.* 1978a. Report May and Baker AR/1400 of May 1978. Fungicide: Iprodione residue studies on white beans (Canada 1977). Unpublished.

24. Brockelsby, C.H. *et al.*1978b. Report May and Baker AR/1389 of May 1978. Fungicide: Iprodione - Residue studies on rape (Canada 1977). Unpublished.

25. Brockelsby, C.H. *et al.*1978c. Report May and Baker AR/1450 of October 1978. Fungicide: Iprodione - Residue studies on potatoes (UK 1978). Unpublished.

26. Brockelsby, C.H. *et al.*1979a. Report May and Baker AR/1494 of January 1979. Fungicides: Iprodione Residue studies on field lettuces (UK 1978). Unpublished.

27. Brockelsby, C.H. *et al.*1979b. Report May and Baker AR/1574 of October 1979. Fungicides: Iprodione. Residue studies on lettuces (Low volume spray)(UK 1979). Unpublished.

28. Brockelsby, C.H. *et al.* 1979c. Report May and Baker AR/1552 of July 1979. Fungicides: Iprodione Residue studies on Red Cabbage (UK 1978 -1979). Unpublished.
29. Brockelsby, C.H. *et al.* 1979d. Report May and Baker no AR/1515 of March 1979. Fungicide: 26019 RP - Residue studies on cherries (Canada 1978). Unpublished.
30. Brockelsby, C.H. *et al.* 1980a. Report May and Baker AR/1631 of February 1980. Fungicides: Iprodione Residue studies on blackcurrants (UK 1979). Unpublished.
31. Brockelsby, C.H. *et al.* 1980b. Report May and Baker AR/1651 of March 1980. Fungicide: Iprodione Residue studies on onions (Canada,1979). Unpublished.
32. Brockelsby, C.H. *et al.* 1980c. Report May and Baker AR/1655 of March 1980. Fungicide: Iprodione residue studies on white beans (Canada 1979). Unpublished.
33. Buys, M. and Laurent, M. 1974. Report SUCRP-DS an nord no 2534 of 30.08.74 26019 RP. Dosage de residues sur framboises (France 1974). Unpublished.
34. Buys, M. *et al.* 1974a. Report SUCRP-DS An Nord no 2529 of 06.09.74. 26 019 RP. Etude de résidus sur laitues (Grande-Bretagne, 1974). Unpublished.
35. Buys, M. *et al.* 1974b. Report SUCRP-DS An Nord no 2530 of 06.09.74. 26 019 RP. Dosage de résidus sur laitues (France, 1974). Unpublished.
36. Chabassol, Y. and Aublet, J.C. 1983a. Report CRV/CNG An no 4720 of 3.03.83. Iprodione (26019 RP) Résidus sur groseilles (Provenance Pays Bas 1981-82). Unpublished.
37. Chabassol, Y. and Aublet, J.C. 1983b. Report CRV/CNG An no 4744 of 12.04.83. Iprodione (26019 RP) Résidus sur colza (France 1982). Unpublished.
38. Chabassol, Y. and Aublet J.C. 1983c. Report CRV/CNG no 4745 of 12.04.83. Iprodione (26 019 RP)Résidus sur fraises, raisins et tomates (Italie, 1982). Unpublished.
39. Chabassol, Y. and Aublet, J.C. 1983d. Report CRV/CNG An no 4746 of 12.04.83. Iprodione (26019 RP). Résidus sur fraises, haricots et pois (France 1982). Unpublished.
40. Chabassol, Y. and Aublet, J.C. 1983e. Report CRV/CNG An no 4782 of 20.05.83. Iprodione + Carbendazime - Détermination des résidus dans les pommes de terre (France 1982). Unpublished.
41. Chow, W. *et al.* 1991. Report RP Ag Co NWZ/90/08F of 10.10.91. Residues of iprodione in/on kiwi fruit resulting from foliar applications of ROVFAL FLO 250 g/l. Unpublished.
42. Clavière, B. and Guyot, C. 1985a. Report AG/CRLD no 15824.85 of 23.09.85 - Iprodione, formulation Kidan. Résidus dans les raisins - Essais (France 1984). Unpublished.
43. Clavière, B and Guyot, G. 1985b. Report AG/CRLD no 15924.85 of 18.10.85. Iprodione - Formulation Rovral - Résidus dans les groseilles - Essais (Hollande 1984). Unpublished.
44. Clavière, B. and Guyot, C. 1985c. Report AG/CRLD no 16159.85 of 6.12.85. Iprodione - Formulation KIDAN. Résidus dans les carottes Essais (Hollande 1985). Unpublished.
45. Clavière, B. and Muller, M.A. 1988a. Report AG/CRLD no 15131.88 of 20.01.88. Iprodione. Formulation Calidan (SC) et Rovral (WP). Essais France 1987. Résidus dans les pêches. Unpublished.

46. Clavière, B. and Muller, M.A. 1988b. Report AG/CRLD no 15061.88 of 13.01.88. Iprodione. Formulations KIDAN (SC), Calidan (SC) et Rovral (WP). Essais France 1987. Résidus dans les pêches. Unpublished.
47. Dobson, K. *et al.* 1979. Report May and Baker AR/1527 of May 1979. Fungicides: Iprodione Residue studies on peaches (Canada 1978). Unpublished.
48. Dupont, C. and Muller, M.A. 1987. Report AG/CRLD no 15828.87 of 31.07.87. Iprodione. Formulation EXP.2265 (SC). Essais Hollande 1987. Résidus dans les tubercules de pommes de terre. Unpublished.
49. Dupuy, M.G. 1981a. Report CRV/CNG-An no 21220 of 16.11.81. Iprodione Dosage de résidus dans les raisins, le vin et l'alcool (France, 1977-1980). Unpublished.
50. Dupuy, M.G. 1981b. Report C.R. Vitry/ C.N.G. no 21272 of 28.12.81. Iprodione (26019 R.P.). Traitements de post-récolte (trempage, thermonébulisation, cuisson conserve). Essais effectués de 1972 à 1981. Dosage des résidus. Unpublished.
51. Gemma, A, A. *et al.* 1986. Report RP Inc 86/BHL/326 AG (ASD NO 86/196) of May 1986. Iprodione - Aquatic field dissipation and field irrigated crop study. Unpublished.
52. Gillings, O. 1991a. Report RP AG CO no USA88RO8 of 29.07.91. Residues of iprodione and its metabolites in/on tomatoes resulting from aerial and ground applications of Rovral 50 WP. Unpublished.
53. Gillings, O. 1991b. Report RP Ag Co no USA89R16 of 1.10.91. Rovral Onions 1989. Aerial residue program. Unpublished.
54. Gillings, O. 1992. Report RP Ag Co no 41054 of 07.07.92. Rovral Canola (rape seed). 1989 Residue program. Unpublished.
55. Gillings, O. and Shaver, J.A. 1989. Report RPAgCo 40524 (Project no: USA 88R05) of 21.03.89. Residues of Iprodione and its metabolites in/on grapes resulting from commercial application of Rovral 50 WP. Unpublished.
56. Gillings, O. and Shaver, J.A. 1990. Report RPAgCo no 40759 of March 1990. Residues of iprodione and its metabolites in/on peppers resulting from broadcast foliar applications of Rovral 4F. Unpublished.
57. Gillings, O. *et al.* 1984. Report RP Inc no 84/BHL/546/AG (ASD no 84/097) of September 1984. Iprodione residue data on succulent dans dry bean, samples 82/83, Field programs. Unpublished.
58. Gillings, O. *et al.* 1985. Report RP Inc. 85/BHL/483/AG (ASD 85/150) of August 1985. Residue data for broccoli treated with Rovral, 1984, Field Program. Unpublished.
59. Gillings, O. *et al.* 1986a. Report RP Inc 86/BHL/380 AG (ASD no 86/201) of June 1986. Iprodione - Residue report for rice and rice fractions, 1985 field programs E-19 and E-20. Unpublished.
60. Gillings, O.J. *et al.* 1986b. Report RP INC. no 86/BHL/482/AG (ASD no 86/207) of September 1986. Residue data for carrots treated with Rovral. Unpublished.
61. Golner, U. 1991. Report from Ministry of Agriculture Plant Protection Department, Israël, of 10.02.91. Rovral residues in Mineola. Unpublished.
62. Guillet, M. and Muller, M.A. 1986. Report AG/CRLD no 15712.86 of 21.07.86. Iprodione-carbendazime. Formulation Calidan (Suspension concentrée). Résidus dans les grains de colza (France 1985). Unpublished.
63. Guillet, M. *et al.* 1986. Report AG/CRLD no 16013.86 of 27.10.86. Iprodione - formulation Calidan (France 1985) Résidus dans les pêches et les pommes. Unpublished.
64. Guyton, C. 1987a. Report RP Inc no 87/BHL/017/AG (ASD no 87/226) of January 1987. Residue data for potatoe

- tubers, chips, flakes and granules treated at 20 and 40 lb ai/ha with Rovral. Unpublished. Addendum to report by Gemma, A.A.1987). Residues of iprodione and its metabolites (RP30228 and RP32490) in stock feed (potatoe peels) from the processing of potatoes treated with Rovral 50 WP.
65. Guyton, C. 1987b. Report RP Inc. no 87/BHL/056/AG (ASD no 87/227) of February 1987. Residue data for celery treated with Rovral. Unpublished.
66. Guyton, C. 1987c. Report RP Inc 87/BHL/091/AG (ASD no 87/228) of March 1987. Iprodione - Residue data for stored corn and corn fractions. Unpublished.
67. Guyton, C. 1987d. Report RP Inc. 87/BHL/213/AG (ASD no 87/233) of April 1987. Rovral residue data on Chinese cabbage, Chinese mustard and Chinese broccoli. Unpublished.
68. Guyton, C. 1987e. Report RP Inc ref. 87/BHL/250/AG (ASD no 87.235) of May 1987. Rovral residue data for tomato and tomato fractions. Unpublished.
69. Guyton, C. 1987f. Report RP Inc ref. 87/BHL/294/AG no ASD/87-236 of May 1987. Rovral residue data for canberries 1986 Field program E 29. Unpublished.
70. Guyton, C. and Guardigli, A. 1983. Report RP Inc. 83/BHL/860/AG (ASD 83/054) of November 1983. Iprodione, Residue data on California broccoli, Special 1983 Program. Unpublished.
71. Guyton, C. and Guargigli, A. 1984. Report RP Inc no 84/BHL/836/AG (ASD no 84/119) of December 1984. Iprodione residue data on beans and cannery waste fraction 1984, Field programs. Unpublished.
72. Guyton, C. and Guardigli, A. 1985a. Report RP Inc no 85/BHL/111/AG (ASD no 85/126 of March 1985. Iprodione - Residue data on almonds receiving multiple applications of Rovral (1984 Field Program E-2). Unpublished.
73. Guyton, C. and Guardigli, A. 1985b. Report RP Inc no 85/BHL/268/AG (ASD no 85/136) of May 1985. Residue data for potatoes treated with Rovral via Spinkler System. Unpublished.
74. Guyton, C. and Guardigli, A. 1985c. Report RP Inc no 85/BHL/352/AG (ASD no 85/142) of June 1985. Iprodione residue data on Texas onions receiving multiple aerial applications of Rovral, 1985, Field Program E-28. Unpublished.
75. Guyton, C. and Guargigli, A. 1985d. Report RP Inc no 85/BHL/425 AG (ASD 85/147) of July 1985. Residue data for bok choy treated with Rovral (special 1984 Field program). Unpublished.
76. Guyton, C. and Guardigli, A. 1986a. Report RP Inc no 86/BHL/033/AG (ASD no 86/169) of February 1986. Residue data for potatoe tubers, culls and processed fractions following multiple applications of Rovral 1985 Field program. Unpublished.
77. Guyton, C. and Guardigli, A. 1986b. Report RP Inc no 86/BHL/544/AG (ASD no 86/211) of September 1986. Iprodione residue data for leaf lettuce. Field programs FD/84 E-3 and FD/85 E-13. Unpublished.
78. Guyton, C.L. and Guargigli, A. 1986c. Report RP Inc no 84/BHL/398/AG (ASD no 86/203) of July 1986. Iprodione residue data on succulent and dry bean, sample from Michigan and California, 1985, Field programs. Unpublished.
79. Heffernan, P. 1990. Report Rhône-Poulenc LTD no D/AG/1382 of 09.01.90. Fungicide: Rovral-Residue levels from post-harvest dicarboximide fungicide spray application (New Zealand, 1988). Unpublished.
80. Laurent, M. 1975. Report RP/RD/CNG-An no 2854-E of 04.11.75. 26 019 RP. Determination of residues in grapes and wine from Portugal (Portugal, 1974). Unpublished.
81. Laurent, M. 1980. Report RP/RD/CNG-An no 3935 of 11.06.80. Iprodione (26019 RP). Résidus sur prunes (France, 1979). Unpublished.

82. Laurent, M. and Aublet, J.C. 1984. Report ST/CRV/An no 5059 of 26.04.84. Iprodione (26019 RP) Résidus sur colza (Hollande 1983). Unpublished.
83. Laurent, M. and Buys, M. 1975a. Report RP/RD/CNG-An no 2662 of 05.02.75. 26 019 RP Dosages de résidus sur tomates de serre (Grande Bretagne, 1974). Unpublished.
84. Laurent, M. and Buys, M. 1975b. Report RP/RD/CNG-An no 2688 of 13.03.75. 26 019 RP. Dosage de résidus sur raisins et vin d'Italie (Italie, 1974). Unpublished.
85. Laurent, M. and Buys, M. 1975c. Report RP/RD/CNG-An no 2691 of 18.03.75. 26 019 RP. Dosage de résidus sur raisins et vin (Allemagne, 1974). Unpublished.
86. Laurent, M. and Buys, M. 1975d. Report RP/RD/CNG-An no 2734 of 07.05.75. 26 019 RP. Dosage de résidus sur laitues de serre (Grande-Bretagne, hiver 74-75). Unpublished.
87. Laurent, M. and Buys, M. 1975e. Report RP/RD/CNG-An no 2808 of 27.08.75. Iprodione (26 019 RP). Dosage de résidus sur laitues (Lyon, 1975). Unpublished.
88. Laurent, M. and Buys, M. 1975f. Report RP/RD/CNG-An no 2893 of 30.12.75. 2 019 RP. Dosage de résidus sur raisins Provenant d'Espagne (Espagne, 1975). Unpublished.
89. Laurent, M. and Buys, M. 1976a. Report RP/RD/CNG-An no 2926 of 20.02.76. 26 019 RP. Dosage de résidus sur raisins (Portugal, 1975). Unpublished.
90. Laurent, M. and Buys, M. 1976b. Report RP/RD/CNG-An no 2927 of 20.02.76. 26 019 RP. Dosage de résidus sur raisins (Italie, 1975). Unpublished.
91. Laurent, M. and Buys, M. 1976d. Report RP/RD/CNG-An no 2939 of 15.03.76. 26 019 RP. Etude de résidus sur raisins, moût et vin (Allemagne, 1975). Unpublished.
92. Laurent, M. and Buys, M. 1976e. Report RP/RD/CNG An no 2942 of 16.03.76. 26019 RP and 30228 RP - Residues in cucumbers (Japan 1975). Unpublished.
93. Laurent, M. and Buys, M. 1976f. Report RP/RD/CNG-An no 3003 of 22.06.76. Iprodione (26 019 RP). Evolution des résidus de 26 019 RP et de ses métabolites possibles sur laitues provenant des Pays-Bas (essais, 1975). Unpublished.
94. Laurent, M. and Buys, M. 1976g. Report RP/RD/CNG-An no 3047 of 20.10.76. 26019 R.P. and 30228 R.P. Residues in rice (grain and straw) from Japan. Unpublished.
95. Laurent, M. and Buys, M. 1977a. Report RP/RD/CNG-An no 3159 of 28.04.77. Iprodione (26 019 RP). Dosage de résidus sur laitues de serre (France, 1976). Unpublished.
96. Laurent, M. and Buys, M. 1977b. Report RP/RD/CNG An no 3255 of 20.09.77. 26019 RP - Dosage de résidus sur ail (France 1976). Unpublished.
97. Laurent, M. and Buys, M. 1977c. Report RP/RD/CNG-An no 3272 of 17.10.77. Iprodione (26 019 RP). Evolution des résidus sur laitues (Allemagne, 1976/77). Unpublished.
98. Laurent, M. and Buys, M. 1977d. Report RP/RD/CNG-An no 3273 of 17.10.77. Iprodione and Vinchlozoline. Etude comparée des résidus sur laitues (Allemagne, 1977). Unpublished.
99. Laurent, M. and Buys, M. 1977e. Report RP/RD/CNG no 3319 of 15.12.77. Rovral 26019 RP. Residues sur fruits à noyau (France 1977). Unpublished.

100. Laurent, M. and Buys, M. 1977f. Report RP/RD/CNG-An no 3321 of 12.12.77. Iprodione (26 019 RP). Etude comparative de résidus sur raisins après traitement avec une poudre mouillable ou une émulsion huileuse. Unpublished.
101. Laurent, M. and Buys, M. 1978. Report RP/RD/CNG-An no 3372 of 15.03.78. Iprodione. Evolution des résidus sur raisins (Allemagne, 1977). Unpublished.
102. Laurent, M. and Chabassol, Y. 1979a. Report RP/RD/CNG-An no 3671 of 20.07.79. Iprodione (26 019 RP) Résidus sur tomates (France, 1978). Unpublished.
103. Laurent, M. and Chabassol, Y. 1979b. Report RP/RD/CNG-An no 3700 of 21.08.79. Iprodione (26 019 RP). Evolution des résidus sur laitues pommées (Allemagne, 1978). Unpublished.
104. Laurent, M. and Chabassol, Y. 1979c. Report RP/RD/CNG An no 3655 of 20.06.79. Iprodione (26019 RP). Dosage de résidus sur fraises, radis et choux raves (Pays Bas 1978). Unpublished.
105. Laurent, M. and Chabassol, Y. 1979d. Report RP/RD/CNG-An no 3672 of 20.07.79. Iprodione (26 019 RP) Résidus sur laitues et concombres (France, 1978). Unpublished.
106. Laurent, M. and Chabassol, Y. 1979e. Report RP/RD/CNG-An no 3703 of 21.08.79. Iprodione (26019 RP) Evolution des résidus sur choux de Chine (Allemagne, 1978). Unpublished.
107. Laurent, M. and Chabassol, Y. 1980a. Report RP/RD/CNG An no 3825 of 16.01.80. Iprodione (26019 RP) Résidus sur oignons (Pays Bas, 1979). Unpublished.
108. Laurent, M. and Chabassol, Y. 1980b. Report RP/RD/CNG An no 3821 of 16.01.80. Iprodione (26019 RP) Evolution des résidus sur haricots nains. (Pays Bas, 1979). Unpublished.
109. Laurent, M. and Chabassol, Y. 1980c. Report RP/RD/CNG-An no 3853 of 11.02.80. Iprodione (26 019 RP) Evolution des résidus sur laitues pommées (Allemagne, 1979). Unpublished.
110. Laurent, M. and Chabassol, Y. 1980d. Report CRV/CNG An no 3860 of 15.02.80. Iprodione (26019 RP) et carbendazime (21065 RP) Détermination des résidus dans l'orge, l'avoine et le blé (Allemagne, 1979). Unpublished.
111. Laurent, M. and Chabassol, Y. 1980e. Report RP/RD/CNG-An no 3936 of 11.06.80. Iprodione (26 019 RP). Résidus sur raisins (France, 1979). Unpublished.
112. Laurent, M. and Chabassol, Y. 1980f. Report RP/RD/CNG-An no 3971 of 31.07.80. Iprodione. Evolution des résidus sur pommes traitées en post-récolte, avant stockage (station d'Emerainville, 1979-1980). Unpublished.
113. Laurent, M. and Chabassol, Y. 1980g. Report RP/RD/CNG-An no 3972 of 04.08.80. Iprodione (26019 RP). Dosage des résidus dans les pommes traitées après récolte puis entreposées 222 jours. Influence de la cuisson sur les résidus. Unpublished.
114. Laurent, M. and Chabassol, Y. 1980h. Report CRV/CNG An no 4065 of 29.12.80. Iprodione (26019 RP) Résidus sur apricots (France, 1980). Unpublished.
115. Laurent, M. and Chabassol, Y. 1980i. Report CRV/CNG-An no 4067 of 31.12.80. Iprodione. Résidus sur raisins (Italie, 1980) Unpublished.
116. Laurent, M. and Chabassol, Y. 1981a. Report CRV/CNG An no 4070 of 08.01.81. Iprodione (26019 RP) Résidus sur colza (R.F.A. et France, 1980). Unpublished.
117. Laurent, M. and Chabassol, Y. 1981b. Report CRV/CNG An no 4098 of 13.02.81. Iprodione (26019 RP). Evolution des résidus sur laitue (R.F.A., 1980). Unpublished.
118. Laurent, M. and Chabassol, Y. 1981c. Report CRV/CNG An no 4120-E of 03.03.81. Iprodione (26019 RP) and

carbendazim (21065 RP) Determination of residues in barley and wheat (Germany, 1980). Unpublished.

119. Laurent, M. and Chabassol, Y. 1981d. Report CRV/CNG An no 4341 of 30.10.81. Iprodione (26019 RP) Résidus sur colza (France, 1981). Unpublished.

120. Laurent, M. and Chabassol, Y. 1982a. Report CRV/CNG-An no 4434 of 15.02.82. Iprodione. Résidus dans les raisins (Italie, 1981). Unpublished.

121. Laurent, M. and Chabassol, Y. 1982b. Report CRV/CNG An no 11562 of 26.05.82. Iprodione - Dosage de résidus sur bananes de Philippines. Unpublished.

122. Laurent, M. and Chabassol, Y. 1982c. Report CRV/CNG An no 4551 of 21.07.82. Iprodione (26019 RP) Résidus sur colza (R.F.A., 1981). Unpublished.

123. Laurent, M. *et al.* 1974a. Report SUCRP DS An Nord no 2582 of 16.10.74. 26019 RP. Dosage de résidus sur prunes et pruneaux (France, 1974). Unpublished.

124. Laurent, M. *et al.* 1974b. Report SUCRP/DS An Nord no 2596- E of 14.11.74. 26019 RP. Residues studies on Canadian peaches (Canada, 1974). Unpublished.

125. Laurent, M. *et al.* 1974c. Report SUCRP-DS An Nord no 2597 of 25.11.74. 26019 RP. Dosage de résidus sur fraises, framboises et cassis. Unpublished.

126. Laurent, M. *et al.* 1974d. Report SUCRP-DS An Nord no 2604 of 02.12.74. 26 019 RP Determination of residues in grapes (France, 1974). Unpublished.

127. Laurent, M. *et al.* 1975a. Report RP/RD/CNG-An no 2687-E of 13.03.75. 2 019 RP. Residue determination in grapes and wine from Spain. (Spain, 1974). Unpublished.

128. Laurent, M. *et al.* 1975b. Report RP/RD/CNG An no 2720 of April 1975. 26019 RP. Dosage des résidus sur endives (C.E. Ag Emerainville, 1974). Unpublished.

129. Laurent, M. *et al.* 1975c. Report RP/RD/CNG-An no 2733 of 07.05.75. Iprodione (26 019 RP). Dosage de résidus sur laitues de plein champ (France, 1974/75). Unpublished.

130. Laurent, M. *et al.* 1975d. Report RP/RD/CNG-An no 2890 of 30.12.75. 26 019 RP and 30 228 RP. Residues in laitues from Japan. Unpublished.

131. Laurent, M. *et al.* 1976a. Report RP/RD/CNG An no 3077 of 13.12.76. 26019 RP and 30228 RP Residues in kidney beans (Japan, 1976). Unpublished.

132. Laurent, M. *et al.* 1976b. Report RP/RD/CNG/ An no 2944 of 16.03.76. 26019 RP and 30228 RP Residues in tomatoes (Japan, 1975). Unpublished.

133. Laurent, M. *et al.* 1976c. Report RP/RD/CNG-An no 2941 of 16.03.76. 26 019 RP and 30 228 RP. Residues in laitues from Japan. Unpublished.

134. Laurent, M. *et al.* 1976d. Report RP/RD/CNG-An no 2970 of 09.06.76. 26 019 RP and 30 228 RP Residues in tomatoes (Japan, 1975). Unpublished.

135. Laurent, M. *et al.* 1976e. Report RP/RD/CNG An no 2971 of 9.06.76. 26019 RP and 30228 RP Residues in apples (Japan, 1975). Unpublished.

136. Laurent, M. *et al.* 1976f. Report RP/RD/CNG An no 2972 of 9.06.76. 26019 RP and 30228 RP-Residues in apples (Japan, 1975). Unpublished.

137. Laurent, M. *et al.* 1976g. Report RP/RD/CNG no 2975 and 2976 of 09.06.76. 26019 RP and 30228 RP. Residues in peaches (Japan, 1975). Unpublished.
138. Laurent, M. *et al.* 1976 h. Report RP/RD/CNG An no 3049 of 20.10.76. 26019 RP and 30228 RP Residues in kidney beans (Japan, 1975). Unpublished.
139. Laurent, M. *et al.* 1977. Report RP/RD/CNG-An no 3146 of 05.04.77. 26 019 RP. Résidus sur raisins (France, 1975) Unpublished.
140. Laurent, M. *et al.* 1978. Report RP/RD/CNG An no 3331 of January 1978. Iprodione (26019 RP). Dosage des résidus sur endives (Pays Bas, 1977). Unpublished.
141. Laurent, M. *et al.* 1979. Report RP/RD/CNG An no 3670 of 20.07.79. Iprodione (26019 RP) Evolution des résidus sur haricots verts (France, 1978). Unpublished.
142. Laurent, M. *et al.* 1980a. Report RP/RD/CNG An no 3835 of 28.01.80. Iprodione (26019 RP) Résidus sur poires (Italie, 1979). Unpublished.
143. Laurent, M. *et al.* 1980b. Report RP/RD/CNG An no 3934 of 10.06.80. Iprodione (26019 RP) Evolution des résidus sur haricots verts (France, 1979). Unpublished.
144. Laurent, M. *et al.* 1980c. Report RP/RD/CNG-An no 3970 of 31.07.80. Iprodione (26019 R.P.). Résidus sur pommes traitées par trempage avant stockage (France, 1979-1980). Unpublished.
145. Laurent, M. *et al.* 1980d. Report CRV/CNG An no 4062-E of 17.12.80. Iprodione (26019 RP) Residues on bananas (T80 The Philippines, 1980). Unpublished.
146. Laurent, M. *et al.* 1980e. Report CRV/CNG An no 4066 of 31.12.80. Iprodione (26019 RP) Résidus sur poires (Italie, 1980). Unpublished.
147. Laurent, M. *et al.* 1981. Report RP/RD/CNG-An no 4100 of 13.02.81. Iprodione (26019 RP) Evolution des résidus sur chou blanc (R.F.A., 1979 -1980). Unpublished.
148. Laurent, M. *et al.* 1982. Report CRV/CNG-An no 4524 of 17.06.82. Iprodione (26 019 RP). Résidus sur raisins (Chili, 1982). Unpublished.
149. Laurent, M. *et al.* 1984a. Report ST/CRV no 5140 of 12.09.84. Iprodione (26019 RP). Evolution des résidus sur fraises et haricots (France, 1983). Unpublished.
150. Laurent, M. *et al.* 1984b. Report ST/CRV/An no 5138 of 6.09.84. Iprodione (26019 RP) Résidus sur poires (France, 1983). Unpublished.
151. Lusson, R. and Muller, M.A. 1989. Report AG/CRLD no 8915727 of 27.04.89. Iprodione. Formulation Calidan (Sc) Essais France 1988. Résidus dans les graines de tournesol. Unpublished.
152. Malcom, C.P. and P.T. Holland, P.T. 1989). Report from Ministry of Agriculture and Fisheries, New Zealand of 29.09.89 by Iprodione on tangelos. Unpublished.
153. Martens, P.H. and Detroux, L. 1977. Rapport CRP Gembloux no LZ 779-7710-7711-7712 of February 1977. Teneurs résiduelles en glyco-phène dans les chicons. Unpublished.
154. Maycey, P.A. (1980a. Report MAY and BAKER no AR 1680 of 14.05.80. Fungicides: Iprodione-Residue studies in kiwi fruit (New Zealand, 1979/80). Unpublished.
155. Maycey, P.A. 1980b. Report May and Baker AR/1690 of May 1980. Fungicide: Iprodione Residue studies in blackcurrants and the associated juice (UK, 1979). Unpublished.

156. Maycey, P.A. 1980c. Report May and Baker no AR 1693 of June 1980. Fungicides: Iprodione. Residue studies in grapes (Canada, 1979). Unpublished.
157. Maycey, P.A. 1980d. Report May and Baker no AR/1711 of 5.08.80. Fungicide: Iprodione-Residue studies in kiwi fruit (New Zealand, 79/80). Unpublished.
158. Maycey, P.A. 1980e. Report May and Baker no AR/1784 of 5.08.80. Fungicide: Iprodione-Residue studies on peeled kiwi fruit (New Zealand, 79/80). Unpublished.
159. Maycey, P.A. 1980f. Report May and Baker no AR 1753 of November 1980. Iprodione - Residue studies on gooseberries - (UK, 1980). Unpublished.
160. Maycey, P.A. 1981a. Report May and Baker AG/Tech 23 of May 1981. Fungicide: Iprodione Residue studies on onions (Canada, 1980). Unpublished.
161. Maycey, P.A. 1981b. Report May and Baker AR/58 of August 1981. Fungicide: Iprodione - Residue studies on oilseed rape (Canada, 1980). Unpublished.
162. Maycey, P.A. 1982a. Report May and Baker AG Tech 108 of January 1982. Fungicides: Iprodione. Residue studies on oilseed rape (UK, 1981). Unpublished.
163. Maycey, P.A. 1982b. Report May and Baker AG Tech 139 of January 1982. Fungicides: Iprodione. Residue studies on raspberries (UK, 1981). Unpublished.
164. Maycey, P.A. 1982c. Report May and Baker AG Tech 145 of February 1982. Fungicide: Iprodione residue studies in blackcurrants. Flowable formulation (UK, 1981). Unpublished.
165. Maycey, P.A. 1982d. Report May and Baker AG Tech 152 of February 1982. Fungicides: Iprodione Residue studies on blackcurrants and associated juice (UK, 1981). Unpublished.
166. Maycey, P.A. 1982e. Report May and Baker AG/Tech/194 of 1982. Iprodione Residue studies on tomatoes (Denmark, 1981). Unpublished.
167. Maycey, P.A. 1982f. Report May and Baker AG Tech 195 of May 1982. Fungicides: Iprodione. Residue studies on lettuce (Denmark, 1981). Unpublished.
168. Maycey, P.A. 1982g. Report May and Baker AG/Tech/196 of May 1982. Iprodione residue studies on cucumbers (Denmark, 1981). Unpublished.
169. Maycey, P.A. 1982 h. Report May and Baker no AG Tech 248 of August 1982. Iprodione - Residue studies on raspberries (Canada, 1981). Unpublished.
170. Maycey, P.A. 1983a. Report May and Baker no AG Tech 333 of February 1983. Fungicide : 26019 RP - Residue studies on cherries (Canada, 1982). Unpublished.
171. Maycey, P.A. 1983b. Report May and Baker AG/Tech/337 of February 1983. Iprodione residue studies on cucumbers (Canada, 1982). Unpublished.
172. Maycey, P.A. 1983c. Report May and Baker AG/Tech/338 of February 1983. Iprodione Residue studies on tomatoes (Canada, 1981/82). Unpublished.
173. Maycey, P.A. 1983d. Report May and Baker no AG Tech 341 of February 1983. Fungicides: Iprodione. Residue studies on raspberries UK 1982 (Flowable formulation). Unpublished.
174. Maycey, P.A. 1983e. Report May and Baker AG Tech 342 of February 1983. Fungicides: Iprodione. Residue

studies on oilseed rape (UK, 1982). Unpublished.

175. Maycey, P.A. 1983f. Report May and Baker AG Tech 346 of March 1983. Fungicide: Iprodione residue studies on white beans (Canada, 1982). Unpublished.

176. Maycey, P.A. 1983g. Report May and Baker AG Tech 347 of March 1983. Fungicide: Iprodione residue studies on kidney beans (Canada, 1982). Unpublished.

177. Maycey, P.A. 1983h. Report May and Baker AG Tech 348 of March 1983. Fungicide: Iprodione residue studies on snap beans (Canada, 1982). Unpublished.

178. Maycey, P.A. 1983i. Report May and Baker no AG Tech 372 of April 1983. Iprodione - Residue studies on raspberries (Canada, 1982). Unpublished.

179. Maycey, P.A. and Brockelsby, C.H. 1982a. Report May and Baker Ag Tech 200 of May 1982. Fungicides: Iprodione. Residue studies in grapes (Canada, 1980/81). Unpublished.

180. Maycey, P.A. and Brockelsby, C.H. 1982b. Report May and Baker AG Tech 261 of October 1982. Fungicide Iprodione - Residue studies on cereals (UK, 1981-82). Unpublished.

181. Maycey, P.A. and Brockelsby, C.H. 1982c. Report May and Baker AG Tech 262 of October 1982. Fungicide Iprodione - Residue studies on barley (UK, 1981). Unpublished.

182. Maycey, P.A. and Outram J.R. 1986a. Report May and Baker D. Ag. 20 of July 1986. Fungicides: Iprodione. Residue studies in plums (South Africa, 1985-86). Unpublished.

183. Maycey, P.A. and Outram, J.R. 1986b. Report May and Baker Ag. Tech. 1254 of June 1986. Fungicides: Iprodione. Residue studies on swedes and turnips (UK, 1985). Unpublished.

184. Maycey, P.A. and Outram, J.R. 1987a. Report May and Baker D.Ag. 577 of September 1987. Fungicides: Iprodione. Residue studies on swedes (UK, 1986). Unpublished.

185. Maycey, P. A. and Outram, J.R. 1987b. Report May and Baker D.Ag 583 of September 1987. Fungicides: Iprodione. Residue studies on sugar beet (UK, 1986). Unpublished.

186. Maycey, P.A. and Sharpe, J.P. 1984. Report May and Baker no AG.Tech 543 of March 1984. Fungicides: Iprodione residue studies on field beans (UK, 1983). Unpublished.

187. Maycey, P.A. and Sharpe, J.P. 1985a. Report May and Baker AG. Tech. 821 of May 1985. Fungicides: Iprodione. Residue studies on turnips (UK, 1984). Unpublished.

188. Maycey, P.A. and Sharpe, J.P. 1985b. Report May and Baker no AG Tech 876 of July 1985. Fungicides: Iprodione residue studies on field beans (UK, 1984). Unpublished.

189. Mestres, R. 1976. Notes Université de Montpellier of February 1976. Unpublished.

190. Mestres, R. 1982. Essai KIDAN - Iprodione/Pois. Unpublished.

191. Mestres, R. and Tourte, J. 1982. Report CR Vitry-CNG no 317751 of December 1982. Essai Rovral/café no 12-82. Unpublished.

192. Muller, M.A. 1987a. Report AG/CRLD no 15915.87 of 04.08.87. Iprodione-Carbendazime. Formulation EXP. 1926 (SC). Essais France 1986. Résidus dans les pois protéagineux. Unpublished.

193. Muller, M.A. 1987b. Report AG/CRLD no 15886.87 of 24.08.87. Iprodione. Formulation KIDAN (SC). Essais France 1986-1987. Résidus dans les Actinidia (Kiwis). Unpublished.

194. Muller, M.A. 1987c. Report RP/AG/CRLD no 15962.87 of September 1987. Iprodione. Formulation Rovral (WP). Essais France 1986-1987. Résidus dans les endives (racines et chicons). Unpublished.
195. Muller, M.A. 1988a. Report AG/CRLD no 88 15969 of July 1988. Iprodione. Formulation KIDAN (SC). Essais France 1987. Résidus dans les choux-fleurs. Unpublished.
196. Muller, M.A. 1988b. Report AG/CRLD no 88 16070 of July 1988. Iprodione. Formulation Rovral (WP). Essais France 1987. Résidus dans les endives (racines et chicons). Unpublished.
197. Muller, M.A. 1988c. Report AG/CRLD no 8815252 of 09.02.88. Iprodione. Formulation Rovral Aqua Flo. Essais Hollande 1987. Résidus dans les petits pois (grains et gousses). Unpublished.
198. Muller, M.A. 1988d. Report AG/CRLD/An no 88 15966 of 17.06.88. Iprodione-Carbendazime. Formulation Calidan (SC). Essais France 1987. Résidus dans les petits pois (grains et gousse) et les pois protéagineux. Unpublished.
199. Muller, M.A. 1988e. Report AG/CRLD/An no 88 16726 of 10.11.88. Iprodione: Formulation Rovral (WP). Essais France 1988. Résidus dans les haricots verts de conserve. Unpublished.
200. Muller, M.A. 1989a. Report AG/CRLD/AN no 89 16842 of 30.11.89. Diniconazole - Iprodione: Formulation EXP. 02189 (SC). Essai France 1989. Résidus dans la betterave sucrière (racine et feuilles). Unpublished.
201. Muller, M.A. 1989b. Report AG/CRLD/AN no 8916841 of 30.11.89. Iprodione carbendazime. Formulation Calidan (SC). Essais France 1988. Résidus dans les pêches. Unpublished.
202. Muller, M.A. 1989c. Report AG/CRLD/AN no 8916804 of 23.11.89. Iprodione. Formulation KIDAN (SC) Essais France 1989. Residus dans les cerises. Unpublished.
203. Muller, M.A. 1989d. Report AG/CRLD/AN no 8916783 of 20.11.89. Iprodione Carbendazime. Formulation Calidan (SC). Essais France 1989. Résidus dans les cerises. Unpublished.
204. Muller, M.A. 1990a. Report AG/CRLD/AN no 9016278 of 11.09.90 Iprodione. Formulation Rovral FLO (SC). France 1989. Essais traitement après récolte par trempage des fruits (pomme et poire). Unpublished.
205. Muller, M.A. 1990b. Report AG/CRLD/AN no 9016636 of 11.11.90. Iprodione/carbendazime. Formulation Calidan (SC). Essais France 1989. Résidus dans la pêche. Unpublished.
206. Muller, M.A. 1991a. Report AG/CRLD/AN no 9115245 of 11.02.91. Iprodione. Formulation Rovral (WP). Essai Italie 1990. Résidus dans le raisin de cuve (étude de décroissance). Unpublished.
207. Muller, M.A. 1991b. Report AG/CRLD/AN no 9115291 of 19.02.91. Iprodione. Formulation Rovral (WP). Essais France 1989. Résidus dans le raisin de table. Unpublished.
208. Muller, M.A. 1991c. Report AG/CRLD/AN no 9116295 of 30.09.91. Iprodione. Formulation Rovral (WP). Essais Italie 1990. Résidus dans la nectarine. Unpublished.
209. Muller, M.A. 1991d. Report AG/CRLD/AN no 9116303 September 1991. Iprodione. Formulation Rovral (WP). Essais France 1991. Résidus dans le pissenlit. Unpublished.
210. Muller, M.A. 1991e. Report AG/CRLD/AN no 9116329 of 02.10.91. Iprodione. Formulation RHAPSODIE (SC). Essais Hollande 1990. Résidus dans la pomme de terre. Unpublished.
211. Muller, M.A. 1991f. Report AG/CRLD/AN no 9116667 of 05.11.91. Iprodione. Formulation Rovral FLO (SC). Essai France 1991. Résidus dans les petits pois. Unpublished.

212. Muller, M.A. 1991g. Report AG/CRLD/AN no 9116777 of 19.11.91. Iprodione. Formulation Rovral FLO (EXP.1862)-(SC). Essai France 1991. Résidus dans la tomate (Etude de décroissance). Unpublished.
213. Muller, M.A. 1991h. Report AG/CRLD/AN no 9116798 of 21.11.91. Iprodione. Formulation Rovral FLO (EXP.1862) (SC). Essai France 1991. Résidus dans la tomate. Unpublished.
214. Muller, M.A. 1991i. Report AG/CRLD/AN no 9116796 of 21.11.91. Iprodione. Formulation Rovral FLO (SC). Essai France 1991. Résidus dans le concombre (étude de décroissance). Unpublished.
215. Muller, M.A. 1991j. Report AG/CRLD/AN no 9116896 of 4.12.91. Iprodione. Formulation Rovral FLO (SC). Essai France, 1991. Résidus dans l'oignon. Unpublished.
216. Muller, M.A. 1991k. Report AG/CRLD/AN no 9116915 of 06.12.91. Iprodione. Formulation Rovral FLO (SC). Essais France 1991. Résidus dans le melon. Unpublished.
217. Muller, M.A. 1992a. Report AG/CRLD/AN no 9215160 of 11.02.92. Iprodione. Formulation EXP.1862 (SC). Essais France 1991. Résidus dans le haricot vert. Unpublished.
218. Muller, M.A. 1992b. Report AG/CRLD/AN no 9215144 of 11.02.92. Iprodione Formulation EXP.1862 (SC). Essai France 1991. Résidus dans le melon (Etude de décroissance). Unpublished.
219. Muller, M.A. 1992c. Report AG/CRLD/AN no 9215154 of 11.02.92. Iprodione. Formulation EXP.1862 (SC). Essais France 1991. Résidus dans le raisin. Unpublished.
220. Muller, M.A. 1992d. Report R&D/CRLD/AN/bd/9217102 of November 1992. Iprodione. Formulation EXP.1560 ou Rovral (WP). Essai Italie 1991. Résidus dans la pêche (Etude de décroissance). Unpublished.
221. Muller, M.A. 1992e. Report R&D/CRLD/AN/vp/9217119 of November 1992. Iprodione. Formulation EXP.1560 ou Rovral (WP). Essai Italie 1991. Résidus dans la poire (Etude de décroissance) Unpublished.
222. Muller, M.A. 1992f. Report R&D/CRLD/AN/vp/9218090 of December 1992. Iprodione-carbendazime-diniconazole. Formulation Sumistar (SC). Essais France 1990. Résidus dans le tournesol (graine-huile-tourteau). Unpublished.
223. Muller, M.A. 1992g. Report R&D/CRLD/AN/vp/9218119 of December 1992. Iprodione-carbendazime-diniconazole. Formulation Sumistar (SC). Essais France 1991. Résidus dans le tournesol (graine). Unpublished.
224. Muller, M.A. 1993a. Report R&D/CRLD/AN/bd/9315788 of April 1993. Iprodione. Formulation EXP.1862 (SC). Essai France 1991-1992. Résidus dans le chou-fleur. Unpublished.
225. Muller, M.A. 1993b. Report R&D/CRLD/AN/bd/9315821 of April 1993. Iprodione. Formulation EXP 01862 Essais France 1991-1992. Résidus dans l'endive (chicon et racine). Unpublished.
226. Muller, M.A. and Guyot, C. 1985. Report AG/CRLD/An no 15731.85 of 20.08.85. Phosethyl-Al - Iprodione. Formulations Mikal et Rovral. Résidus dans les raisins (Maroc, 1984). Unpublished.
227. Muller, M.A. and Guyot, C. 1987. Report AG/CRLD no 15732-87 of July 1987. Iprodione, formulation KIDAN (SC) Essais France 1986. Residus dans les choux-fleurs. Unpublished.
228. Nangniot, P. 1989. Report CRUPA no 89/328 of 3.05.89. Residus d'Iprodione et de Carbendazime sur pois protéagineux secs. Unpublished.
229. Petrinko, P. and Requier, A. 1977. Report RP/RD/CNG no 19191 of 27.06.77. Rovral 26019 RP - Dosages de résidus dans les légumes. Unpublished.
230. Piznik, M. and Wargo, J.P. 1983. ASD Report No 83/002 Lab Ref No 83/043/BHL/AG. Metabolism of ¹⁴C

Iprodione in rice and the determination of ¹⁴C residues in irrigated crops. Unpublished.

231. Prost, F. 1985. Report RP/CRLD of 27.11.85. Iprodione (26019 RP) residues on kiwi fruit. Unpublished.
232. Robin, J. and Muller, M.A. 1988. Report ref. AG/CRLD no 88 16247 of 09.08.88. Iprodione. Formulation EXP.2265 (SC). Essais Hollande 1987. Résidus dans les tubercules de pommes de terre. Unpublished.
233. Robles, J.M. *et al.* 1987a. Report RPA/AG/CRLD no 15045 of 15.01.87. Iprodione: Formulation Rovral. Résidus dans les graines de tournesol (Italie 1986). Unpublished.
234. Robles, J.M. *et al.* 1987b. Report AG/CRLD no 15309-87 of 19.03.87. Iprodione - formulation Rovral. Essais Italie 1986 - Résidus dans les grains de riz. Unpublished.
235. Robles, J.M. *et al.* 1987c. Report AG/CRLD no 15336-87 of 23.03.87. Iprodione: formulation Rovral. Essais Italie 1986. Résidus dans les grains de maïs. Unpublished.
236. Robles, J.M. *et al.* 1987d. Report AG/CRLD no 15349/87 of 24.03.87. Iprodione-diniconazole. Formulation EXP. 2119 (SC). Essais Hollande 1986. Residus dans le blé (paille et grains). Unpublished.
237. Woods, L.S. 1976a. Report May and Baker RES/2591 of April 1976. Fungicides: 26019 RP. Residue studies on nectarines (Australia, 75/76). Unpublished.
238. Woods, L.S. 1976b. Report May and Baker RES/2588 of April 1976. 26019 RP. Residue studies on plums (Australia, 1975/76). Unpublished.
239. Woods, L.S. 1976c. Report May and Baker RES/2596 of May 1976. Fungicides: 26019 RP. Residue studies on peaches (Australia, 75/76). Unpublished.
240. Woods, L.S. 1977a. Report May and Baker AR/1237 of April 1977. Fungicides: Iprodione Residue studies on Dutch White cabbage (UK, 1976). Unpublished.
241. Woods, L.S. 1977b. Report May and Baker AR/1298 of October 1977. Fungicides: Iprodione Residue studies on glass house lettuces (UK, 1977). Unpublished.
242. Woods, L.S. 1977c. Report May and Baker AR/1338 of December 1977. Fungicides: Iprodione Residue studies on Dutch White cabbage (UK, 1977). Unpublished.
243. Woods, L.S. 1978. Report May and Baker AR/1339 of January 1978. 26019 RP Residue studies on sprayed tomatoes (UK, 1977). Unpublished.
244. Prost, F. 1993a. Iprodione monograph residues JMPR 1993. Rhone Poulenc 23.03.1993. Unpublished.
245. Prost, F. 1992a. Iprodione monograph JMPR 1992. Residue data. Rhone Poulenc 03.04.1992. Unpublished.
246. Prost, F. 1993b. Iprodione monograph residues JMPR 1993. Rhone Poulenc 28.04.1993. Unpublished.
247. Prost, F. 1992b. Iprodione monograph JMPR 1992. Fate of residues. Rhone Poulenc 03.04.1992. Unpublished.
248. Prost, F. 1992c. Iprodione monograph JMPR 1992. Processed food. Rhone Poulenc 03.04.1992. Unpublished.
249. Prost, F. 1992d. Iprodione monograph JMPR 1992. Methods of analysis. Rhone Poulenc 03.04.1992. Unpublished.
250. Laurent, M. and Buys, M. 1978. Report RP/RD/CNG.An no 3346 of February 1978. Iprodione (26019RP) dosage de residus sur laitues (France, 1977). Unpublished.

251. Laurent M. and Buys, M. 1975. Report RP/RD/CNG.An no 2733 of May 1975. Dosage de residus sur laitues de plein champ. Unpublished.
252. Laurent, M. and Chabassol, Y. 1982. Report RP/RD/CNG.An no 4429 of February 1982. Iprodione (26019 RP): Evolution des residus sur haricots verts (France, 1981). Unpublished.
253. Laurent, M. and Chabassol, Y. 1980. Report RP/RD/CNG.An no 3859 of 14.02.19980. Iprodione (26019 RP). Evolution des residus sur haricots verts nains (Germany, 1979). Unpublished.
254. Nangniot, P. 1989. Report CRUPA 89/327 of 03.05.1989. Residus d'iprodione et de carbendazime sur haricots. Unpublished.
255. Laurent, M. and Buys, M. 1976. Report RP/RD/CNG.An no 3048 of 20.10.1976. 26019 R.P. and 30228 R.P. Residues in rice (grain and straw) from Japan. Unpublished.
256. Cappy, J.J. 1992. Report RP Ag Co no 41214 of 09.06.1992. Rovral 4F/Beans/ground/cannery waste/Magnitude of residue. Unpublished.
257. Maycey, P.A. *et al.* 1991. Report RP Agriculture D. Ag.1666 of July 1991. Fungicides: Iprodione. Residue studies in raw and processed potatoes (UK, 1991). Unpublished.
258. Golner, U. and Goldschlag, P. 1992. Report from Ministry of Agriculture Department of Plant Protection and Inspection State of Israel, of March 1992. Iprodione (Rovral). Residues in citrus fruit. Unpublished.
259. Piznik, M. and Wargo, J.P. 1983b. Rhône-Poulenc INC. Lab Ref. 83/088/BHL/AG, ASD No. 83/007. Identification of major unknowns from goat tissues and urine through the metabolism of ¹⁴C-iprodione (RP-26019). Unpublished.
260. Laurent, M. and Chabassol, Y. 1980. Report RP/RD/CNG An no 3898 of 8.04.80. Iprodione (26019RP) Residus sur houblon et biere (Allemagne Federale 1979). Unpublished.
261. Laurent, M. and Chabassol, Y. 1981. Report CRV/CNG An no 4132 of 17.03.81. Iprodione (26019RP) Determination des residus presents dans le houblon et la biere (Allemagne Federale 1980). Unpublished.
262. Chow, W. *et al.* 1982. Report Inc 82/397 BHL/AG ASD 82/039 of July 1982. Iprodione residues in/on grapes and its fractions juice, pomace and raisins. Unpublished.
263. Brockelsby, C.H. *et al.* 1987. Report May and Baker no D. Ag.337 of March 1987. Fungicides: Iprodione. Residue studies on flour and bread (UK, 1986). Unpublished.
264. Wilkes, L.C., Herrera, R.E. and Bache, B. 1982. Research report 317786 vom 14.12.1982 ADC Project 675. Metabolism of ¹⁴C-Iprodione (¹⁴C-RP26019) in laying hens for Rhône-Poulenc Chemical Co. Unpublished.
265. Craig, L.D., Servatius, L., Bache, B. and Herrera, R. 1982. Research Report ADC Project 623. Analysis of milk and tissues of treated dairy cattle for iprodione and its metabolites. Unpublished.
266. Wargo, J.P. and Guardigli, A. 1983. Rhône-Poulenc INC. Lab Ref. 83/547/BHL/AG, ASD No. 83/028. Analysis of tissues and eggs from treated laying hens fed iprodione. Unpublished.
267. Wilkes, L.C. and Herrera, R.E. 1981. Research report ADC Project 543-B. Metabolism of ¹⁴C-iprodione (¹⁴C-RP-26019) in the dairy cow. Unpublished.
268. Wilkes, L.C. and Herrera, R.E. 1982. Research report ADC Project 622. Metabolism of ¹⁴C-iprodione (¹⁴C-RP-26019) in the lactating goat. Unpublished.

269. Gouot, J.M. *et al.* 1977. Report RP/RD/CNG An no 19201-E of June 1977. Pathway and degradation rates of RP 26019 absorbed by roots or applied to plant leaves (strawberries and wheat) with ¹⁴C labelled RP 26019. Unpublished.

270. Gereck, D. *et al.* 1981. Report RP Inc no 81/001 of May 1981. Metabolism study of iprodione in/on peaches. Unpublished.

271. Gouot, J.M. *et al.* 1981. Report CRV/CNG no 21234 E of November 1981. Study of the degradation of iprodione (26019 RP) applied by foliar spray to lettuce under glass, using ¹⁴C labelled iprodione. Unpublished.