

Table 7. Feed formulae (ingredient composition) and proximate composition of commonly used formulated made feeds for different life stages of striped catfish in intensive farming system.^{1,2}

Ingredient/proximate composition (% dry matter)	Life stages/size class			
	Fry	Fingerling	Grower	Broodstock
Ingredient composition (%)				
Rice bran-full fat	23.3	28.1	33.5	37.8
Fishmeal 55%	8	5	5	3
American meat-bone meal 45%	12.0	12.9	12.5	13.9
Defatted soybean meal Arg 47%	35.8	33.2	28.2	19.8
Canola meal	5	5	5	5
Cassava meal	15	15	15	20
Fish oil	0.2	0.1	0.1	0.1
DL-methionine	0.051	0.095	0.069	0.031
Choline chloride 60% China	0.1	0.1	0.1	0.1
Premix-Nutriway	0.5	0.5	0.5	0.3
Proximate composition (%)				
Crude protein	30	28	26	22
Crude lipid	6.21	6.33	6.85	7.14
Ash	10	10	10	10
Crude fiber	5.61	5.84	5.94	5.95
Nitrogen free extract (NFE)	48.2	49.8	51.2	54.9
Gross energy (kJ/g feed)	17.8	17.8	17.8	17.8
Cost (VND/kg)	8 367	7 904	7 657	6 988
Cost (USD/kg)	0.440	0.416	0.403	0.368

¹ Refer to Table 12 for details of commonly used vitamin premix.

² Refer to Table 13 for details of commonly used mineral premix.