

**RISK ASSESSMENT REPORT  
OF THE GENETIC MODIFICATION  
ADVISORY COMMITTEE (GMAC)**

***FOR***

**AN APPLICATION FOR APPROVAL FOR  
RELEASE OF PRODUCTS OF J163  
ALFALFA FOR SUPPLY OR OFFER TO  
SUPPLY**

**NBB REF NO: JBK(S) 600-2/1/17**

**APPLICANT: MONSANTO MALAYSIA  
SDN. BHD.**

**DATE: 12 JANUARY 2022**

## ***I - Summary of Assessment Process***

On 22 September 2021, the Genetic Modification Advisory Committee (GMAC), received from the Department of Biosafety an application for the approval for importation for release [sale/placing on the market for direct use as food, feed and for processing (FFP)] of a product of a Living Modified Organism Glyphosate tolerant J163 alfalfa. The application was filed by Monsanto Malaysia Sdn. Bhd. (hereafter referred to as “the applicant”). After an initial review, GMAC requested for additional information from the applicant.

A public consultation for this application was conducted from 24 February 2021 to 25 March 2021 via advertisements in the local newspapers. Comments were received from Consumers Association of Penang (CAP) and Malaysian Agroecology Society (SRI-Mas). GMAC took into consideration the comments that were relevant to the risk assessment regarding genetic contamination of the environment, and impact on human health such as glyphosate residues, as well as animal toxicity studies. GMAC also took into consideration the comment on the risk and uncertainty associated of the genetic modification technology which may pose unintended consequences and need for proper assessment of J163 alfalfa.

GMAC had four (4) meetings pertaining to this application and prepared the Risk Assessment Report and Risk Assessment Matrix along with its recommended decision, for consideration by the National Biosafety Board.

## ***II - Background of Application***

This application is for approval to import and release products of a Living Modified Organism glyphosate tolerant J163 alfalfa. The aim of the import and release is to supply or offer to supply for sale/placing on the market for direct use as food, feed and for processing (FFP). According to the applicant, J163 alfalfa has been registered in a number of countries for cultivation as well as for food, feed and for processing. J163 alfalfa is approved in a few countries including the United States of America, Australia, New Zealand, Canada, Japan, Korea, Mexico, Philippines, Singapore, Taiwan and Vietnam and may be imported, stored and processed for use in food, animal feed and industrial products in the same way as other conventional, non-transgenic alfalfa. The type of expected use of the products derived from J163 alfalfa in Malaysia will be the same as the expected usage for products derived from conventional alfalfa.

### **Information about J163 alfalfa**

The recipient or parental plant is alfalfa (*Medicago sativa* L.). Alfalfa is among the most important forage crops in the United States and ranks as the fourth most widely grown crop by acreage, after corn, soybean, and wheat. In certain regions, alfalfa is cultivated as a mixture with perennial grasses where it may be harvested as forage or used for grazing livestock. As a legume, it is also

desired for rotational use to improve soil characteristics such as nitrogen content (Undersander *et al.*, 2011).

Many alfalfa plants exhibit various forms of genetic self-incompatibility or self-sterility and will not successfully self-pollinate (Viands *et al.*, 1988). Alfalfa is adversely affected by inbreeding, i.e., self-fertilized plants commonly demonstrate a dramatic reduction in forage and seed yield potential (Rumbaugh *et al.*, 1988). Inbreeding depression may be due to the loss of heterosis and/or accumulation and unmasking of deleterious recessive alleles that occur as a result of self-pollination and/or pollination among close relatives. Flowers do not shed pollen to the wind. After pollination, alfalfa seed requires four to six weeks of adequate growing conditions to ripen. Rainfall during the ripening period will cause poor seed quality and decrease seed yield (Canada Biology Document, 2005).

Alfalfa has a history of minor use as human food, dietary supplements and herbal remedies (OECD 2005). Vast majority of alfalfa is grown and harvested for animal feed (Higginbotham *et al.*, 2008). In Malaysia, alfalfa is used as feed in the form of pellet, dry blocks or bales of hay. For example, pellets are used as feed for rabbits and guinea pigs.

J163 alfalfa contains the *cp4 epsps* gene from *Agrobacterium* sp. strain CP4 that encodes for the 5-enolpyruvylshikimate-3-phosphate synthase (CP4 EPSPS) protein, which confers tolerance to glyphosate, the active ingredient in Roundup® agricultural herbicides.

### **III - Risk Assessment and Risk Management Plan**

GMAC evaluated the application with reference to the following documents:

- (i) CODEX Guideline for the Conduct of Food Safety Assessment of Foods Derived from Recombinant-DNA Plants.
- (ii) Roadmap for Risk Assessment of Living Modified Organisms, (according to Annex III of the Cartagena Protocol on Biosafety produced by the *Ad Hoc* Technical Expert Group (AHTEG) on Risk Assessment and Risk Management of the Convention on Biological Diversity).
- (iii) The risk assessment and risk management plan submitted by the applicant.

GMAC also referred to the following recommendations within the AHTEG guidelines:

- (i) That the risk assessment exercise be specific to the details of this particular application
- (ii) That the risk assessment exercise be specific to the receiving environment in question, and
- (iii) That any risk identified be compared against that posed by the unmodified organism.

In conducting the risk assessment, GMAC identified potential hazards, and then added a value/rank for the likelihood of each hazard as well as its consequences. The likelihood of each hazard occurring was evaluated qualitatively on a scale of 1 to 4, with 1 for 'highly unlikely', and 4 for 'highly likely'. The consequences of each hazard, if it were to occur, were then evaluated on a scale of 1 to 4, with 1 for 'marginal' and 4 to denote a 'major consequence'. A value was finally assigned for the overall risk from the identified potential hazard. The general formula: Overall Risk = Likelihood x Consequence was employed. GMAC also proposed risk management strategies for potential hazards, where appropriate. This methodology of assessment follows the procedure of Risk Assessment in Annex III of the Cartagena Protocol on Biosafety.

The potential hazards were identified in three main areas:

(i) **Effects on human health**

Relevant scientific publications on the genetic modifications were reviewed for potential human health risks and issues pertaining to acute toxicity of novel protein / altering / interference of metabolic pathways, potential allergenicity of the novel protein, reproductive toxicity, potential transfer of antibiotic resistance genes in digestive tract, pathogenic potential of donor microorganisms, nutritional equivalence and anti-nutritional content.

(ii) **Effects on animal health**

Issues pertaining to allergenicity, toxicity, anti-nutritional content, survivability and animal product contamination.

(iii) **Effects on the environment**

Issues pertaining to accidental release of seeds, unintentional release and planting, potential of transgenes being transferred to bacteria (soil bacteria, bacterial flora of animal gut), increased fitness, weediness and invasiveness, accumulation of the protein in the environment via feces from animals fed with the GM plant/grain and cross pollination leading to transfer of transgenes were examined.

Based on the above, a final list of 20 potential hazards was identified. Most of these hazards were rated as having an Overall Risk of 1 or "negligible".

GMAC also took caution and discussed a few of the hazards that required further evaluation and data acquisition. Some of these risks are expected to be managed effectively with the risk management strategies proposed (please refer to section IV of this document).

Some of the potential hazards are highlighted below along with the appropriate management strategies:

**a) Accidental release of viable seeds**

Seeds may be accidentally released during transportation. In the conducive warm and humid climate of Malaysia, there is a high likelihood of these volunteers maturing to the flowering and seed-setting stages. However rainfall causes poor seed quality and decrease, seed yield (Canada Biology Document, 2005). Therefore, the likelihood of a feral population is quite low.

**b) Planting of seeds**

Plants may be grown by uninformed farmers and perpetuated through small scale cultivations. This GM alfalfa may pollinate the non-GM alfalfa. There should also be clear labeling of the product to state that it is only for the purpose of food, feed and processing, and is not to be used as planting material.

**c) Compromised nutritional content**

Compositional analyses of the forage and grain samples showed no significant difference in nutritional composition between J163 alfalfa and conventional alfalfa.

However, applicant is required to update the National Biosafety Board immediately if additional information indicate potential adverse effects from the J163 alfalfa.

#### ***IV - Proposed Terms and Conditions for Certificate of Approval***

Based on the 20 potential hazards identified and assessed, GMAC has drawn up the following terms and conditions to be included in the certificate of approval for the release of this product:

- a) There shall be clear documentation by the exporter describing the product which shall be declared to the Royal Malaysian Customs.
- b) There shall be clear labeling of the product from importation to all levels of marketing stating that it is only for the purpose of food, feed and processing, and is not to be used as planting material.
- c) Should the approved person receive any information that indicates any adverse effect of J163 alfalfa, the National Biosafety Board shall be informed immediately.
- d) Any spillage (during loading/unloading/transportation) shall be collected and cleaned up immediately.
- e) Transportation of the consignment from the port of entry to any destination within the country shall be in secured and closed condition.

## ***V - Other Regulatory Considerations***

- a) Administrative regulatory procedures shall be arranged between the Department of Biosafety, Royal Malaysian Customs Department and relevant agencies to ensure accurate declaration of product information and clear labeling of the product is implemented.
- b) Administrative regulatory procedures shall be arranged between the Department of Biosafety and the Malaysian Quarantine and Inspection Services (MAQIS) to impose post entry requirements for accidental spillage involving the GM product.
- c) Administrative regulatory procedures shall be arranged between the Department of Biosafety and the Malaysian Quarantine and Inspection Services (MAQIS) and other competent agencies to impose post entry requirements for food safety compliance.
- d) Administrative regulatory arrangements shall be carried out between the Department of Biosafety and the Department of Veterinary Services (DVS) so that any unanticipated adverse effects in animals caused by any consumption of the GM products shall be reported immediately.
- e) Administrative regulatory arrangements shall be carried out by Food Safety and Quality of Ministry of Health to monitor compliance to the Food Regulations 1985 for labelling of GM food.
- f) Administrative regulatory procedures shall be arranged between Department of Biosafety and Ministry of Health to ensure that herbicide residues in alfalfa consignments are below the acceptable maximum residual level established. It is recommended that importers are required to provide certificate of analysis for herbicide residues prior to shipment.

## ***VI - Identification of issues to be addressed for long term use release of this product***

- a) Continuous monitoring is required from the approved person and any unanticipated adverse effect caused by the J163 alfalfa shall be reported to the National Biosafety Board.

## ***VII –Conclusion and Recommendation***

GMAC has conducted a thorough evaluation of the application for approval for importation for release [sale/placing on the market for direct use as food, feed and for processing (FFP)] of a product of a Living Modified Organism glyphosate tolerant J163 alfalfa and has determined that the release of this product does not endanger biological diversity or human, animal and plant health. GMAC recommends that the proposed application for release be **APPROVED WITH TERMS AND CONDITIONS** as listed in section IV - Proposed Terms and Conditions for Certificate of Approval.

## VIII – Bibliography

1. Bora, K.S. and A. Sharma. 2011. Phytochemical and pharmacological potential of *Medicago sativa*: A review. *Pharmaceutical Biology* 49:211-220.
2. Caetano-Anollés, G., M. Wang, D. Caetano-Anollés and J.E. Mittenenthal. 2009. The origin, evolution and structure of the protein world. *Biochemical Journal* 417:621-637.
3. CFIA. 2005. The Biology of *Medicago sativa* L. (Alfalfa). Canadian Food Inspection Agency, Ottawa, Ontario. <https://inspection.canada.ca/plant-varieties/plants-with-novel-traits/applicants/directive-94-08/biology-documents/medicago-sativa-l-/eng/1330981151254/1330981232360>
4. CFIA. 2007. Code of practice for the hygienic production of sprouted seeds. Canadian Food Inspection Agency, Ottawa, Ontario. <http://www.inspection.gc.ca/english/fssa/frefra/safsal/sprointe.shtml> [Accessed May 18, 2012].
5. CFIA. 2010. The Biology of *Medicago sativa* L. (Alfalfa). Canadian Food Inspection Agency, Ottawa, Ontario.
6. Church, D.C. 1984. Table of contents. Pages iii-iv in *Livestock Feeds and Feeding*. Second Edition. O & B Books, Inc., Corvallis, Oregon.
7. Codex Alimentarius. 2009. Foods derived from modern biotechnology. Second Edition. Codex Alimentarius Commission, Joint FAO/WHO Food Standards Programme, Food and Agriculture Organization of the United Nations, Rome, Italy.
8. Conrad, H.R. and T.J. Klopfenstein. 1988. Role in livestock feeding-Greenchop, silage, hay, and dehy. Pages 539-551 in *Alfalfa and Alfalfa Improvement*. A.A. Hanson, D.K. Barnes, and R.R. Hill (eds.). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, Wisconsin.
9. Dill GM, CaJacob CA, Padgette SR (2008) Glyphosate-resistant crops: adoption, use and future considerations. *Pest Manag Sci* 64:326–331.
10. Duke, J.A., M.B. Forde, C.F. Reed and J.K.P. Weder. 1981. *Medicago sativa* L. Pages 157- 161 in *Handbook of Legumes of World Economic Importance*. J.A. Duke (ed.). Plenum Press, New York, New York.
11. FAO-WHO. 1991. Strategies for assessing the safety of foods produced by biotechnology. Report of joint FAO/WHO consultation. World Health Organization, Geneva, Switzerland.
12. FAO-WHO. 2001. Evaluation of allergenicity of genetically modified foods. Report of a joint FAO/WHO expert consultation on allergenicity of foods derived from biotechnology. Food and Agriculture Organization of the United Nations, Rome, Italy.
13. Hammond, B.G. and R.L. Fuchs. 1998. Safety evaluation for new varieties of food crops developed through biotechnology. Pages 61-79 in *Biotechnology and Safety Assessment*. Second Edition. J.A. Thomas (ed.). Taylor & Francis, Philadelphia, Pennsylvania.
14. Harrison, L.A., M.R. Bailey, M.W. Naylor, J.E. Ream, B.G. Hammond, D.L. Nida, B.L. Burnette, T.E. Nickson, T.A. Mitsky, M.L. Taylor, R.L. Fuchs and S.R. Padgette. 1996. The expressed protein in glyphosate-tolerant soybean, 5-enolpyruvylshikimate-3-phosphate

- synthase from *Agrobacterium* sp. strain CP4, is rapidly digested in vitro and is not toxic to acutely gavaged mice. *Journal of Nutrition* 126:728-740.
15. Haslam, E. 1993. *Shikimic acid : metabolism and metabolites*. Wiley Chichester; New York.
  16. Higginbotham, G.E., C.L. Stull, N.G. Peterson, A.V. Rodiek, B.A. Reed and J.N. Guerrero. 2008. Alfalfa utilization by livestock. Pages 265-279 in *Irrigated Alfalfa Management for Mediterranean and Desert Zones*. C.G. Summers and D.H. Putnam (eds.). University of California Agriculture and Natural Resources, Oakland, California.
  17. Hileman, R.E., A. Silvanovich, R.E. Goodman, E.A. Rice, G. Holleschak, J.D. Astwood and S.L. Hefle. 2002. Bioinformatic methods for allergenicity assessment using a comprehensive allergen database. *International Archives of Allergy and Immunology* 128:280-291.
  18. Holden, L.A., B.P. Glenn, R.A. Erdman and W.E. Potts. 1994. Nutrition, feeding, and calves: Effects of alfalfa and orchardgrass on digestion by dairy cows. *Journal of Dairy Science* 77:2580-2594.
  19. Howarth, R.E. 1988. Antiquality factors and nonnutritive chemical components. Pages 493- 514 in *Alfalfa and Alfalfa Improvement*. A.A. Hanson, D.K. Barnes, and R.R. Hill (eds.). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, Wisconsin.
  20. Illergård, K., D.H. Ardell and A. Elofsson. 2009. Structure is three to ten times more conserved than sequence - A study of structural response in protein cores. *Proteins* 77:499- 508.
  21. Julier, B., C. Huyghe and C. Ecalte. 2000. Within- and among-cultivar genetic variation in alfalfa: Forage quality, morphology, and yield. *Crop Science* 40:365-369.
  22. Keese P. Risks from GMOs due to horizontal gene transfer. *Environmental Biosafety Research*. 2008;7(3):123–149.
  23. Klee et al., 1987. *Agrobacterium*-mediated plant transformation and its further applications to plant biology. *Plant Physiology* 38: 467-486
  24. Kumar, K., Gambhir, G., Dass, A. et al. Genetically modified crops: current status and future prospects. *Planta* 251, 91 (2020)
  25. Marten, G.C., D.R. Buxton and R.F. Barnes. 1988. Feeding value (Forage quality). Pages 463-491 in *Alfalfa and Alfalfa Improvement*. A.A. Hanson, D.K. Barnes, and R.R. Hill (eds.). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, Wisconsin.
  26. Metcalfe, D.D., J.D. Astwood, R. Townsend, H.A. Sampson, S.L. Taylor and R.L. Fuchs. 1996. Assessment of the allergenic potential of foods derived from genetically engineered crop plants. *Critical Reviews in Food Science and Nutrition* 36:S165-S186.
  27. Michaud, R., W.F. Lehman and M.D. Rumbaugh. 1988. World distribution and historical development. Pages 25-91 in *Alfalfa and Alfalfa Improvement*. A.A. Hanson, D.K. Barnes, and R.R. Hill (eds.). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, Wisconsin.



28. NRC. 1982. Feed composition tables. Pages 8-124 in United States-Canadian Tables of Feed Composition: Nutritional Data for United States and Canadian Feeds. Third Revision Edition. National Research Council, National Academy Press, Washington, D.C.
29. OECD. 2005. Consensus document on compositional considerations for new varieties of alfalfa and other temperate forage legumes: Key feed nutrients, anti-nutrients and secondary plant metabolites. ENV/JM/MONO (2005)13. Organisation for Economic Co-operation and Development, Paris, France.
30. Pariza, M.W. and E.A. Johnson. 2001. Evaluating the safety of microbial enzyme preparations used in food processing: Update for a new century. *Regulatory Toxicology and Pharmacology* 33:173-186.
31. Putnam, D.H., P. Robinson and E. DePeters. 2008. Forage quality and testing. Pages 241-264 in *Irrigated Alfalfa Management for Mediterranean and Desert Zones*. C.G. Summers and D.H. Putnam (eds.). University of California Agriculture and Natural Resources, Oakland, California.
32. Quiros, C.F. and G.R. Bauchan. 1988. The genus *Medicago* and the origin of the *Medicago sativa* complex. Pages 93-124 in *Alfalfa and Alfalfa Improvement*. A.A. Hanson, D.K. Barnes, and R.R. Hill (eds.). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., Madison, Wisconsin.
33. Rumbaugh, M. D., Caddel, J. L., and Rowe, D. E. 1988. Breeding and quantitative genetics. In: *Alfalfa and Alfalfa Improvement*. pp. 777–808. Hanson, A.A., Barnes, D. K. and Hill, Jr. R. R., Eds. American Society of Agronomy, Inc. Publishers, Madison, Wisconsin, USA.
34. Schönbrunn, E. *et al.* 2001. Interaction of the herbicide glyphosate with its target enzyme 5-enolpyruvylshikimate 3-phosphate synthase in atomic detail. *Proceedings of the National Academy of Sciences of the United States of America*. 98(4):1376-1380
35. Silvanovich, A., M.A. Nemeth, P. Song, R. Herman, L. Tagliani and G.A. Bannon. 2006. The value of short amino acid sequence matches for prediction of protein allergenicity. *Toxicological Sciences* 90:252-258.
36. Sjoblad, R.D., J.T. McClintock and R. Engler. 1992. Toxicological considerations for protein components of biological pesticide products. *Regulatory Toxicology and Pharmacology* 15:3- 9.
37. Smith, D. 1969. Influence of temperature on the yield and chemical composition of 'Vernal' alfalfa at first flower. *Agronomy Journal* 61:470-472.
38. Steinrücken HC, Amrhein N. 1984. 5-Enolpyruvylshikimate-3-phosphate synthase of *Klebsiella pneumoniae*. 1. Purification and properties. *Eur J Biochem*;143(2):341-9. doi: 10.1111/j.1432-1033.1984.tb08378.x.PMID: 6381056
39. Thomas, K., M. Aalbers, G.A. Bannon, M. Bartels, R.J. Dearman, D.J. Esdaile, T.J. Fu, C.M. Glatt, N. Hadfield, C. Hatzos, S.L. Hefle, J.R. Heylings, R.E. Goodman, B. Henry, C. Herouet, M. Holsapple, G.S. Ladics, T.D. Landry, S.C. MacIntosh, E.A. Rice, L.S. Privalle, H.Y. Steiner, R. Teshima, R. van Ree, M. Woolhiser and J. Zawodny. 2004. A multilaboratory evaluation of a common in vitro pepsin digestion assay protocol used in assessing the safety of novel proteins. *Regulatory Toxicology and Pharmacology* 39:87-98.

40. Thomas, K., G. Bannon, S. Hefle, C. Herouet, M. Holsapple, G. Ladics, S. MacIntosh and L. Privalle. 2005. In silico methods for evaluating human allergenicity to novel proteins: International Bioinformatics Workshop Meeting Report, 23-24 February 2005. *Toxicological Sciences* 88:307-310.
41. Townsend, M.S., J.A. Henning, D.W. Smith, I.M. Ray and C.G. Currier. 1998. Differential concentration of ten minerals among four ancestral alfalfa germplasms. *Crop Science* 39:574- 578.
42. Undersander, D., D. Cosgrove, E. Cullen, C. Grau, M. Rice, M. Renz et al. 2011. Alfalfa management guide. <https://www.agronomy.org/files/publications/alfalfa-management-guide.pdf>
43. U.S. EPA. 1996. Plant pesticide inert ingredient CP4 enolpyruvylshikimate-3-D and the genetic material necessary for its production in all plants. *Federal Register* 61:40338-40340.
44. U.S. FDA. 1992. Statement of policy: Foods derived from new plant varieties. *Federal Register* 57:22984-23005.
45. Viands, D. R., Sun, P., and Barnes, D. K. 1988. Pollination control: Mechanical and sterility. In: *Alfalfa and Alfalfa Improvement*. pp. 931–960. Hanson, A.A., Barnes, D. K. and Hill, Jr. R. R., Eds. American Society of Agronomy, Inc. Publishers, Madison, Wisconsin, USA.

**GENETIC MODIFICATION ADVISORY COMMITTEE (GMAC) MEMBERS INVOLVED IN  
SPECIFIC RISK ASSESSMENT AREAS FOR THE APPROVAL FOR RELEASE OF  
PRODUCTS OF J163 ALFALFA FOR SUPPLY OR OFFER TO SUPPLY**

Genetic Modification Advisory Committee (GMAC) members divided the task of looking up more information for the Risk Assessment matrix based on three broad categories which were environment, human health and animal health. Each sub-committee had a nominated leader to coordinate the work and report back to the main GMAC. The GMAC members involved in the risk assessment are as below:

- **Prof. Dr. Mohd. Faiz Foong bin Abdullah (Universiti Teknologi MARA) (GMAC Chairman)**
- **Dr. Kodi Isparan Kandasamy (Industry Representative) (Environment sub-committee Leader)**
- **Madam T.S. Saraswathy (Institute of Medical Research - retired) (Human Health sub-committee Leader)**
- **Prof. Dr Jothi Malar Panandam (Universiti Putra Malaysia - retired) (Animal Health sub-committee Leader)**
- **Dr. Rahizan Issa (Institute of Medical Research - retired) (Notification Assessment sub-committee Leader)**
- Dato' Dr. Sim Soon Liang (Academy of Sciences Malaysia)
- Prof. Dr. Abd Rahman Milan (Universiti Malaysia Sabah - retired)
- Prof. Dr. Chan Kok Gan (Universiti Malaya)
- Assoc. Prof. Dr. Choong Chee Yen (Universiti Kebangsaan Malaysia)
- Dr. Adiratna Mat Ripen (Institute of Medical Research)
- Dr. Norliza Tendot Abu Bakar (Malaysian Agricultural Research & Development Institute)
- Dr. Norwati Muhammad (Forest Research Institute of Malaysia)
- Dr. Saifullizam bin Abdul Kadir (Department of Veterinary Services)
- Dr. Teo Tze Min (Entomological Society of Malaysia)
- Dr. Mohd Hefni Rusli (Malaysian Palm Oil Board)
- Madam Shafini Abu Bakar (Ministry of Health)
- Madam Sabariah Kamis (Department of Agriculture)
- Mr. Harun bin Ahmad (Department of Chemistry Malaysia)