

EUROPEAN INLAND FISHERIES ADVISORY COMMISSION

Report of the

**EIFAC AD HOC WORKING PARTY ON HANDLING OF FISHES IN FISHERIES
AND AQUACULTURE**

Utrecht, Netherlands, 24–26 March 2004



Copies of FAO publications can be requested from:

Sales and Marketing Group

Communication Division

FAO

Viale delle Terme di Caracalla

00153 Rome, Italy

E-mail: publications-sales@fao.org

Fax: +39 06 57053360

Web site: <http://www.fao.org>

EUROPEAN INLAND FISHERIES ADVISORY COMMISSION

Report of the

EIFAC AD HOC WORKING PARTY ON HANDLING OF FISHES IN FISHERIES AND AQUACULTURE

Utrecht, Netherlands, 24-26 March 2004

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

ISBN 978-92-5-106016-2

All rights reserved. Reproduction and dissemination of material in this information product for educational or other non-commercial purposes are authorized without any prior written permission from the copyright holders provided the source is fully acknowledged. Reproduction of material in this information product for resale or other commercial purposes is prohibited without written permission of the copyright holders. Applications for such permission should be addressed to:

Chief
Electronic Publishing Policy and Support Branch
Communication Division
FAO
Viale delle Terme di Caracalla, 00153 Rome, Italy
or by e-mail to:
copyright@fao.org

© FAO 2008

PREPARATION OF THIS DOCUMENT

The handling of fishes is increasingly subject to legislation and regulation. In 2002, the twenty-second session of the European Inland Fisheries Advisory Commission of FAO (EIFAC) realized the need to identify current legislation and regulations in the EIFAC member countries. Also, it was felt there was a need to exchange information on the practice of handling fishes in fisheries and aquaculture, e.g. catching and handling, husbandry conditions, transport and slaughter of fishes. The session established an EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture as a forum to exchange information about the handling of fishes for those working in fisheries and aquaculture. The terms of reference of the ad hoc Working Party are presented in the Preface to this document.

This EIFAC Occasional Paper contains two parts. The first part provides the report of the EIFAC Workshop on Handling of Fishes in Fisheries and Aquaculture, which was held in Utrecht, Netherlands, from 24 to 26 March 2004. The second part provides an overview of the handling of fishes in commercial inland fisheries, recreational fisheries, aquaculture and fisheries research, compiled by Mr A.J.P. Raat (Dutch Organization for the Improvement of Inland Fisheries (OVV) and Mr P. van de Nieuwegiessen (Wageningen University Research).

In March 2003, Mr A.J.P. Raat, the convener of the ad hoc Working Party, sent an inquiry on the current practice of handling of fishes in EIFAC member countries to EIFAC National Correspondents. The inquiry was designed to compile data on handling of fishes from EIFAC member countries. The results of the inquiry are included in this report. The convener is grateful to the following persons for their reaction to the questionnaire: U.A. Grosch (Berlin, Germany), H. Wedekind (Groß Glienicke, Germany), M. Pietrock (Groß Glienicke, Germany), E. Dupont (Marloie, Belgium), J.W. van de Vis (IJmuiden, Netherlands), E. Lambooy (Lelystad, Netherlands), P. Hickley (Kidderminster, United Kingdom), N. Tredwin (Weymouth, United Kingdom), R. Poole and colleagues (Newport, Ireland).

The revision of the draft version of the overview was carried out by Mr A.J.P. Raat. The document was circulated for review to the participants of the Utrecht workshop. The contributions to the present document by the participants of the Utrecht workshop are acknowledged with great appreciation. Finally, Mr P. Hickley (Vice-chairperson of EIFAC), Mr I. Cowx (Chairperson of EIFAC Sub-Commission IV), Mr R. Roesch (Langenargen, Germany) and Mr B.P. North (Stirling, Scotland) reviewed the document. The convener acknowledges the input of Mr H. van de Vis (IJmuiden, Netherlands). The comments and suggestions of the above mentioned persons are gratefully acknowledged by the convener of the Working Party.

The report of the Utrecht Workshop of the EIFAC ad hoc Working Party was presented in June 2004 to the Twenty-third Session of EIFAC in Wierzba, Poland. It was decided to publish the report of the workshop and the overview of the handling of fishes as an EIFAC Occasional Paper. The overview of the handling of fishes in fisheries and aquaculture is based on a technical document of the Dutch Organisation for the Improvement of Inland Fisheries (the Organisatie ter Verbetering van de Binnenvisserij OVB), Nieuwegein, Netherlands. OVB offered the facilities for the activities of the Working Party.

FAO/European Inland Fisheries Advisory Commission.
Report of the EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture.
Utrecht, Netherlands, 24–26 March 2004.
EIFAC Occasional Paper. No. 40. Rome, FAO. 2008. 88p.

ABSTRACT

This document presents the results of the EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture, including the report of its workshop held from 24 to 26 March 2004 in Utrecht, Netherlands. The welfare of fishes has become a subject of attention in recent years. A concise definition of welfare is inherently difficult and instead, definitions are feelings-based, function-based and nature-based. However, none of these three concepts offers a comprehensive view on the major ethical concerns over the quality of life of animals. Concerns have been raised about the potential for infringed fish welfare in fisheries, aquaculture and fisheries research. Most of these concerns have been related to farm escapees, post-release mortality, and some capture methods which have been shown to induce stress responses. Fish welfare problems in recreational fisheries deal with the physiological effects of angling and subsequent hooking mortality. Several factors may influence fish welfare during angling: physiological factors, environmental factors and fishing method factors. Husbandry practices in aquaculture have the potential to impact on the welfare of cultured fish including: handling, inappropriate stocking density, confinement, transportation, water quality deterioration, social interaction, light regime, slaughter. Many issues also apply to the use of fish in scientific investigations. Here, fish are commonly held in captivity for experimentation and fish are caught or sampled in surveys and monitoring programmes. Fish are being tagged using invasive methods. Concerns have been raised about the effects of electric fishing techniques, commonly used to sample fish from fresh waters. Indicators often used to assess the welfare status of an animal are stress physiology, animal behaviour and health status. There are similarities and differences between fishes and mammals with regard to the subjective experience of pain and suffering.

Independent of the question on pain and suffering of fishes, members of the ad hoc Working Party agreed that handling of fishes must be carried out in a responsible way. Fishes show reactions to injurious stimuli. These stimuli result in physiological stress responses. Depending on the duration and intensity of the exposure to the stressor, the stress response varies from acute to chronic. Handling causing injuries and damage to the fishes should be avoided in fisheries, aquaculture and fisheries research. Careful handling of the fishes implies prevention of damage or unnecessary injuries, prevention of unnecessary stress and prevention of unnecessary prolongation of the period before the target fish is killed. Those responsible for handling of fishes in fisheries, aquaculture and fisheries research should have basic knowledge of the effects of handling on fishes and have justifiable objectives (“reasonable purpose”), within the context of the situation. Furthermore the rules and regulations with regard to these activities must be taken into account. To ensure the welfare of fish in commercial fisheries, recreational fisheries and aquaculture, sufficient legislation and regulation is necessary. All EIFAC member countries discussed in this report have legislation and regulations concerning animal welfare. However, some of the legislation and regulations do not provide specific attention to fish welfare.

Key words: Handling, Fish Welfare, Inland Capture Fisheries, Aquaculture, EIFAC, European Inland Fisheries Advisory Commission, Europe

CONTENTS

	Page
Preparation of this document	iii
Abstract	iv
Preface	viii
 PART ONE	 1
REPORT OF THE EIFAC WORKSHOP ON HANDLING OF FISHES IN FISHERIES AND AQUACULTURE	1
INTRODUCTION	1
AIMS OF THE EIFAC WORKSHOP	1
OUTPUT OF THE EIFAC WORKSHOP	1
GENERAL CONSIDERATIONS	2
WELFARE AND FISHES	3
PAIN AND SUFFERING OF FISHES	4
RESPONSIBLE HANDLING OF FISHES	5
HANDLING OF FISHES: FISHERIES RESEARCH	5
HANDLING OF FISHES IN PRACTICE: FISHES IN AQUACULTURE	5
HANDLING OF FISHES IN PRACTICE: ORNAMENTAL FISHES	6
HANDLING OF FISHES IN PRACTICE: FISHERIES MANAGEMENT	6
RELEVANT INFORMATION	7
REFERENCES	7
ANNEX List of participants	8
 PART TWO	 11
HANDLING OF FISHES IN COMMERCIAL INLAND FISHERIES, RECREATIONAL FISHERIES, AQUACULTURE AND FISHERIES RESEARCH. AN OVERVIEW	11
CHAPTER 1 FISH WELFARE	11
1.1 Defining animal welfare	11
1.2 Criteria for fish welfare	11
1.2.1 Five freedoms	11
1.2.2 Discussion on perception of pain by fishes	12
1.3 Indications of fish welfare	16
1.3.1 Stress physiology	17
1.3.2 Behaviour	17
1.3.3 Health status	18
1.3.4 Pathological changes	18
1.3.5 Product quality	18
CHAPTER 2 EFFECTS OF FISHERIES	19
2.1 General aspects of commercial inland fisheries	19
2.2. Actions for commercial inland fisheries	20
2.3 Fisheries management	20
2.4 Catch and release fisheries	21
2.5 Physiological effects	21
2.6 Environmental effects	22
2.7 Effects of fishing methods	23
CHAPTER 3 EFFECTS OF AQUACULTURE	24
3.1 Handling	24
3.2 Stocking density	24
3.3 Water quality	25

3.3.1	Dissolved oxygen	25
3.3.2	Ammonia	25
3.3.3	Effects of deterioration in water quality on welfare	26
3.3.4	Effect of environmental factors on water quality deterioration	26
3.4	Social interaction	27
3.5	Light regime	28
3.6	Slaughter	28
CHAPTER 4 FISH IN SCIENTIFIC RESEARCH		31
4.1	Research and monitoring of fish and fish ecology	31
4.2	Fields of interest	31
4.3	Electric fishing	31
CHAPTER 5 LEGISLATION AND REGULATION		33
5.1	International legislation and regulation	33
5.1.1	European Union	33
5.1.2	European Food Safety Authority (EFSA)	34
5.1.3	Council of Europe	34
5.1.4	World Organisation for Animal Health (OIE)	35
5.2	National legislation and regulation	36
5.2.1	Austria	36
5.2.2	Finland	36
5.2.3	France	36
5.2.4	Germany	37
5.2.5	Ireland	37
5.2.6	Netherlands	38
5.2.7	Norway	38
5.2.8	Poland	38
5.2.9	Sweden	39
5.2.10	Switzerland	39
5.2.11	United Kingdom	39
5.2.12	Denmark	41
CHAPTER 6 CURRENT CODES AND GUIDELINES		42
6.1	Aquaculture	42
6.2	Commercial fisheries	42
6.3	Angling	42
6.4	Ornamental fish	43
CHAPTER 7 CITED REFERENCES		44
CHAPTER 8 AVAILABLE INFORMATION ON THE INTERNET		55
8.1	Angling organizations	55
8.2	Animal rights organizations	55
8.3	Animal welfare organizations	56
8.4	Aquaculture organizations	57
8.5	Fisheries organizations	57
8.6	Governmental and intergovernmental organizations	58
8.7	Scientific organizations	59
8.8	Codes and guidelines	60
8.9	Others	61
APPENDIX		62
BIBLIOGRAPHY		62
1.	References for “Animal welfare”	62
2.	References for “Painperception of fish”	65

2.1	Painperception	65
2.2	Anatomical, pharmacological and neurophysiological studies pertaining to pain, fear and stress in fish	66
2.3	Cognitive ability, fear and awareness in fish	68
3.	References for “Commercial fisheries”	70
3.1	Electric fishing	70
3.2	Bycatch	72
3.3	Fishing gear escape	72
3.4	Physiological effects	72
3.5	Mortality	72
3.6	Other	73
4.	References for “Recreational fisheries”	73
4.1	Catch and release	73
4.2	Physiological effects/stress	75
4.3	Mortality	76
4.4	Retention in keepnets/confinement	78
4.5	Other	79
5.	References for “Aquaculture”	79
5.1	Handling	79
5.2	Slaughter	80
5.3	Stocking density	82
5.4	Physiological effects/stress	82
5.5	Tagging	85
5.6	Transportation	85
5.7	Social interaction/behaviour	86
5.8	Food	86
5.9	Light regime	87
5.10	Health	87
5.11	Other	88

PREFACE

Attention about animal welfare has increased in recent years, and the welfare of fishes has also become a subject of attention (Kestin, 1993; FAWC, 1996; Lymbery, 2002; Erickson, 2003; FSBI, 2002). Furthermore, the handling of fishes and fish stocks is increasingly subject to legislation and regulation. The European Inland Fisheries Advisory Commission (EIFAC) felt there was a need to exchange information on the practice of handling fishes in fisheries and aquaculture and to identify current legislation and regulations in EIFAC member countries. The twenty-second session¹ of EIFAC in Windermere, United Kingdom, in 2002, decided to establish an ad hoc Working Party on Handling of Fishes in Fisheries and Aquaculture, with the following terms of reference:

- compilation of the basic scientific knowledge on the effects of handling of fishes in fisheries and aquaculture (fishery methods, harvest, holding and rearing, use of anaesthetics, slaughter);
- compilation of the current practice of handling of fishes in fisheries and aquaculture, including training programmes, and collation of existing codes of practice and guidance in EIFAC member countries and relevant information from other countries;
- listing of the current and proposed legislation and regulation on handling in fisheries and aquaculture and in experimental settings in EIFAC member countries (including EU regulations);
- identification of the relevant Web sites and other sources for information on welfare of fishes and handling of fishes and other relevant topics;
- distribution of information about the discussion on fish welfare in EIFAC member countries.

Part one of this report presents the result of the EIFAC Ad Hoc Working Party, in particular the Workshop held in Utrecht in March 2004. Part two gives an overview of current knowledge available in scientific literature, grey literature and on the World Wide Web, on several issues relating to fish welfare, and provides comprehensive reference lists and relevant Web sites dealing with aspects of fish welfare.

Nieuwegein, February 2006

Lex Raat

Convener, EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture

¹ FAO 2002. Report of the twenty-second session of the European Inland Fisheries Advisory Commission. Windermere, United Kingdom, 12–19 June 2002. FAO Fisheries Report. No. 681. Rome, FAO. 42p.
<ftp://ftp.fao.org/docrep/fao/005/y3930e/y3930e00.pdf>

PART ONE

REPORT OF THE EIFAC WORKSHOP ON HANDLING OF FISHES IN FISHERIES AND AQUACULTURE, 24–26 MARCH 2004

INTRODUCTION

1. The handling of fishes is increasingly subject to legislation and regulation. In 2002 the twenty-second session of European Inland Fisheries Advisory Commission of FAO (EIFAC) in Windermere, United Kingdom, realized the need to identify current legislation and regulations in the EIFAC member countries. Also, a need was felt to exchange information on the actual practise of handling fishes in fisheries and aquaculture, e.g. catching and handling, husbandry conditions, transport and slaughter of fishes. The Session decided to establish an EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture. The ad hoc Working Party is a forum to exchange information about the handling of fishes for people working in fisheries and aquaculture. The ad hoc Working Party was convened by Mr A.J.P. Raat, Organization for the Improvement of Inland Fisheries (OVB), Netherlands.

2. In March 2004 the convener of the ad hoc Working Party, Mr A.J.P. Raat, organized an international EIFAC Workshop on Handling of Fishes in Fisheries and Aquaculture in Utrecht (Netherlands). The workshop was attended by 16 persons from six European countries. The list of participants can be found in the Annex to the workshop report.

AIMS OF THE EIFAC WORKSHOP

3. The meeting had the following aims:

- Compilation of the current practice of handling of fishes in fisheries and aquaculture, including training programmes, and collation of existing codes of practice and guidance in EIFAC member countries and relevant information from other countries;
- Compilation of the current and proposed legislation and regulation on handling in fisheries and aquaculture and in experimental settings in EIFAC member countries (including EU regulations);
- Distribution of information about the discussion on welfare of fishes in EIFAC member countries.

OUTPUT OF THE EIFAC WORKSHOP

4. The EIFAC Workshop covered a wide range of interests in the handling of fishes with emphasis on freshwater fishes. The Workshop explored how new knowledge, guidelines and codes of practice, regulations and legislation might influence inland fisheries, fisheries management, inland fisheries research and aquaculture. The participants gave short presentations about their professional experience of handling fishes and the current state of regulation and legislation in their countries. Trends and developments were also identified. The general outcome is an improved understanding and awareness of the effects of handling of fishes in inland fisheries, fisheries research and aquaculture. Furthermore positive suggestions for better management and a direction set for future work were supplied by the participants.

5. This report gives a summary of the EIFAC Workshop Handling of Fishes. The conclusions and recommendations of the participants of the Workshop were presented to the twenty-third session of EIFAC in Wierzba, Poland, June 2004, for future actions.

6. Furthermore the participants of the Workshop amended and updated the draft document *Handling of fishes* that was prepared for discussion by the convener of the Working Party. The revised document *Handling of fishes in commercial inland fisheries, recreational fisheries, aquaculture and fisheries research. An overview*, consolidates the available information on handling of fishes and supplies basic information for a future position statement of EIFAC. The document was presented to the twenty-third session of EIFAC in 2004 for information and publication.

GENERAL CONSIDERATIONS

7. The participants of the Workshop recommend that in fisheries, fisheries research and aquaculture one should be aware that fishes are sophisticated animals, which makes a fish subject to careful handling.

Box 1: Valuation of fishes

- (Human) use values (fishes object). These values embed an anthropocentric, moral position and perpetuate a relationship of domination between people and nature:
 - fishes as objects for fisheries;
 - fishes as objects in laboratory and field experiments.
- (Human) use values (market value). These assigned values are market prices or their equivalent (observable terms of exchange between goods and services):
 - fishes in commercial (recreational) fisheries;
 - fishes in aquaculture;
 - fishes for consumption;
 - trade in ornamental fishes.
- (Fish) inherent values (fishes subject). These values reside in the object – the fish – itself.
 - fishes are unique individual creatures;
 - fishes cannot be made by humans;
 - fishes can live, reproduce and grow;
 - fishes in the aquatic resource.

8. The participants to the Workshop noted that animal welfare, including the welfare of fishes, is subject to debate in our society. The attitude of the public towards fishes has been changed in the EIFAC member countries during the past fifty years. The valuation of the fishes has been broadened. In our present society fishes represent both human use values as well as inherent values. The human use values are complex and difficult to formulate in generally acceptable and operational terms. The inherent values are by definition intangible. This has brought the use and handling of fishes not only in the realm of fish biologists, fisheries workers and aquaculturists, but also in the public debate. This is reflected in the Animal Welfare legislation in various EIFAC member countries. Handling of fishes is a human action which needs a responsible approach by those who are in charge of the handling and use of the fishes.

9. The present public debate in many EIFAC member countries deals with the acceptability of (recreational) fisheries or special forms of these fisheries, such as catch-and-release fishing (forbidden in some parts of Germany), use of fishing methods (e.g. electric fishing), holding of fishes (e.g. in cages, keepnets) or live fish bait fishing (forbidden in The Netherlands).

10. In aquaculture and commercial fisheries the perceptions of consumers and the public regarding the conditions in which the fishes are raised and captured, play a role in the acceptance of the products. Here the experience with human welfare and the perceptions of the welfare of mammals and birds is often extrapolated to fishes. However, this does not exclude a critical analysis of the handling procedures in practice and, when needed, an improvement of the applied methods and technology, by those responsible for the handling of the animals.

11. The participants of the Workshop observed that although there is a critical public attitude to the handling of fishes, the behaviour of the consumer until now seems to be mainly directed by the price and the perceived safety and quality of the products.

12. The participants of the Workshop noted that there is an important role for the fisheries and aquaculture sectors to respond to the awareness of the public concern about the use and handling of the fishes.

13. In many countries the use of animals, including fishes, for experimental purposes, seems to be more strictly regulated than their use in other human activities. The participants of the Workshop

observed that this is an additional reason for the fisheries and aquaculture sectors to take their responsibility for improving the way that fishes are used, kept, sampled, captured, handled and killed.

14. The participants of the Workshop noted that aquaculture and fisheries organizations are aware of the need to improve the conditions from the perspective of the welfare of the fishes. There are many codes of conduct and guidelines that take the welfare of the fishes into consideration. However, further efforts are needed to improve the translation of these codes of conduct and guidelines to the way in which fish are handled in practice. Training and education programmes can improve the awareness among individual persons. The certification of actions (to “clarify the welfare criteria”) by the sector itself is also a good marketing instrument, which endorses the acceptability of fisheries and aquaculture products.

15. The participants of the Workshop noted that part of the public perception of fish welfare issues in fisheries and aquaculture is based on anthropocentric assumptions. Communication with stakeholders such as consumers, trade, non-governmental organization (NGOs), fishermen, angling organizations, aquaculturists and government, can lead to a better understanding of the way these issues are taken up by the fisheries and aquaculture sectors.

WELFARE AND FISHES

16. The participants of the Workshop noted the difficulty of finding a comprehensive, generally acceptable definition of fish welfare. However, in the EIFAC member countries, scientific considerations and ethical standards mandate a respect for all life forms. Thus the framework based on the “five freedoms” (defined in the United Kingdom Farm Animal Welfare Council), that recognises five domains in which welfare may be compromised, is relevant for the identification of fish welfare criteria in fisheries, fisheries research and aquaculture.

Box 2: Five “freedoms” defined in the United Kingdom Farm Animal Welfare Council.

Domain 1. Water and food deprivation, malnutrition. Fishes should have ready access to (fresh) water and an appropriate diet in sufficient quantities and with a composition that maintains full health and vigour.

Domain 2. Environmental challenge. Fishes should have a suitable environment, including shelter and a comfortable resting area, whether in their natural environment or in holding conditions.

Domain 3. Disease, injury and functional impairment. Disease should be prevented or rapidly diagnosed and treated.

Domain 4. Behavioural/interactive restriction. Fishes should have sufficient space, proper facilities and where appropriate, the company of the fishes own kind.

Domain 5. Mental and physical suffering. Conditions that produce potentially unacceptable levels of anxiety, fear, distress, boredom, sickness, pain, thirst, hunger and so on should be minimised.

17. The participants of the Workshop emphasize that there is a lack of scientifically sound criteria to assess fish welfare. In addition, they noted that not all criteria are applicable in field circumstances and that many of the criteria are restricted for use under controlled conditions e.g. laboratory experiments or intensive aquaculture. Moreover, criteria with respect to welfare of fish are:

- species specific (e.g. eel and catfish);
- individual/ genetically specific (e.g. catfish);
- gender specific (e.g. catfish and carp);
- depending on size and age;
- depending on the environment.

*Box 3: Criteria for fish welfare.***Behavioural:**

- swimming activity;
- escape;
- elevated gill ventilation;
- stereotype behaviour (aquaculture only);
- coping styles (agonistic behaviour: aquaculture only).

Physiological:

- stress physiology: blood: neuroendocrine responses, muscle outer appearance;
- brain activity for assessment of stunning/killing methods;
- heart rate;
- *post mortem* biochemistry of muscle tissue.

Morphological:

- injuries;
- fin damage;
- body condition indices.

Performance-based indicators:

- mortality;
- feed conversion efficiency (aquaculture only);
- flesh quality;
- growth rate;
- reproductive success (fisheries only; long-term exposure to a stressor);
- resistance against diseases, disease state (aquaculture only);
- pathological changes (aquaculture only).

Environmental:

- water quality;
- water availability;
- endemic diseases/ parasites.

PAIN AND SUFFERING OF FISHES

18. The participants of the Workshop discussed the similarities and differences of fishes and mammals with regard to the subjective experience of pain and suffering. The argument was put forward that the cortical regions, that are essential for human pain experience, are not present in the brains of the fishes, and therefore the “human” experience of pain is a neurological impossibility. However, following this reasoning it can be argued that birds are blind, because they do not possess a visual cortex (FSBI, 2002). In the case of the birds this task (processing of visual stimuli) is performed by the mid brain optic tectum. It is therefore not impossible that parts of the brain other than the cerebral cortex have evolved the capacity for generating negative emotional states/suffering in non-mammalian vertebrates, including fishes.

19. Recent studies, which have caused controversy in the scientific community, were presented to the participants at the Workshop suggesting that fishes are capable of perceiving pain. The results of these studies by a scientific group from the Roslin Institute in Edinburgh, albeit refuted by others, are based on similarities between fishes and higher vertebrates in their basic structure of neurones and their neuronal biochemistry. There are also similarities in stress responses and in behavioural and decision-making responses to tissue damaging, or potentially tissue damaging, incidents. Recent anatomical and electrophysiological studies have further demonstrated that fishes are capable of nociception: the simple detection of a noxious, potentially painful stimulus and the reflex response to this. The pain-related behaviour which was observed however, is not a simple reflex. Thus, the presence of a neocortex as a brain structure, seems not necessarily required for the perception of pain by fishes.

20. The discussion on pain and suffering of fishes during the Workshop did not result in a shared opinion among the participants. However, independent of the question on pain and suffering of fishes, the participants agreed that handling of fishes has to be done in a responsible way.

RESPONSIBLE HANDLING OF FISHES

21. The participants of the Workshop noted that fishes show reactions to injurious stimuli. These stimuli result in behaviour leading to physiological stress responses. Depending on the duration and intensity of the exposure to the stressor, the stress response varies from acute to chronic.

22. The participants of the Workshop agreed that handling causing injuries and damage to the fishes should be avoided in fisheries, aquaculture and fisheries research. This means that careful handling of the fishes implies prevention of damage or unnecessary injuries, prevention of unnecessary stress and prevention of unnecessary prolongation of the period before the killing takes place.

23. The participants of the Workshop noted that those responsible for handling of fishes in fisheries, aquaculture and fisheries research should have basic knowledge of the effects of handling on fishes and have justifiable objectives (“reasonable purpose”), within the context of the situation, and furthermore take into account the rules and regulations with regard to these activities.

HANDLING OF FISHES: FISHERIES RESEARCH

24. The participants of the Workshop noted that in many EIFAC member countries there is legislation dealing with animal experimentation, including those with fishes. Most countries do not have specific regulations or legislation for fishes in experimental settings.

25. Special fields of interest relating to fish welfare that were identified during the Workshop were:

- Tagging and marking. It was noted, that recently a relevant report of concerted action within the EU framework FAIR Tagging methods for stock assessment and research in fisheries was published (Davenport *et al.*, 2002).
- Stunning and consciousness of fish.
 - It was noted that most anaesthetics and related chemicals used for stunning the fishes do result in enhanced brain activity and increase in heart rate.
 - It was further noted that in most EIFAC member countries anaesthetics for fisheries use are not certified (in several countries only MS 222 is certified).
 - It was concluded that there is a need for better understanding and certification of anaesthetics for use in fish handling and fish research (e.g. clove oil, Benzocaine, 2-Phenoxyethanol).
- Implantation techniques:
 - It was noted that the practice of implantation is subject to wide variation within the EIFAC member countries with regard to legislation, experience, procedures and protocols.
 - It was concluded that international co-operation in training for the use of invasive methods and techniques in EIFAC member countries is advisable. EIFAC should play a co-ordinating and stimulating role for initiatives for concerted action.
- Parasite infections.
- Sampling of fishes:
 - It was noted that there is a need for more information on the aspects of fish welfare affected by fisheries methods (e.g. electric fishing, gill netting).
 - It was noted that standardization of fishing methods should take into account the available information on fish welfare.
 - It was advised that in monitoring and sampling programmes more attention should be paid to alternative methods, which leave the fishes in their natural environment (e.g. acoustic surveys).

HANDLING OF FISHES IN PRACTICE: FISHES IN AQUACULTURE

26. The participants of the Workshop identified several topics relevant from the perspective of fish welfare:

- Rearing conditions:
 - water quality;
 - stocking density/ Loading rate;
 - light regime;
 - feeding regime;
 - design of a tank.
- Transportation.
- Fasting, sorting and crowding, transport.
- Harvest: the process of catching and handling of farmed fish. It was noted that more research effort must be invested in the technical aspects of fish capture in aquaculture situations.
- Slaughter: the process of stunning and killing. It was noted that more research effort must be invested in the technical aspects of fish killing, taking into account industrial constraints.
- Stunning/killing and flesh quality. It was noted that an efficient and rapid stunning/killing of the fishes usually leads to a better quality of the flesh. Prolonged storage of the fishes prior to stunning or killing, may have a negative effect on the flesh quality.
- Problems with parasites, predators and birds.
- Disease treatment and prevention of diseases.
- Technical aspects: optimise conditions.

27. The participants of the Workshop noted that the Council of Europe standing committee of the European Convention for the Protection of Animals kept for Farming Purposes is presently working on farmed fish, including a *Draft recommendation concerning farmed fish* (which was adopted in December 2005 and published by the Council of Europe²). It was recommended by the participants that EIFAC should have a liaison with the Council of Europe regarding their activities in the field of aquaculture.

28. It was further noted that in the European Union, retailers, animal welfare associations and governments are increasingly demanding humane methods to convert live farmed fishes into food. The European Food Safety Authority (EFSA) is currently reviewing current stunning and killing methods for all animals (including fishes). Various European research institutes are taking part in the expert working group. It was recommended by the participants of the Workshop that EIFAC should have a liaison with these activities.

HANDLING OF FISHES IN PRACTICE: ORNAMENTAL FISHES

29. The participants of the Workshop noted that about 80 percent of ornamental fishes are captured from natural waters and are not cultured. The following topics are relevant from the perspective of fish welfare:

- capture;
- transportation;
- holding conditions.

30. Although mortality due to the trade in ornamental fishes can be substantial on global, European, national and local scale, actions by EIFAC have no priority when considered from the terms of reference of this advisory body.

HANDLING OF FISHES IN PRACTICE: FISHERIES MANAGEMENT

31. Welfare of fishes deals not only with individual fishes but also with wild fish populations. The participants of the Workshop identified several topics relevant from the perspective of fish welfare:

² Recommendation Concerning Farmed Fish, adopted by the Standing Committee on 5 December 2005. Standing Committee of the European Convention for the Protection of Animals kept for Farming Purposes (T-AP). http://www.coe.int/t/e/legal_affairs/legal_co-operation/biological_safety,_use_of_animals/Farming/Rec%20fish%20E.asp

- catch and release fisheries;
- put and take fisheries;
- stocking practices;
- use of live bait fish;
- keepnets;
- harvest: the process of catching, handling, slaughter for wild fish;
- protection from predators;
- disease and parasites;
- eradicating fish stocks (e.g. rotenone);
- biomanipulation fisheries;
- effect of hydro power plants;
- effects of damming on (migratory) species;
- enforcement of regulations.

32. The participants of the Workshop recommended that EIFAC should take action to extend knowledge and guidance about the fish welfare aspect of fisheries management.

RELEVANT INFORMATION

33. The participants of the Workshop noted that relevant information on handling of fishes is summarized in the following documents:

- *Handling of fishes*, discussion paper prepared by A.J.P. Raat and P. van de Nieuwegiessen for the EIFAC ad hoc Working Party on Handling of Fishes in Fisheries and Aquaculture.
- *Guidelines for the use of fishes in research* of the American Fisheries Society (Nickum *et al.*, 2004).
- *Fish welfare. Briefing paper 2* of the Fisheries Society of the British Isles (FSBI, 2002).
- *Information resources on fish welfare 1970-2003* of the Animal Welfare Information centre of the US department of Agriculture (Erickson, 2003).

34. All persons professionally engaged with handling of fishes should be aware of the existence of these papers and should apply the information and guidelines in their handling of the fishes.

REFERENCES

Davenport, J., Baras, E., Fabi, G. & Jonsson, G., 2002. Chapters 6 (Legislation) & 7 (Fish welfare and health in relation to tagging) In: Thorsteinsson V. (coordinator) 2002. *Tagging Methods for Stock Assessment and Research in Fisheries. Report of Concerted Action FAIR CT.96.1394 (CATAG)*. Reykjavik. *Marine Research Institute Technical Report* No. 79. pp 183.

Erickson, H.S., 2003. *Information Resources on Fish Welfare: 1970-2003*. AWIC Resource Series No. 20. U.S. Dept. of Agriculture, National Agricultural Library, Animal Welfare Information Center. Beltsville, MD. 436 pp.
<http://www.nal.usda.gov/awic/pubs/Fishwelfare/fishwelfare.htm>

FSBI, 2002. *Fish Welfare*. Briefing Paper 2. Fisheries Society of the British Isles, Granta Information Systems, 82A High Street Sawston Cambridge. 25 pp.
<http://www.leicester.ac.uk/biology/fsbi/welfare.pdf>

Nickum, J.G., Bart, H.L., Bowser, P.R., Greer, I.E., Hubbs, C., Jenkins, J.A., MacMillan, J.R., Rachlin, J.W., Rose, J.D., Sorensen, P.W. & Tomasso, J.R. 2004. *Guidelines for the use of fishes in research*. American Fisheries Society; American Institute of fisheries research biologists; American Society of Ichthyologists and Herpetologists.
http://www.fisheries.org/html/Public_Affairs/Sound_Science/Guidelines2004.shtml

ANNEX

List of participants

Anduleit, Martine

VWA
 Postbus 19506
 2500 CM Den Haag
 The Netherlands
 Tel.: 0031-704484493
 Fax.: 0031-704484430
 E-mail: martina.anduleit@vwa.nl

Jepsen, Niels

DFU, Department of Inland Fisheries
 Vejlsøvej 39
 DK-8600 Silkeborg
 Denmark
 Tel.: 0045-89213150
 Fax.: 0045-89213150
 E-mail: nj@dfu.min.dk

Kane, Frank

Aquaculture Section, ACMS Marine Institute
 Galway Technology Park, Parkmore
 Galway
 Ireland
 Tel.: 0035-39841107
 E-mail: frank.kane@marine.ie

Klein Breteler, Jan

Organisatie ter Verbetering van de
 Binnenvisserij
 Postbus 433
 3434 AK Nieuwegein
 The Netherlands
 Tel.: 0031-306058411
 Fax.: 0031-306039874
 E-mail: kleinb@ovb.nl

Lambooij, Bert

Animal Sciences Group of the Wageningen-
 UR, Division Nutrition
 and Food
 Postbus 65
 8200 AB Lelystad
 The Netherlands
 Tel.: 0031-320238970
 Fax.: 0031-320237320
 E-mail: bert.lambooij@wur.nl

Mueller, Rudolf

(Chair of EIFAC)
 EAWAG, Department of Limnology
 Fisheries Sciences
 Limnological Research Center
 CH-6047 Kastanienbaum
 Switzerland
 Tel.: 0041-41-349 21 30
 Fax.: 0041-41-349 21 62
 E-mail: rudolf.mueller@eawag.ch

Nieuwegiessen, Pascal van de

WUR
 Oude Kerkstraat 13a
 4271 BB Dussen
 The Netherlands
 Tel.: 0031-416392254
 E-mail: Pascal.vandeNieuwegiessen@wur.nl

O'Donohoe, Pauline

Aquaculture Section, ACMS Marine Institute
 Galway Technology Park, Parkmore
 Galway
 Ireland
 Tel.: 0035-39841107
 E-mail: pauline.odonohoe@marine.ie

Pietroock, Michael

Institute for Inland Fisheries
 Jägerhof am Sacrower See
 D-14476 Groß Glienicke
 Germany
 Tel.: 0049-3320140613
 Fax.: 0049-3320140640
 E-mail: michael.pietroock@ifb-potsdam.de

Raat, Lex (A.J.P.)

(Convenor of Working Party)
 Organisatie ter Verbetering van de
 Binnenvisserij
 Postbus 433
 3400 AK Nieuwegein
 The Netherlands
 Tel.: 0031-302933529
 lexraat@planet.nl

Roesch, Roland

Fischereiforschungsstelle Baden-
Württemberg
Untere Seestr. 81
D-88085 Langenargen
Germany
Tel.: 0049-7543-930816
Fax.: 0049-7543-930820
E-mail: roland.roesch@lvvg.bwl.de

Schultze-Wiehenbrauck, Hardy

Ministerium Für Umwelt und Naturschutz,
Landwirtschaft und Verbraucherschutz
Nordrhein-Westfalen
D-40190 Düsseldorf
Deutschland
Tel.: 0049-2114566245
Fax.: 0049-2114566415
E-mail: schulze-w@munlv.nrw.de

Schrama, Johan

WUR
Postbus 338
6700 AH Wageningen

Shields, Brian

Environmental Agency
Knutsford Road
Warrington WA4 1HG
United Kingdom
Tel.: 0044-1925653999
Fax.: 0044-1925415961
E-mail: brian.shields@environment-
agency.gov.uk

Tredwin, Neil

Cefas Laboratory
The Nothe, Barrack Road
Weymouth, United Kingdom DT4 8UB
United Kingdom
Tel.: 0041-01305 206676
E-mail: n.j.tredwin@cefas.co.uk

Vis, Hans van de

Netherland Institute for Fisheries Research
(RIVO)
Postbus 68
1970 AB Ijmuiden
The Netherlands
Tel.: 0031-255564613
Fax.: 0031-255564644
E-mail: hans.vandervis@wur.nl

PART TWO

HANDLING OF FISHES IN COMMERCIAL INLAND FISHERIES, RECREATIONAL FISHERIES, AQUACULTURE AND FISHERIES RESEARCH. AN OVERVIEW

CHAPTER 1 FISH WELFARE

1.1 Defining animal welfare

In recent years, animal welfare has been the subject of intense debate and public concern. However, to discuss animal welfare objectively, a definition of animal welfare is required. Producing such a definition has not been easy, particularly because the concept of animal welfare is complex and the phrase has been used in different contexts. Most definitions can be divided into three broad categories (Duncan and Fraser, 1997; Fraser *et al.*, 1997).

- *Feelings-based definitions:* These definitions are normally based on subjective mental states. The requirement for good welfare is that a certain animal should feel mentally well, it should be free of pain and fear and it should have access to positive experiences, such as social contacts. Naturally, such definitions are only valid if the animal of concern is able to experience these subjective mental states. Furthermore, the use of these definitions depends on understanding of these subjective mental states.
- *Function-based definitions:* These definitions are based on the physical state of an animal. Here, for an animal to experience good welfare it should be healthy and its biological systems should be functioning appropriately. However, these definitions are based on the assumption that animals in good physical health are also in a good mental state.
- *Nature-based definitions:* These definitions are based on the biological functioning of an animal. Good welfare requires that an animal is allowed to perform its natural behaviour and live a natural life. These definitions assume that animals are suffering if they cannot express their full behavioural repertoire. However, a part of their natural behaviour is based on escaping from adverse situations, such as escaping from a predator, and it may be hard to argue that an animal is suffering if this behaviour cannot be expressed. Furthermore, the concept provides little guidance on many important animal welfare issues, such as the use of analgesia, euthanasia and medication.

Unfortunately, none of the concepts of animal welfare offer a comprehensive view on the major ethical concerns over the quality of life of animals; an ideal definition would combine each of the three stated concepts.

1.2 Criteria for fish welfare

1.2.1 Five freedoms

To overcome the problems with producing a comprehensive definition of fish welfare, criteria were defined that should be fulfilled for an animal to experience good welfare. These “five freedoms” were elaborated by the Brambell Committee of the United Kingdom Parliament in 1965 and updated by the Farm Animal Welfare Council (1992):

1. freedom from hunger, thirst and malnutrition;
2. freedom from physical and physiological discomfort;
3. freedom from injury, disease and functional impairment;
4. freedom to express normal behaviour and social interactions;
5. freedom from fear and chronic stress.

The Fisheries Society of the British Isles (FSBI, 2002) reworked these “five freedoms” into a form more appropriate for fishes. These freedoms can be applied to all fish, although the details mentioned under each freedom depend on the species concerned, age, gender and reproductive condition.

1. *Thirst, hunger and malnutrition.*
 - Fish should have access to an adequate, nutritionally complete diet, taking into account the fact that fish vary greatly in their natural diet, that they do not need to maintain a constant body temperature and that in many cases they show marked differences in requirements with season and life history stage.
 - Fish should be presented with food in a manner that is appropriate to the natural feeding behaviour of the species concerned (e.g. pellets of the correct size) and that avoids undue competition.
2. *Physical and physiological discomfort.*
 - Fish should be kept under water quality, flow rates and temperature appropriate for the species concerned.
 - Seasonal and daily patterns of light intensity should be appropriate for the species concerned.
 - Fish should be provided with cover and shelter.
3. *Injury, disease and functional impairment.*
 - Disease should be prevented or rapidly diagnosed and treated when possible.
4. *Natural behaviour/social interactions.*
 - Fish should have sufficient space to allow a degree of freedom of movement, but the definition of “sufficient” will be species-specific.
 - For shoaling species, the company of their own kind is important for welfare, but for territorial species, this may not be true.
 - A degree of environmental complexity may be important, depending on the species concerned.
5. *Fear and chronic stress.*
 - Conditions that potentially produce unacceptable levels of anxiety, fear, distress, boredom, sickness, pain, thirst, hunger and so on should be minimised in fish, as in other vertebrates.

The fifth freedom in particular may be subject of debate, because the ability of fish to experience pain, fear and suffering is a highly contentious issue. Arguments have been made both to support and refute the claim that fish are capable of experiencing pain and suffering. Rose (2002) argues against the ability of fish to experience pain and suffering, based on the distinction between reaction to injury (nociception) and the psychological experience of pain (see section 1.2.2).

To comprehend this distinction it is important to understand the main components of the pain perception mechanism. Components of this mechanism in mammals are shown in Figure 1. In this figure the first component of experiencing pain is nociception. This term refers to the detection of noxious stimuli by the nervous system. Nociception is controlled by the spinal cord and brainstem. The two types of nociceptors are: A-delta fibres with thinly myelinated axons; and C-fibres with slowly conducting unmyelinated axons. The occurrence of these fibres in fish varies according to species groups:

- Elasmobranchs: Although nociceptors were found in several studies of elasmobranchs, the studies failed to detect pain in the studied fish species (Sneddon, Braithwaite and Gentle, 2003).
- Bony Fish: C- and A-delta fibres present in the peripheral nerves. Very few C-fibres, only around 4 percent of afferents are C-fibres. Primarily A-fibres.
- Hagfish: Nociceptors present, primarily C-fibres.

The occurrence of these fibres in bony fish and hagfish implies that these species groups must be able to react to injury. Consequently, without doubt both fishes and humans are capable of responding to noxious stimuli. For instance, a fish that has been hooked is responding to a stimulus. Likewise, if someone burns themselves, he or she will very quickly respond to the stimulus, but this response occurs before one feels any pain.

1.2.2 Discussion on perception of pain by fishes

Rose (2002) stated that the difference in the perception of pain and fear of fishes and humans results from the different structure of the brains. The human brain has a massively developed cerebral cortex. Pain and fear in humans results from the stimulation of several regions of this cerebral cortex

(Bermond, 1997; Rose, 2002). The tiny cerebral cortex of fish brains lack these regions (Figure 2). Reaction to injury, therefore, does not necessarily mean that psychological pain is experienced.

Recent studies, however, have caused controversy in the scientific community by suggesting that fishes are capable of perceiving pain (Braithwaite, 2004). The results of these studies are based on similarities between fishes and higher vertebrates in their basic structure of neurones and their neuronal biochemistry. There are also similarities in stress responses and in behavioural and decision-making responses to tissue damaging, or potentially tissue damaging, incidents. Anatomical and electrophysiological studies further demonstrated that fishes are capable of nociception: the simple detection of a noxious, potentially painful stimulus and the reflex response to this. The pain-related behaviour that was observed, however, was not a simple reflex. Thus, the presence of a neocortex as a brain structure, may not necessarily be a prerequisite for the perception of pain by fishes (Sneddon, 2003; Braithwaite and Huntingford, 2004; Chandroo *et al.*, 2004).

Modern brain imaging technology has made it possible to observe directly and quantify neural events that relate systematically with the subjective experience of pain (Broom, 2001). Although some of these methods (brain imaging technology), in principle, could be applied to fishes, according to Nickum *et al.* (2004) the effort would be fruitless due to fundamental differences between fish and human brain structure.

Sneddon (2003) stated: In a review concluding that fish are not capable of pain perception, Rose (2002) defined pain in terms of brain structure and stated that an animal must have the necessary brain structure, a neocortex, to perceive pain. With this definition only humans and primates are capable of experiencing pain and Rose (2002) ignored the bird and amphibian literature that has proven these animals are capable of pain and yet they do not have a neocortex.

The argument proposed by Rose (2002) is based on the observation that the cortical regions essential for human pain experience, are not present in the brains of fishes, and therefore the “human” experience of pain is a neurological impossibility. However, following this reasoning it can be argued that birds are blind, because they do not possess a visual cortex (FSBI, 2002). In the case of birds this task (processing of visual stimuli) is performed by the mid brain optic tectum. Other examples exist of similar functions being carried out by different parts of the brain in different animal groups (FSBI, 2002). It is therefore possible that parts of the brain other than the cerebral cortex have evolved the capacity for generating negative emotional states or suffering in non-mammalian vertebrates, including fish.

Statements and conclusions made in a review by Chandroo *et al.* (2004) of the perception of sentience, pain, fear and stress in fishes are summarized below:

Fish behaviour and internal representations:

- There is evidence that fish can remember, recognize and interpret the fighting skills of future opponents to assess their own probability of winning a future conflict, as well as signal their motivational state of familiar conspecifics.
- This forms a preliminary basis for suggesting that primary consciousness and cognition exists in fishes.

Motivational affective:

- Despite differences in ontogeny, there is a growing body of anatomical, behavioural and pharmacological evidence to suggest that the brain structures and neural systems associated with motivational affective states in tetrapods are present in fishes.
- Therefore, it is reasonable to suggest that motivational affective states could have developed at some level of fish evolution.




Figure 1. The main components of the pain perception mechanisms in mammals (FSBI, 2002).




Figure 2. Comparison of the structure of the human brain (A) and the trout brain (B). (C) Brain of a 30 cm rainbow trout (*Oncorhynchus mykiss*) shown at the same scale as the human brain diagram (Rose, 2002).

Pain and the telencephalon:

- That pain in fish may be experienced in ways similar to tetrapods.
- The cytoarchitecture of the fish brain is one that permits integration of differentiated pallial and sub-pallial regions, it is therefore reasonable to suggest that modern hypothesis pertaining to the neural basis of consciousness may apply to fish.
- Fish have the ability to construct cognitive representations of noxious events and are able to form expectancies of the noxious stimuli as a result (also suggested by Overmier and Hollis, 1990).
- It is conceivable that nociception (and other senses) in fish is consciously experienced, and is responsible for motivated behaviour.

Fear and anxiety:

- The cognitive neurophysiological and behavioural features of fear responses of fish suggest that they have some capacity to consciously experience fear.

Psychological aspects of stress:

- The neurobehavioral similarities between fish and other vertebrates with respect to social stress further supports the suggestion that fish have some capacity to experience psychological stress.

In another paper on fear reactions of the domesticated rainbow trout, Yue *et al.* (2004) concluded that: “This evidence suggests that trout can experience fear and that they can learn to avoid frightening stimuli. It implies that they are sentient animals, more complex than previously thought”.

Arguments presented in support of the belief that fishes are capable of experiencing pain can be divided into three forms.

- *Arguments stating that behavioural reactivity to presumed noxious stimuli are evidence of pain perception.* A recent study by Sneddon *et al.* (2003) looked at the behavioural response of rainbow trout to administration of noxious substances (bee venom and acetic acid) to the lips. Both the physiology and behaviour of the fish were affected. A significant increase in opercular beat rate and the time taken to resume feeding were found. Furthermore, fish injected with venom and acid showed anomalous behaviour. Fish performed “rocking” where they moved from side to side balancing on either pectoral fin while resting on the gravel. Fish injected with acid also rubbed their lips into the gravel and against the tank walls. The affected physiology and behaviour of the fish over a prolonged period of time may suggest discomfort and pain. The authors concluded that they had proved that fish experience pain. This conclusion, however, implicitly starts from the assumption that the “human” brain is not necessarily the requirement for the perception of pain. Nickum *et al.* (2004) and Rose (2003) argue that until conscious reaction to injurious stimuli can be demonstrated, the behaviour demonstrated by the fish in this study is not proof of nociceptive behaviour.
- *Arguments stating that learning by fishes to avoid noxious stimuli or to develop Pavlovian conditioned responses to these stimuli are evidence of pain experience.* Several studies have been performed on common carp (*Cyprinus carpio*; Beukema, 1970; Raat, 1985) to support this argument. Carp, initially hooked once and then released, learned to associate the bait with the hooking experience and subsequently avoided the bait. Up to three years later, the previously hooked fish were more difficult to catch than naïve fish. Furthermore, the remaining fish also became progressively more difficult to catch as the others were caught, implying that the unhooked fish were able to learn from the hooked fish that the experience was aversive. However, this behaviour, also known as implicit learning, is a virtually universal capability of vertebrate and invertebrate animals, including those lacking brains (Macphail, 1998, cit. in Rose, 2002).
- *Arguments stating that the same or equivalent neural structures and neurochemicals mediating pain in humans are present in fish.* Ehrensing *et al.* (1982) performed a study to support this argument. The response (agitated swimming) of goldfish (*Carassius auratus*) to an electric shock was studied. Morphine was used to induce analgesia, and naloxone was used to reverse this analgesia. Increases in the concentration of morphine injected intracranially increased the intensity of shock needed to stimulate an agitated swimming response in a dose dependent way,

and naloxone reversed this effect. The compounds thus behaved in a manner analogous to the way they would behave in a rat subjected to a similar experimental protocol. This implies that fish react to analgesics similarly, and may thus feel pain similarly. Opiates, such as morphine, are known to produce their anti-nociceptive effects by acting at subcortical brainstem and spinal cord sites (Prices, 1999, cit. in Rose, 2002). Thus, opiate effects on nociceptive processing are not evidence of a capacity for consciously perceived pain (Rose, 2002).

Braithwaite and Huntingford (2004) summarized the arguments for and against the ability of fish to perceive pain. They agreed with Rose (2002) that fish are unlikely to perceive pain in the same way that humans do. However, they believed that currently available evidence indicates that fish have the capacity for pain perception and suffering. As such, it would seem timely to reflect on the implications of fish pain and suffering, and to consider what steps can be taken to ensure the welfare of the fish that we exploit.

The arguments to support and refute the claim that fish are capable of experiencing pain and suffering currently largely depend on the definition of pain that is used. The definition of pain used by Bermond (1997), Rose (2002) and Nickum et al. (2004) implies that only humans and primates are capable of experiencing pain. When this line of reasoning is followed, even new born babies lack a capacity to perceive feelings (Spruijt, 1999). The alternative approach starts from the awareness that internal and external states of fishes are probably not an all or nothing phenomenon, but that they are rather graded or limited to a range of subject matter appropriate for the animal's ecology. The authors following this line of reasoning, provided evidence that suggests fish have a capacity to perceive feelings.

Finally it is important to remember that "absence of evidence is not evidence of absence" (Sherwin, 2001). In the future, further research is necessary to come to a solid robust conclusion about the ability of fishes to experience pain and suffering. Only then can an objective assessment of fish welfare be accomplished.

1.3 Indications of fish welfare

To determine if an animal's welfare is compromised, different aspects of an animal's condition can be studied. Stress physiology, behaviour, health status, pathological changes of intestines and, in the case of food fish, flesh quality are often used to assess the welfare status of an animal. These five indicators are related to the function-based and nature-based definitions of animal welfare. Indicators related to feeling-based definitions are not common because science cannot yet, and perhaps cannot in principle, give empirical answers to many ethically relevant questions regarding the subjective experience of animals (Fraser *et al.*, 1997). The following section refers to the five indicator groups.

There is a consensus that no single parameter can provide adequate representation of an animals' welfare status, but there is no consensus on how to incorporate the different measurements into meaningful indices (B.N. North personal communication). Spooler *et al.* (2003) summarized ways that this can be done for terrestrial animals. Principal Components Analysis (PCA) combines the strengths of univariate post-hoc interpretation and multivariate analysis to objectively identify relative weightings of variables. It is difficult to apply systems for assessing welfare in terrestrial animals to fish. This has been particularly due to problems of observing fish especially in production systems. Here, measurements have been concentrated on individual aspects of welfare such as growth, stress or damage. However, these are influenced by factors other than welfare, thus they are often imprecise or noisy indicators of fish welfare. Turnbull *et al.* (2005) and North *et al.* (*Aquaculture* in press) used the statistical tool of multivariate analysis to combine simultaneously recorded measures of welfare on the basis of the observed statistical relationships among them. Turnbull *et al.* (2005) examined welfare in Atlantic salmon in cages on a commercial marine farm, exposed to different stocking densities over a period of 10 months. Four commonly used measures of fish welfare (e.g. condition of body and fins and plasma concentrations of glucose and cortisol) were registered from each cage and combined by multivariate analysis into one welfare score. North *et al.* applied the PCA to study the impact of stocking density on the welfare of rainbow trout in cages. Welfare was assessed by measuring a range of population (mortality, growth, size variation, FCR), individual morphometric (mass, fin condition) and physiological (haematocrit, plasma cortisol, lysozyme activity) indicators.

The studies suggested that while stocking density can influence the welfare of the fish in the cages, this is only one influence on the welfare and on its own cannot be used accurately predict or to control welfare.

1.3.1 Stress physiology

In fish, studies on stress responses play a prominent role in welfare research. A stress response is often seen as an indicator for impaired welfare. However, before relating the stress response of a fish to its welfare status, more information on the stress response of fish is required. Stress can be defined as a condition caused by external stimuli that disturbs the internal environment. To cope with these external stimuli a defence reaction, or stress response, is started. Stress responses can be divided into primary, secondary and tertiary stress responses.

- The *primary stress response* is characterised by a neurally stimulated release of adrenaline and noradrenaline from the chromaffin tissue and an activation of the hypothalamic-pituitary-interrenal axis (HPI). The hypothalamus is stimulated by a stressor to produce corticotropin releasing hormone (CRH) and thyrotropin-releasing hormone (TRH). Both hormones cause a release of pituitary peptides. The main peptides produced by the pituitary are adrenocorticotrophic hormone (ACTH), α -melanocyte-stimulating hormone (α -MSH) and endorphin. ACTH is known to stimulate the release of cortisol from interrenal cells (Wendelaar Bonga, 1997).
- *Secondary stress responses* occur due to the neuroendocrine changes of the primary stress response. A number of physiological processes associated with metabolism and immune capacity are involved. Under hormonal control, including cortisol and catecholamines, energy compounds such as glucose and free fatty acids (FFA) are mobilized (Wendelaar Bonga, 1997), and ammonical products, such as urea and ureic acid, discharged. In addition to these metabolic changes, the immune capacity is affected by the HPI-axis activity (Weyts *et al.*, 1999).
- *Tertiary stress responses* involve effects on the whole animal, disease outbreaks after prolonged stress, reduced growth rates and a lower reproductive capacity (Schreck *et al.*, 2001; Snieszko, 1974). Although stress responses are related to animal welfare, it is important to realize that a physiological stress response does not necessarily mean that an animal is suffering. A short-term stress response is a natural reaction that may be beneficial for the fish, depending on the situation. In the case of tertiary effects such as reduced growth rate and reproductive capacity, it is possible that they do indicate a compromised welfare status.

1.3.2 Behaviour

The fourth freedom of the Brambell Committee (see 1.2) states that animals should be free to express their natural behaviour and social interactions. In animal welfare research, behavioural studies have always played an important role. These studies include the possibility to use specific behavioural responses as an indicator for impaired welfare (such as ‘freezing’ in presence of a predator, flashing to reduce ectoparasites). Furthermore, choice tests have been performed to allow animals to express their natural preferences. Behavioural elements relevant to welfare in fish are:

- *Feeding activity*. The initial reaction of a fish to impaired welfare is often a sudden drop in food consumption (Plumb, 1994).
- *Swimming*. This involves the integrated effects of numerous physiological processes (Schreck, 1990). Assessment of swimming ability can provide a sensitive index to general stress in fish. Different swimming abilities are observed between fish exposed to stressful conditions compared with those not subjected to stress. The pattern of swimming – swimming into shallow water, swimming lethargically at the surface, lying listlessly on the pond or tank bottom, floating downstream or swimming erratically – can be indicators (Plumb, 1994). Critical swimming speed and the length of time a certain swimming velocity can be maintained may be used as indicators of welfare. Swimming performance can also be used as an indicator in toxicity tests (e.g. Shingles *et al.*, 2001; Wicks *et al.*, 2002; McKenzie *et al.*, 2003). From the viewpoint of aquaculture,

swimming behaviour offers a practical means by which farmers can assess the welfare of their stock e.g. crowding of fish around an inlet may indicate low dissolved oxygen.

- *Respiration* is also thought to be a useful indicator of fish welfare. A difference between resting and active metabolism can be quantified simply by counting breaths opercular beats per minute, or more precisely, by measuring respiratory gases. Respiratory movements and even oxygen consumption all provide measures of toxicological stress, which is related to welfare of the fish.

Other direct observations of behaviour, such as determination of a fish's ability to learn or remember a particular behaviour may also be a feasible indicator. However, considerable differences in behavioural responses between different fish species exist. Behaviour patterns may be used as a useful index of pain and can be validated if they are observed in groups exposed to noxious stimuli, but absent from untreated control groups.

1.3.3 Health status

Increased susceptibility to disease in fish may be the result of prolonged stress periods and is therefore an indicator of compromised welfare. The causes of diseases in fish are complex and the occurrence of diseases may indeed suppress the welfare status of an animal. However, diseases are not necessarily the result of poor environmental conditions and also occur in fish experiencing optimal conditions. Also the incidence of disease may not necessarily be caused by human mismanagement, since marked losses due to diseases also occur in populations of wild fish.

Stress physiology, behaviour and health status all provide information about the welfare status of an animal. However, it remains difficult to relate clearly each of these indicators to fish welfare. Preferably, all three aspects should be taken into account when assessing the welfare status of an animal.

1.3.4 Pathological changes

Chronic stress may also result in morphological and morphometric alterations. In European eels (*Anguilla anguilla*) it has been shown that stress can result in marked changes in the stomach (Peters *et al.*, 1980). The findings demonstrate that the functional capability of the stomach is heavily interfered with when the eels are subjected to severe chronic stress. Analogous observations can be made in mammals and humans.

1.3.5 Product quality

For farmed fish species such as Atlantic salmon (*Salmo salar*; Byrne, 2002; Berg *et al.*, 1997, Roth *et al.*, 2002; Sigurgisladottir, 2001), trout (Azam *et al.*, 1989, Byrne, 2002), eel (Morzel and Van de Vis, 2003) and turbot (Morzel *et al.*, 2003), it is shown that reduced stress during slaughter has a positive influence on flesh quality. Freshness, which is a major product quality attribute in the opinion of consumers, can be improved by reducing stress and muscle activity during slaughter (Morzel and Van de Vis, 2003; Byrne, 2002). Thus, it is of interest to the fishing industry to improve welfare aspects of the processing of catching and killing fish, as this may not only have the potential to improve product quality, but also the image of the fish farming industry, as perceived by consumers, governments and animal welfare associations.

CHAPTER 2 EFFECTS OF FISHERIES

2.1 General aspects of commercial inland fisheries

Commercial freshwater fisheries in most EIFAC countries are generally small scale operations. The fisheries usually operate with fyke-nets, gillnets, seines, traps and trawls. Eels are usually captured with fyke-nets and electrofishing. In many countries, commercial fishermen often leave the nets unattended for long periods, which results in confinement of the fish under bad conditions, and the eventual death of many less hardy (than eel) fish species. The information in the following section is not only from small scale inland fisheries but encompasses a much wider perspective because the information about handling of fish is mainly available from large scale marine fisheries. Consequently, the information must be interpreted with care when applied to inland fisheries.

In recent years concerns have been raised about the possible negative effects of commercial fisheries on fish stocks. Most of these concerns have been about the escape and post-release mortality. This mortality has been acknowledged in several fisheries, e.g. fish escaping through the meshes of a bottom trawl, fish discarded from a hand-line fishery or slipping by purse seines. A number of factors affect the mortality of discarded bycatch. One main factor is the selected gear type (Table 1). Problems with trawled gear depend on tow duration, catch composition and weight of the catch (Pálsson *et al.*, 2003). In catch and release fisheries problems depend on bait type, size of the fish and water temperature (Chopin and Arimoto, 1995). Defining escape and post-release mortality may be essential for stock assessments and for the improvement of fishing gear selectivity.

Table 1. Mortalities of fish escaping from fishing gears (based on Chopin and Arimoto, 1995)

Fishing gear	Species	Mortality (%)	Reference
Surrounding gear	<i>Scomber</i> sp.	0-90	Lockwood <i>et al.</i> , 1983
Seine nets	Cod, haddock	0, <10	Soldal and Isaksen, 1993
Seine nets	Striped bass	1-17	Dunning <i>et al.</i> , 1989
Seine nets	Freshwater drums	84.7	Fritz and Johnson, 1987
Trawls	Various	Varied	Wassenberg and Hill, 1989
Trawls	Haddock, whiting	9-27, 10-35	Sangster and Lehmann, 1989
Trawls	Gadoids	14-100	Main and Sangster, 1990
Trawls	Haddock, whiting	9-27, 10-35	Anonymous, 1993
Trawls	Cod, haddock	0, 1-32	Soldal <i>et al.</i> , 1991
Trawls	King and Tanner crabs	21-22	Stevens, 1990
Trawls	Lobster	21	Smith and Howell, 1987
Trawls	Atlantic halibut	65	Nelson <i>et al.</i> , 1989
Trawls	Herring	85-90, 75-85	Suuronen <i>et al.</i> , 1993
Trawls	Scup, flounder, cod	0-50, 0-15, 0	DeAlteris and Reifsteck, 1993
Trawls	Haddock, whiting	48-89, 60-86	Sangster <i>et al.</i> , 1996
Trawls	Herring	30-72	Suuronen <i>et al.</i> , 1996
Trawls	Sablefish, walleye pollock	0-25, 0-100	Olla <i>et al.</i> , 1997
Dredges	<i>Pecten</i> sp.	78-88	McLoughlin <i>et al.</i> , 1991
Dredges	<i>Placopecten</i> sp.	10-17	Caddy, 1973
Hoop nets	Paddlefish	35	Dieterman <i>et al.</i> , 2000
Gillnets and entangling nets	Pacific salmon	80-100	Thompson <i>et al.</i> , 1971
Gillnets and entangling nets	Pacific salmon	80	Thompson and Hunter, 1973
Gillnets and entangling nets	<i>Clupea</i> sp.	1.9	Hay <i>et al.</i> , 1986
Gillnets and entangling nets	Coho salmon	35-70	Buchanan <i>et al.</i> , 2002
Longlines	Pacific halibut	3-34	Kaimmer and Trumble, 1998
Handlines	Cod	32-54	Pálsson <i>et al.</i> , 2003

In addition to escape and post-release mortality, other negative effects of commercial fisheries have been identified. Several capture methods induce stress responses. Striped bass (*Morone saxatilis*) caught in gill nets and fyke nets exhibit significantly increased lactate, glucose, potassium and P_{CO2},

but lowered blood pH. Both capture methods induced secondary stress responses, although responses were greater in gillnetted fish (Hopkins and Cech, 1992). Capture of sea bream (*Pagrus major*) by hook and line and trammel nets significantly elevated cortisol concentrations (Chopin *et al.*, 1996).

2.2 Actions for commercial inland fisheries

With regard to the welfare of the fishes in commercial (inland) fisheries the following actions should be taken into account.

- All kinds of handling of landed fish on the vessel: sorting, discards, slaughtering. This not only refers to the mortality of the fishes, but also to the conditions in which the fishes are kept after capture (in tanks, buckets or alive without being kept in water), transported (density in tanks, temperature and oxygen conditions, tranquilizers) and brought to the market.
- The effects of the different fishing methods in respect to animal welfare. This refers to the optimization of fishing equipment and fishing methods to improve conditions for the captured fishes.

2.3 Fisheries management

Inland fisheries have a high degree of inter-relatedness with other users of the aquatic resource. In most areas of the world, the principal impacts on fisheries do not originate from the fishery itself, but from outside the fishery. Most aspects directed at the conservation and sustainability of the resource are under control of a wide range of interests with superior social and financial implications for society. The fishery has to be managed within the constraints imposed by other sectors and, while there is space for conventional management of fisheries, much attention is now paid to techniques for mitigation or rehabilitation of external impacts (FAO, 1997). In this context, the issue of fish welfare arises where the external activities have serious negative effects on fishes. These problems result from social and economic uses of river systems, and include:

- effects of hydro power plants;
- effects of damming on (migratory) species;
- flow regulation;
- river channelization;
- industrial and urban discharges (i.e. pollution);
- navigation;
- water abstraction.

Besides the negative effect on migratory movements within the river system, resulting in the disappearance of species like Atlantic sturgeon, salmon from rivers in Europe, the hydropower plants cause serious damage to down-stream migrating fish like eel.

Several measures in fisheries management affect fish welfare (Cowx and Gerdeaux, 2004; Cooke and Cowx, 2006). The well-being of individual fishes can be impaired and the reproductive potential of fish populations can be diminished. Several topics relevant to fish welfare can be identified in the management of recreational fisheries:

- Catch and release fisheries;
- Put and take fisheries;
- Stocking;
- Use of live bait fish;
- Keepnets;
- Harvest: the process of catching, handling and slaughter of wild fish;
- Protection from predators;
- Disease and parasites;
- Eradicating fish stocks (e.g. by poisoning);
- Biomanipulation of fisheries.

2.4 Catch and release fisheries

A growing public attention to animal welfare has been seen in the last decades. In recent years recreational fisheries has also become a subject of attention. In response to the growing practice of “Catch and Release” fishing, concerns have been raised about the physiological effects of angling and subsequent hooking mortality. Table 2 provides an overview of hooking mortality caused by angling in different fish species. Several studies investigated factors influencing fish welfare during angling. These factors can be divided in physiological factors, environmental factors and fishing method factors.

Table 2. Mortalities of angled fish (based on Chopin and Arimoto, 1995)

Fishing gear	Species	Mortality (%)	Reference
Hooks and lines	<i>Oncorhynchus</i> sp.	12-69	Vincent-Lang <i>et al.</i> , 1993
Hooks and lines	<i>Oncorhynchus</i> sp.	34-52, 40-86	Parker <i>et al.</i> , 1959
Hooks and lines	<i>Salmo</i> sp.	0	Wydoski, <i>et al.</i> , 1976
Hooks and lines	Rainbow trout	39, 3-5	Barwick, 1985
Hooks and lines	Cutthroat trout	0.3, 3	Schill <i>et al.</i> , 1986
Hooks and lines	Trout	0-8.6	Dotson, 1982
Hooks and lines	Smallmouth bass	0, 11	Clapp and Clark, 1989
Hooks and lines	<i>Esox</i> sp.	3	Schwalme and Mackay, 1985
Hooks and lines	Chinook salmon	9-32	Wertheimer, 1988
Hooks and lines	Pacific salmon	41	Milne and Ball, 1956

2.5 Physiological effects

Angling of a fish will result in a classical stress response characterised by an activation of the hypothalamus-pituitary-interrenal cells axis (HPI-axis), resulting in an increased plasma cortisol concentration (Ruane, 2002). Hooking and playing of red drum (*Sciaenops ocellatus*) resulted in increased plasma cortisol, plasma glucose, plasma lactate and plasma osmolality (Gallman *et al.*, 1999). Similar stress responses have been observed in blue mao mao (*Scorpius violaceus*; Lowe and Wells, 1996) carp (*Cyprinus carpio*; Pottinger, 1998), largemouth bass (*Micropterus salmoides*; Gustavson *et al.*, 1991) and kahawai (*Arripis trutta*; Davidson *et al.*, 1997). Increased plasma adrenaline, plasma noradrenaline, haemoglobin concentration and haematocrit have also been observed (Lowe and Wells, 1996). Furthermore, cardiac output, heart rate and stroke volume may increase after hooking and playing during angling simulations (Schreer *et al.*, 2001).

Tertiary stress responses may result in growth retardation and impaired reproduction. Growth of smallmouth bass (*Micropterus dolomieu*) was negatively related to the number of hooking events (Clapp and Clark, 1989). However, this relationship was not noticeable among older fish. An effect on reproduction was found in snapper (*Pagrus auratus*). Angling significantly effects sex steroids 17 β -estradiol, testosterone, cortisol and 17 α -, 20 β -dihydroxy-4-pregnen-3-one and suppresses ovulation (Carragher and Pankhurst, 1991). Suppressed estradiol levels were found in rainbow trout (*Oncorhynchus mykiss*; Pankhurst and Dedual, 1994).

Mortality of fish may occur after extreme physiological disturbance. Survival of the fish is dependent on the amount of stress caused by hooking and playing. Hooking location, water temperature and playing time are important factors affecting survival (Nelson, 1998; Thorstad *et al.*, 2003; Bettoli and Osborne, 1998). Hooking of fish in vital organs (oesophagus, gills) may result in high mortality. A mortality rate of 96 percent was found in deeply hooked Atlantic salmon (*Salmo salar*; Warner, 1979). In brown trout (*Salmo trutta*) highest mortalities were found in fish hooked in oesophagus or stomach (46.6 percent) or gills (39.6 percent) (Hulbert and Engstrom-Heg, 1980). In smallmouth bass (*Micropterus dolomieu*), 75 percent of the mortality resulted from fish being hooked in the oesophagus (Clapp and Clark, 1989).

Handling may also influence the welfare of fishes. Handling the fish will often result in air exposure, inhibiting oxygen and CO₂ exchange. The duration of handling will affect survival (Thorstad *et al.*, 2003). This duration depends on factors such as the experience of the angler, fish size and the type of

terminal tackle. Handling may take a relative long time when hooks are engorged or fish are manipulated for measurement of length and weight or for photographing. Besides oxygen deprivation, fish may lose scales and mucus.

Handling also induces physiological disturbances. In rock bass (*Ambloplites rupestris*), handling and air exposure resulted in increased heart rate and stroke volume (Cooke *et al.*, 2001). Recovery times varied between 2-4 hours depending on total air exposure. However, handling and air exposure did not lead to hooking mortality. Several other studies showed a minimal effect of handling stress on hooking mortality. No correlation was found between lake trout (*Salvelinus namaycush*) hooking mortality and the degree of handling (Loftus *et al.*, 1988). No mortality was observed in fish kept out of the water for as long as five minutes. Similar results were found for spotted sea trout (*Cynoscion nebulosus*; Hegen *et al.*, 1987). However, mortalities were found in rainbow trout (*Oncorhynchus mykiss*) after exposure to exercise and air (Ferguson and Tufts, 1992). In their study, no control fish died, 12 percent of exercised fish died and fish exposed to exercise and air exposures of 30 and 60 seconds, respectively, exhibited 38 percent and 72 percent mortality.

Different retention gears are used for storing the fish until release. Retaining rainbow trout (*Oncorhynchus mykiss*), brook trout (*Salvelinus fontinalis*) and carp (*Cyprinus carpio*) in keepnets causes a significant stress response, resulting in an increased plasma cortisol, plasma glucose and disturbed osmoregulation (Pottinger, 1998; Ruane, 2002). However, in these studies no mortalities were observed after retention in keepnets. Long-term effects of keepnet retention were also studied. Growth and survival were not affected in carp (*Cyprinus carpio*), roach (*Rutilus rutilus*), bream (*Abramis brama*), rudd (*Scardinius erythrophthalmus*) and ide (*Leuciscus idus*) after a 4-hour retention (Raat *et al.*, 1997). Indirect mortality may occur due to fungal lesions associated with abrasion (Cooke and Hogle, 2000). Other factors affecting survival after keepnet retention are water temperature and changing water quality (Pottinger, 1997; Hartley and Moring, 1993; Cooke and Hogle, 2000). Other retention gears used, such as gill stringers, may cause gill damage, which affects survival (Cooke and Hogle, 2000). The literature on fishes kept in keepnets is reviewed in the papers by Schreckenbach and Wedekind (see Bibliography in Appendix).

2.6 Environmental effects

An environmental factor known to affect fish welfare during angling is temperature: mortality increases with temperature, and lower mortalities at lower temperatures are associated with lower metabolic rates and physiological activity. Hooking mortality of striped bass (*Morone saxatilis*) during catch and release angling was 14 percent in November and December (average temperature 15°C) but 67 percent in July and August (average temperature 27–31°C) (Bettoli and Osborne, 1998). Cooke and Hogle (2000) found an increasing mortality of smallmouth bass (*Micropterus dolomieu*) with increasing water temperature (reaching 61 percent at 21.8°C). Comparable results were found in striped bass (*Morone saxatilis*; Nelson, 1998). Survival rates of angled Atlantic salmon (*Salmo salar*) were 20 percent at 20±2°C, 100 percent at 16.5±1°C and 100 percent at 8±1°C (Anderson *et al.*, 1998). Mortality of brook trout (*Salvelinus fontinalis*) caught on single- and treble-hook artificial lures was positively correlated to water temperature (ranging between 5.6-17.8°C; Nuhfer and Alexander, 1992). However, Hegen *et al.* (1987) found similar mortalities of spotted sea trout (*Cynoscion nebulosus*) between summer (26.9 percent) and winter (16.9 percent). Furthermore, channel catfish (*Ictalurus punctatus*) mortality in reservoirs was unrelated to water temperature (Ott and Storey, 1993).

Another factor influencing the fish welfare during angling is dissolved oxygen. Anglers using boats tend to store their fish in boat live-wells. This may result in the storage of fish in these wells for long periods at high densities. Poor water quality (low dissolved oxygen) and metabolites (ammonia) resulting from these high fish densities may significantly affect the mortality of caught and released fish. These problems may be more severe during warmer periods of the year when the dissolved oxygen content of the water is even lower. High fish density is an important factor in largemouth bass (*Micropterus salmoides*) and smallmouth bass (*Micropterus dolomieu*) mortality rates (Hartley and Moring, 1991). However, Schramm *et al.* (1987) did not find a relation between mortality rates and

fish density, although they reported initial, delayed and total mortalities of 9, 6 and 14 percent, respectively.

Salinity is a third factor to influence the welfare of angled fishes: hooking mortality in striped bass (*Morone saxatilis*) increases with decreasing salinity. Mortalities of striped bass (*Morone saxatilis*) caught using similar gear and techniques were 1.87, 38.81 and 70.39 percent at salinities of 8.0, 2.3 and 0.0 ppt., respectively (May, 1990).

2.7 Effects of fishing methods

Different types of hooks are used by anglers and they have differing effects. For example, longer handling times are expected in fishes caught on treble hooks because of multiple hook penetration, and added stress and difficulty associated with hook removal (Falk *et al.*, 1974). However, hooking mortality varies between single and treble hooks in different taxonomic groups. Brook trout (*Salvelinus fontinalis*) mortality was significantly higher in fish caught on single hooks than fish caught on treble hooks (8.3 and 2.4 percent, respectively; Nuhfer and Alexander, 1992). Hooking mortality of cutthroat trout (*Oncorhynchus clarki lewisi*) was 59 percent for fish caught on single hooks against 35-48.1 percent for fish caught on treble hooks (Titus and Vanicek, 1988). Payer *et al.* (1989) found walleye (*Stizostedion vitreum*) mortality was significantly higher in fish caught on baited hooks compared with those caught on treble-hook artificial lures.

However, no differences in hooking mortality was found between single and treble hooks in other studies. Mortality of red drum (*Sciaenops ocellatus*) caught on single and treble hooks was similar (4.92-3.33 percent, respectively) Similar results were found for cutthroat trout (*Oncorhynchus clarki lewisi*) caught with treble-hook artificial lures and baited single hooks (Hunsaker *et al.*, 1970) and for spotted sea trout (*Cynoscion nebulosus*) caught on single (4.69 percent) or treble (10 percent) hooks (Matlock *et al.*, 1993).

CHAPTER 3 EFFECTS OF AQUACULTURE

Aquaculture has grown substantially since the early 1990s. In 2001, 31 percent of all fish eaten in the European Union was supplied by aquaculture (European Commission, 2001). In recent years, attention has focused also on animal welfare aspects of aquaculture. Several aspects of husbandry practices in aquaculture have the potential to impact on the welfare of cultured fishes (e.g. site selection, confinement, stocking density, water quality, slaughter). As discussed in section 1.3 no single parameter can provide adequate representation of an animal's welfare status.

3.1 Handling

In aquaculture, fish may be handled during weighing, grading, vacinating and comparable husbandry practices. This physical disturbance evokes physiological stress responses in several fish species (Pickering, 1998; Waring *et al.*, 1996; Grutter and Pankhurst, 2000). There is also evidence showing that transportation can affect the disease resistance of a fish (Stangeland *et al.*, 1996; Mazur and Iwama, 1993), probably as a result of the immunosuppressing effect of corticosteroids release during stress response. Additional affects may occur through confinement or transportation associated with handling practices.

Short-term crowding or confinement may be used in aquaculture during weighing, grading and comparable husbandry practices. Physical confinement has been shown to increases plasma cortisol and glucose levels in most commercially valuable fish species (Barnett and Pankhurst, 1998; Montero *et al.*, 1999; Ortuño *et al.*, 2001; Vijayan *et al.*, 1997). Furthermore, it has an effect on the immune system of several species of fish (Garcia-Garbi *et al.*, 1998; Mazur and Iwama, 1993; Vazzana *et al.*, 2002). Reproduction is also negatively affected by confinement stress (Clearwater and Pankhurst, 1997). Finally, crowding can result in elevated fin erosion (Winfrey *et al.*, 1998; North *et al.* *Aquaculture* in press).

During transportation fishes are exposed to a range of stressful stimuli (handling, netting, loading at high densities, unloading, inadequate water exchange and poor water quality) that may cause adverse physiological reactions affecting essential life functions (Schreck *et al.*, 1995; Erikson *et al.*, 1997; Garcia *et al.*, 2000; Paterson *et al.*, 2003; Barton, 2000; Bandeen and Leatherland, 1997; Rouger *et al.*, 1998; Iversen *et al.*, 1998; Sandodden *et al.*, 2001; Robertson *et al.*, 1988). In 1986, EIFAC published a technical paper on the transport of live fish (Berka, 1986) containing guidelines. Prolonged recovery periods may be necessary after transportation. There is a need for safeguards in practise (e.g. water quality monitoring equipment) and standard operating procedures (e.g. starving fish before transport to minimise stress).

3.2 Stocking density

Stocking density is a factor in aquaculture that is commonly perceived to affect the welfare of fish. According to Ellis *et al.* (2002) and North (personal communication) it is almost certainly going to continue to be a major issue of contention that will be used against the aquaculture industry.

The rearing of fish under crowded conditions, such as high stocking density (defined as the weight per unit of volume, kg m^{-3}) or loading (defined as the weight per unit of flow rate, $\text{kg L}^{-1} \text{min}$), negatively affects growth (Pickering and Stewart, 1984; Procarione *et al.*, 1999), and influences metabolite levels (Vijayan *et al.*, 1990) and immune functions (Yin *et al.*, 1995; Tort *et al.*, 1996) in a number of fish species. This can be indirectly linked to increased competition for food, environmental factors such as deterioration in water quality, or more directly, due to an induction of a stress response through activation of the HPI-axis. Elevation of plasma cortisol and glucose levels was found for many fish species subjected to crowding (Yin *et al.*, 1995; Tort *et al.*, 1996; Rotllant and Tort, 1997; North *et al.*, *Aquaculture* in press). However, not all fish species react negatively on increased stocking densities. Arctic charr (*Salvelinus alpinus*) and halibut (*Hippoglossus hippoglossus* L.) reacted positively to increased stocking densities (Jørgensen *et al.*, 1993; Greaves, 2001; Alanärä and Brännäs, 1996). In addition, social interactions of fish strongly depend on stocking densities.

The review of Ellis *et al.* (2002) about the relationships between stocking density and welfare in farmed rainbow trout gives an insight in the key issues involved. Increased stocking density does not appear to cause prolonged crowding stress in rainbow trout. However, commonly reported effects of increasing density are reductions in food conversion efficiency, nutritional condition and growth, and an increase in fin erosion (North *et al.*, *Aquaculture* in press). Such changes, and especially the effects of density on fin damage, which is the most consistently reported affect in most studies, are indicative of a reduced welfare status. According to the authors there is dispute as to the cause of the observed effects of increasing density, with water quality deterioration and/or an increase in aggressive behaviour being variously proposed. Turnbull *et al.* (2005) in a study on cage farmed Atlantic salmon concluded, that stocking density can influence the welfare of the fish in production cages. However, this is only one of the husbandry factors that have an influence on their welfare and density on its own cannot be used to accurately predict or to control welfare.

3.3 Water quality

Ellis *et al.* (2002) summarized the importance of water quality in aquaculture. This section is therefore mainly based on the review of these authors.

Water is the medium in which fish have to meet both their physiological and spatial needs. The water is the source of oxygen and serves to dilute and remove metabolic wastes. Theoretically, increasing density will degrade water quality because greater numbers of fish are respiring and metabolizing in a given volume of water. Attention has been drawn to oxygen and ammonia as the water quality parameters generating the observed effects of increasing density (e.g. Rosenthal *et al.*, 1984). However, with the increasing use of recirculating and oxygenation systems, it has been recognized that carbon dioxide and nitrogenous waste products other than ammonia can also have adverse effects. In addition, an increase in density will increase the amounts of suspended solids in the water column - due to both greater faecal production and more fish movement (which will prevent particles from settling). Alabaster and Lloyd (1980), Boyd (1982), Meade (1985) and Wedemeyer (1996) have comprehensively reviewed water quality in relation to freshwater fish.

3.3.1 Dissolved oxygen

Dissolved oxygen (DO) is essential for fish respiration. It is assumed that above a critical DO concentration, consumption is independent of concentration. However, below the critical level, oxygen consumption is increasingly restricted by concentration, until the fish can only extract that amount which is required for standard metabolism. Below this, asphyxia occurs at the lethal level. Recommendations for a minimum DO level for farmed trout to avoid sub-lethal effects are similar to levels recommended for warm water fish, typically 5 to 6 mg L⁻¹ (e.g. Shepherd and Bromage, 1988). In contrast, Wedemeyer (1996) has suggested a minimum of between 7 and 9 mg L⁻¹ depending upon temperature, and Alabaster and Lloyd (1980) suggested an annual 50 percentile value of 9 mg L⁻¹ for wild salmonids.

3.3.2 Ammonia

Fish excrete nitrogen as ammonia, which reacts with water to form ammonium ions in an equilibrium reaction which is temperature and pH dependant (Boyd, 1982; Piper *et al.*, 1982).



The un-ionized ammonia (UIA) is highly toxic to fish, whereas ionized ammonium (IA) is considered relatively non-toxic (Alabaster and Lloyd, 1980; Boyd, 1982; Piper *et al.*, 1982). This difference in toxicity is thought due to the greater potential for UIA to diffuse across the gill membrane (Smart, 1981). The two forms are not differentiated by analytical methods, which measure total ammonia. In order to gauge whether levels are likely to be detrimental to fish, total ammonia needs to be converted to UIA concentration using published tables of percentage UIA in relation to temperature and pH (Boyd, 1982; Piper *et al.*, 1982). Trout are amongst the most sensitive fish species to UIA (Alabaster and Lloyd, 1980). Reports of levels of UIA that affect rainbow trout are variable, being dependent upon environmental factors (e.g. DO and CO₂ concentrations, temperature, concentration of other

ions, prior exposure), duration of exposure and the ontogenetic stage (Alabaster and Lloyd, 1980; Meade, 1985).

3.3.3 Effects of deterioration in water quality on welfare

The potential for water quality deterioration to lead to acute welfare infringement is well known. Increasing density increases the probability of episodic mortality when water supply, aeration or oxygenation systems fail and therefore requires greater supervision and appropriate back-up equipment (Piper, 1970; Pickering *et al.*, 1991). Water quality deterioration may also act chronically to reduce welfare status. There is evidence that both low DO and high UIA levels can act as chronic stressors to rainbow trout, elevating plasma cortisol levels (Scapigliati *et al.*, 1999). Pickering *et al.* (1991) noted that plasma cortisol levels were raised in fish exposed to low DO levels, being higher in fish held at a higher density. Interpretation is complicated by the facts that plasma cortisol levels change with exposure duration (Swift, 1981), and various linked water quality factors (pH, UIA, DO, CO₂) interact to affect the magnitude of the stress response (Donaldson, 1981).

Stress in fish farming can be associated with a variety of environmental conditions, such as reduced oxygen level (Wedemeyer, 1996;), abrupt variation in temperature (Wagner *et al.*, 1997; Davis and Parker, 1990), appearance of contaminants (Wedemeyer, 1996) or elevated ammonia levels.

A common sublethal effect of ammonia is damage to the gills (Alabaster and Lloyd, 1980). Rosenthal *et al.* (1984) related a decrease in length of the primary gill lamellae to increasing ammonia and density. Soderberg *et al.* (1984) and Soderberg (1985) related observed pathologies of secondary gill filaments (epithelial oedema, blood filled aneurysms, hyperplasia, fusion) to UIA levels. Ammonia exposure has also been shown to decrease the number of erythrocytes (Alabaster and Lloyd, 1980). Such effects on the gills and blood may account for the increased ventilatory rate observed in response to ammonia (Alabaster and Lloyd, 1980). Soderberg *et al.* (1984) and Soderberg (1985) associated a reduction in vacuolation of liver hepatocytes with chronic exposure to ammonia, and suggested that this was due to a reduction in food intake. Reduced food intake would reduce growth rate, which is a common effect of chronic exposure to ammonia (see Soderberg *et al.*, 1983; Meade, 1985).

Bosakowski and Wagner (1994) correlated UIA levels with increasing fin erosion. Soderberg *et al.* (1983) correlated mortality with ammonia levels and suggested that exposure to ammonia predisposed the fish to infection by ectoparasitic protozoans. In an epidemiological survey of commercial farms, Ortega *et al.* (1996) demonstrated ammonia level as a factor affecting the risk of bacterial infection. Indeed low DO and elevated UIA levels are commonly associated with a range of diseases (Wedemeyer, 1996). Water quality deterioration therefore has the potential to reduce welfare status by reducing nutritional status and causing physiological stress, injury to gills and fins and increasing susceptibility to disease.

One of the possible secondary effects of poor water quality is the development of cataracts. Cataracts have been mainly studied in salmonids (Bjerkås *et al.*, 2001; Bjerkås and Bjornestad, 1999; Iwata *et al.*, 1987), but are also found in seabass and sea bream (Bjerkås *et al.*, 2000). They can be caused by variations in water temperature (Bruno and Raynard, 1994; Bjerkås and Bjornestad, 1999; Bjerkås *et al.*, 2001) or toxic agents in the water (Fraser *et al.*, 1990). However, these cataracts may also be caused by nutritional deficiencies (Hargis, 1991), parasites (Ashton *et al.*, 1969), exposure to sunlight (Cullen *et al.*, 1994), hereditary factors (Kincaid, 1989) or rapid growth (Waagbø *et al.*, 1996; Waagbø *et al.*, 1998).

3.3.4 Effect of environmental factors on water quality deterioration

It is well recognized that the biomass of fish that any culture system can support depends upon environmental factors. Carrying capacity is determined by the rates of provision and consumption of DO, and the production, removal and toxicity of metabolic products (Piper, 1970). Hence carrying capacity is dependent upon factors such as water exchange rate, temperature, pH, fish size and feeding rate (Piper *et al.*, 1982). The quality and amount of inflow water will determine whether density results in water quality deterioration to sub-lethal or lethal levels. Natural water sources vary in DO,

NH₃ and CO₂, and the quality of inflow will be of greatest significance when farms reuse water, either via passage through successive systems or by recirculation. Inflow rate and system volume will determine the rate of water replacement, and hence provision of DO and dilution of metabolic wastes. The importance of inflow rate in affecting the biomass of fish that can be supported is well recognized (Piper *et al.*, 1982). Weight specific metabolic rate is well known to decrease with increasing fish size. DO consumption, therefore, has been reported to decrease with increasing fish size (Piper *et al.*, 1982). It is therefore commonly assumed that the physiological requirement for water decreases with increasing fish size. Temperature is a vital factor, determining the rate of metabolism of fish, solubility of DO, and the toxicity of ammonia (Alabaster and Lloyd, 1980). Temperature and fish size have been factored into estimates of carrying capacity (Piper, 1970). Similarly, pH will interact with density by altering the equilibrium of the two forms (IA, UIA) of ammonia (Israeli-Weinstein and Kimmel, 1998). Feeding reduces the DO level of the water (Wagner *et al.*, 1995). It has been suggested that densities may be constrained by limiting the amount of food that is fed (Wedemeyer, 1996). DO levels in the water are commonly increased by aeration and oxygenation. Traditionally, paddle wheels or air stones were used to aerate, although the addition of pure oxygen is now becoming increasingly common. Numerous recent studies have demonstrated that such supplemental oxygen enables trout rearing at higher densities thereby increasing the overall production potential of a system.

3.4 Social interaction

Ellis *et al.* (2002) in their review of 43 studies on the effects of density on production concluded that only inferential evidence exists that aggressive behaviour generates the observed effects of increasing density, whereas there is direct experimental evidence that water quality degradation is responsible. They also conclude that the potential for welfare to be detrimentally affected by non-aggressive behaviour interactions (abrasion, collision, obstruction) and low densities (due to excessive aggressive behaviour and a poor feeding response) have been largely overlooked. According to the authors legislation directly limiting stocking densities is likely to be unworkable, and a more practical option might be to prescribe acceptable levels of water quality, health, nutritional condition and behavioural indicators.

Social interactions have significant influence on the stress responses of fishes. Classification of stressors affecting behavioural, physiological and performance responses:

- Animal-related factors: genetics, coping styles, sex, age;
- Nutritional factors: composition, feeding strategy;
- Water quality related factors: temperature, pH, O₂;
- Environmental factors: stocking density, photo-period, light intensity.

Indicators of behavioural stress responses are:

- Swimming activity;
- Air breathing;
- Agonistic behaviour;
- Escaping behaviour;
- Feeding behaviour;
- Occurrence of stereotypes.

Hierarchies are formed, comprised of a group of dominant individuals at the top, followed by a number of sub-dominants and thereafter a number of subordinates with low rank positions in the hierarchy (Symons (1970). An elevation of plasma cortisol concentration is characteristic for subordinate fishes (Laidley and Leatherland, 1988; Pottinger and Pickering, 1992; Øverli *et al.*, 1999). Other physiological changes shown by subordinate fishes are decreases in weight (Pottinger and Pickering, 1992), hepatic glycogen content (Ejike and Schreck, 1980), and disease resistance (Peters *et al.*, 1988) and increases in interrenal cell activity (Noakes and Leatherland, 1977), plasma glucose concentration (Peters *et al.*, 1988) and standard metabolic rate (Sloman *et al.*, 2000). Furthermore, aggression among fishes may cause injury, especially when competition for food is strong (Greaves

and Tuene, 2001). North et al. (*Aquaculture* in press) demonstrated that stocking density had a significant effect on fin condition.

3.5 Light regime

In aquaculture lights are used to control smoltification of Atlantic salmon, promote growth (most salmonids), reduce maturation (salmon, cod, haddock) and brood stock management (most other species). Light plays an important role in the growth and development of fishes. Light can influence the behaviour of fishes, through variation in intensity, wavelength and photoperiod. This influence differs between species. Bromage *et al.* (2001) summarized the environmental regulation of maturation in farmed finfish with special reference to the role of photoperiod and melatonin. According to these authors current evidence indicates that photoperiod is the principal proximate cue in many fish and that this entrains an endogenous rhythm which in turn controls reproduction. This information is thought to be transmitted to the reproductive axis by diel and seasonal changes in melatonin. It is probably that temperature, nutritional status or size together with other factors act in a permissive way, possibly by gating mechanisms, to enable maturation to proceed.

Haddock larvae (*Melanogrammus aeglefinus*) showed better growth under higher light intensity (Downing and Litvak, 1999). By contrast, halibut (*Hippoglossus hippoglossus*) yolk-sac larvae developed abnormally in the presence of light (Bolla and Holmefjord, 1988). Studies on the effect of photoperiod on the growth and survival of fish larvae have produced mixed results. In a continuous light regime, rabbitfish larvae (*Siganus guttatus*; Duray and Kohno, 1988) and gilthead sea bream larvae (*Sparus aurata*; Tandler and Helps, 1985) grow and survive better while seabass (*Dicentrarchus labrax*; Barahona-Fernades, 1979) larvae exhibit reduced growth and survival.

According to North (personal communication) in terms of direct effects on fish welfare, there are studies that showed that lights can be used to adjust swimming depth and stocking density in Atlantic salmon (Juell and Fosseidengen, 2004). There are lots of aspects of welfare that are negatively affected when fish pass through sexual maturation, so if this can be controlled by lights there is the argument that lights can be used to promote welfare. North further commented that although there isn't much literature to date, the increased use of artificial lights in the industry and rapid developments in lighting technology are sure to make this an issue for the future.

3.6 Slaughter

All slaughter methods are stressful, but some are better than others (Robb *et al.*, 2000). Fishes can be killed by injectable agents, external agents, carbon dioxide, physical methods or cooling. However, not all of these methods are used in commercial aquaculture. Methods used in aquaculture are:

- Percussive stunning by blow to the head: carried out manually using a hand-held club called a "priest" for stunning, followed by cutting of the gill arches to bleed the fish.
- Carbon dioxide stunning: fish are placed in a seawater bath saturated with carbon dioxide. This environment is highly aversive with fish being observed to shake their heads and tails vigorously trying to escape (Robb *et al.*, 2000). Movement ceases after 30 seconds, but sensibility may not be lost for 4 – 5 minutes (Robb *et al.*, 1997). Bleeding after CO₂ stunning is essential to avoid fish recovering.
- Bleeding without prior stunning: gill arches are cut with a knife and fish bleed to death.
- Electrocution: electric current is passed through the water containing the fish. The electric current must be sufficient to stun and kill the fish, otherwise the fish are only paralysed and could suffer considerably. This method can also cause spinal fractures and haemorrhaging.
- Asphyxiation in air or on ice: fishes are hauled out of water and allowed to die through suffocation in air. Alternatively, they are removed from water into bins containing ice.

Many of the slaughter methods currently used for farmed fish (e.g. death in air, death in an ice slurry, freezing, bleeding, carbon dioxide narcosis in combination with bleeding) do not provoke immediate unconsciousness until death occurs (Robb and Kestin, 2002; Van de Vis *et al.*, 2003). When a fish is effectively stunned (immediate and permanent loss of consciousness) prior to killing, the slaughter process may be considered humane. It has been established that percussive stunning (a blow to the

head by using an instrument), captive needle stunning (injection of air under pressure into the brains) and the application of electrical current (provided that appropriate conditions are used) can provoke immediate and permanent loss of consciousness in fish (Robb and Kestin, 2002; Lambooij *et al.*, 2002; Van de Vis *et al.*, 2003). Several researchers reported that various species recovered from the application of an electrical current for one second (Lambooij *et al.*, 2002, Lines *et al.*, 2002; Robb and Kestin, 2002; Van de Vis *et al.*, 2003b). Recovery can be prevented when fish are exposed to electricity for 10 seconds or more and subsequently a killing method, e.g. rapid cooling is applied. This combination of methods results in permanent loss of consciousness until death occurs.

In Australia, a chemical substance, eugenol, is added to the water to stun fish prior to killing. Eugenol is a food grade substance, which is based on clove oil. Barriers to its use in, for instance the United Kingdom, include the cost of overcoming the legislative requirements to introducing a new medication and the possible public response to eating fish that could be perceived as having been poisoned (Lines *et al.*, 2003).

Small warm-water fishes, such as seabass, killed by chilling on ice water had lower plasma glucose and lactate levels and showed less marked behavioural responses than those killed by other methods, especially asphyxiation (Poli *et al.*, 2002; Skjervold *et al.*, 2001). Electrocutation may be less harmful for larger fishes such as trout (FSBI, 2002).

For practical application of electrical methods one has to take into account the effect that water chemistry has on the requirements for current and voltage for stunning fish (North, personal communication). However, it is anticipated that electrical methods for stunning will be feasible for industrial application, as the large numbers of individual fish that need slaughtering after harvest make individual percussion or captive needle stunning and bleeding impractical. Percussive stunning of Atlantic salmon is an exception, as equipment for automated percussive stunning was developed, especially for this species by the Australian company Sea Food Innovations in collaboration with Marine Harvest, of the major producer of Atlantic salmon in Scotland. In 2002, Marine Harvest won a welfare award for introducing percussive stunning as a method for humane slaughter of Atlantic salmon (Anonymous, 2002). For other fish species, which differ substantially in anatomy and morphology, high-speed machinery to separate, orientate, percuss and bleed them would be complex and expensive (Lines *et al.*, 2003), as the developed equipment for salmon is not been adapted to other species such as turbot. Moreover, for fish species such as gilt-head seabream and eel, percussive stunning is not feasible without severe damage to the head, which is not acceptable for the industry (Van de Vis *et al.*, 2003). For fish species with a thick skull, such as African catfish, the application of percussive stunning does not result in immediate loss of consciousness and is therefore unfit for use (Van de Vis and Lambooij, unpublished results).

To date, at least one device is commercially available for electrical stunning of Atlantic salmon (Roth, pers. comm.). For rainbow trout a prototype for electrical stunning was developed (Lines *et al.*, 2003). However, the stunners were designed primarily for Atlantic salmon and trout and it is unlikely that these stunners are suitable to use for other fish species such as, eel, tilapia, seabass and turbot. This is because the optimal conditions for electrical stunning with respect to immediate loss of consciousness and prevention of carcass damage depend on the fish species. Laboratory trials found that the optimal conditions for eel are an electric field strength of 1250 V/m for 1 s followed by 313 V/m for 5 min in combination with deoxygenation of the water (Lambooij *et al.*, 2002; Morzel and Van de Vis, 2003), whereas the established optimal conditions for Atlantic salmon and trout are stunning for 6-12 s in a field of 75 V/m using sinusoidal 1000 Hz current (Robb and Roth, 2002; Roth, *et al.*, 2002) and 60 s in a field strength of 250 V/m, using a sinusoidal 1000 Hz current (Lines *et al.*, 2003), respectively. When a 50 Hz sinusoidal current is applied for salmon or trout, this will cause carcass damage, which is not observed if the same current is applied to eel (Morzel and Van de Vis, 2003) or African catfish (Van de Vis *et al.*, 2003b). Similar to salmon and trout, applying a 50 Hz current for stunning of turbot also causes carcass damage (Morzel *et al.*, 2003). The conditions for effective stunning of tilapia, seabass and turbot, whilst maintaining high product quality standards differ from those established for other species.

In Germany, only electrical stunning is the only permitted method of slaughter for eels (since April 1999). However, the method is not considered humane, according to prescribed conditions in the German legislation, as the eels are not rendered unconscious immediately (Lambooij *et al.*, 2002; Van de Vis *et al.*, 2003), and most animals die of exhaustion during application of the electric field. These authors have also established that eel cannot be rendered unconscious permanently when only electricity is used, as prescribed in the German legislation. Equipment available for electrical stunning of eel on a commercial scale, which is designed to meet the demands of the German legislation, is therefore not suitable for humane slaughter. The current method for slaughtering of eel in the Netherlands involves desliming live eels using salt followed by gutting (evisceration). This is also not a humane method (Verheijen and Flight, 1997; Van de Vis *et al.*, 2001). Current industrial methods applied to the other species selected in the proposed research (tilapia, seabass and turbot) are described by Robb and Kestin (2002). For seabass and tilapia, the use of an ice/water slurry is the first step in the slaughter process. The rapid cooling does not result in an immediate loss of consciousness and is most probably stressful, as demonstrated in salmon, eel and gilt-head seabream (Lambooij *et al.*, 2002b; Skjervold *et al.*, 2001; Van de Vis *et al.*, 2003). Tilapia is also killed by removing the fish from water, which results in asphyxiation. Temporary removal from the water is widely used as an experimental stressor in studies of the stress response of fish. Turbot is killed by gutting the gills or the caudal vein and subsequently placing the fish in an ice/water slurry. During the procedure, vigorous activity occurs, which may indicate stress (Morzel *et al.*, 2003).

Stressful slaughter methods generally lead to fish flesh that is prone to gaping, with an unappealing colour, a soft texture and a low water holding capacity. More humane handling at slaughter, with reduced stress and physical activity immediately prior to death, results in better carcass quality (Robb and Kestin, 2002). For various farmed fish species, it has been observed that reduced stress at the process of slaughter has a positive influence on the flesh quality (Azam *et al.*, 1989; Berg *et al.*, 1997; Byrne, 2002; Kestin *et al.*, 1995; Roth *et al.*, 2002; Sigurgisladottir, 2001; Morzel and Van de Vis, 2003). Reduction of stress at slaughter by applying electrical stunning under specific conditions was found to be humane and generally promote better fish quality (Kestin *et al.*, 1995; Lines *et al.*, 2003; Morzel and Van de Vis, 2003; Robb and Kestin, 2002).

Research on electrical stunning of fish has resulted in a few patents in the area (Van de Vis, 2001, 2001b; Moller 2001). However, the parameters associated with electrical stunning are complex and a patent is not a protocol for effective electrical stunning with high standards of product quality for fish species. The patents do not contain details for optimal electrical stunning of eel, tilapia, seabass and turbot under various conditions. Major factors that determine whether electrical stunning is optimal (immediate loss of consciousness and high product quality) are the size, shape of the fish, fish species and the electrical conductivity of the fish and of the water). Hence, no satisfactory electrical stunning equipment exists for stunning of eel, tilapia, seabass and turbot on a commercial scale.

CHAPTER 4 FISH IN SCIENTIFIC RESEARCH

4.1 Research and monitoring of fish and fish ecology

Large numbers of fish are sampled or reared for scientific purposes throughout Europe. Only a very small proportion of these are used as experimental animals in the traditional sense (to test for effects of various treatments to predict effects on humans). Most fish are caught in the process of monitoring the status of stocks in rivers and lakes. This shows significant linkages to fish ecology studies and includes resource and stock assessment for studies on population dynamics. There has been some focus on the welfare of fish brought to the laboratories and kept for experimentation, and in most countries rules apply for the application of invasive tagging methods on fish. It is, however, also important to consider the ethical aspects of collecting fish in field studies, both in terms of numbers killed and the methods used.

4.2 Fields of interest

Special fields of interest relating to fish welfare in scientific studies are summarized below based on the discussion during the Workshop on Handling of Fishes in Utrecht, The Netherlands, 24-26 March 2004. Information about the welfare aspects for many of these topics is scarce in the literature.

- Tagging and marking. A relevant report of concerted action within the EU framework FAIR was published (Davenport *et al.*, 2002).
- Stunning and consciousness of fish.
 - Most anaesthetics and related chemicals used for stunning the fishes result in enhanced brain activity and increase in heart rate.
 - In most EIFAC member countries anaesthetics for fisheries use are not certified (in several countries, only MS 222 is certified).
 - There is a need for better understanding and certification of anaesthetics for use in fish handling and fish research (e.g. clove oil, benzocaine).
- Implantation techniques:
 - The practice of implantation differs widely in EIFAC member countries with regard to legislation, experience, procedures and protocols.
 - International cooperation in training for the use of invasive methods and techniques in EIFAC member countries is advisable. EIFAC should play a co-ordinating and stimulating role for initiatives for concerted action.
- Parasite infections.
- Sampling of fishes:
 - There is a need for more information on the fish welfare aspects of fishing methods (e.g. electrofishing, gillnetting).
 - Standardization of fishing methods should take into account basic information on fish welfare aspects.
 - In monitoring and sampling programmes, more attention should be paid to alternative methods, which leave the fishes in their natural environment (e.g. acoustic surveys).

4.3 Electric fishing

Concerns have also been raised about the possible negative effects of electric fishing. Electric fishing is a highly effective and widely used method for collecting fish in streams, rivers or lakes. Although this fishing method has some advantages over other, more conventional, methods it may pose some problems regarding the welfare of fish. Several studies showed that electric fishing causes haemorrhaging to varying degrees depending on species and type of current. Percentages of haemorrhages varied between 0-50 percent in crappies (*Pomoxis* sp.; Dolan *et al.*, 2002), 60 percent in brown trout (*Salmo trutta*; Habera *et al.*, 1999) and 78-92.4 percent in rainbow trout (*Oncorhynchus mykiss*; Schill and Elle, 2000). Furthermore, spinal injuries were observed in 9-45 percent of studied rainbow trout (*Oncorhynchus mykiss*), brown trout (*Salmo trutta*), cutthroat trout (*Oncorhynchus clarki lewisi*) and crappies (*Pomoxis* sp.) (Ainslie *et al.*, 1998; Habera *et al.*, 1999; Dwyer *et al.*, 2001; Dolan *et al.*, 2002).

Negative effects were also observed in early developmental stages. Mortality of 85 percent was observed in brook trout (*Salvelinus fontinalis*) eggs 21 days post-fertilization; at 37 days post-fertilization, 14 percent mortality was observed. Furthermore, 7-22 percent physical anomalies were observed (Keefe and Whitesel, 2000). A similar study on Arctic grayling eggs (*Thymallus arcticus*) showed that mortality was affected by electroshock, although the cost to the population was low (Roach, 1999). A study on chinook salmon (*Oncorhynchus tshawytscha*) showed 12-93 percent mortality, with highest mortalities in the early eyed stage (Cho *et al.*, 2002).

Despite electric fishing being harmful for fish (and people), it is a good, non-lethal method to sample fish in small rivers and streams, if performed correctly under good conditions (Cowx and Lamarque, 1990). There are several cases where the same individuals are sampled again and again during repeated electric fishing, and recent telemetric studies where fish were caught by electric fishing did not show any negative effects of the capture method (Jepsen and Aarestrup, 1999; Jepsen and Berg, 2002). (The draft document Best Practice in Electric Fishing is posted on the EIFAC Web site at <http://www.fao.org/WAICENT/FAOINFO/FISHERY/body/eifac/WGFishMon/default.asp>).

CHAPTER 5 LEGISLATION AND REGULATION

To ensure the welfare of fish in aquaculture, commercial fisheries and recreational fisheries, sufficient legislation and regulation is necessary. In this chapter legislation and regulation concerning animal welfare in Europe are reviewed, and to some extent, also discussed with reference to handling of fishes.

5.1 International legislation and regulation

5.1.1 European Union

The European Commission, Directorate-General for Fisheries published a report on “Farmed fish and welfare” (Wolffrom, 2004). This report gives references to the present opinions and regulations on animal welfare and the relevance to fish culture and the various institutions involved in this issue. The following section is taken from this EC report.

Definitions (www.euabc.com)

Resolution: A resolution is a non-binding statement, which defines objectives and makes political declarations. The European Council’s resolutions set out the direction of future policy initiatives. Resolutions may be used by the EU Court to interpret laws. They may be referred to as a form of “soft law”.

Recommendation: A non-binding decision, which only urges Member States to comply. A Member State cannot be fined for the breach of recommendations.

Convention: Conventions are international legal instruments laying down ethical principles for the transport, the farming, the slaughtering of animals as well as for their use for experimental purposes and as pet. They have been used as a basis for and continue to influence all the relevant legislation in Europe.

Directive: Directives are to be transferred into national law through the Member States’ parliaments and governments within 18 months. Through the years, the EU Court has proclaimed many directives to be directly applicable and even declared that countries are liable to pay compensation if they have not implemented a directive in time. Directives are normally transformed into national laws by the national parliaments or most often by the governments through delegated acts.

Decision: A EU decision is binding on the persons, companies or Member States mentioned in the decision. It is not generally binding, as is the case with a regulation.

Regulation: A EU decision that directly binds all Member States and citizens in the whole of the EU. Whereas directives need to be “transformed” into national law, regulations are directly applicable. It is therefore forbidden to change EU regulations when putting them into national laws.

EC legislation and regulations

European Convention for the protection of animals kept for farming purposes *Official Journal* L 323 , 17/11/1978 p. 0014 - 0022

78/923/EEC: Council Decision of 19 June 1978 concerning the conclusion of the European Convention for the protection of animals kept for farming purposes *Official Journal* L 323 , 17/11/1978 p. 0012 - 0013

Council Directive 98/58/EC of 20 July 1998 concerning the protection of animals kept for farming purposes *Official Journal* L 221 , 08/08/1998 p. 0023 – 0027

2000/50/EC: Commission Decision of 17 December 1999 concerning minimum requirements for the inspection of holdings on which animals are kept for farming purposes (notified under document number C(1999) 4534) (Text with EEA relevance) *Official Journal* L 019 , 25/01/2000 p. 0051 – 0053

Council Directive 91/628/EEC of 19 November 1991 on the protection of animals during transport and amending Directives 90/425/EEC and 91/496/EEC *Official Journal* L 340 , 11/12/1991 p. 0017 - 0027

Council Directive 95/29/EC of 29 June 1995 amending Directive 91/628/EEC concerning the protection of animals during transport *Official Journal* L 148 , 30/06/1995 p. 0052 – 0063

Council Resolution of 19 June 2001 on the protection of animals during transport, Official Journal C 273 , 28/09/2001 P. 0001 – 0001

Council Regulation (EC) No 1040/2003 of 11 June 2003 amending Regulation (EC) No 1255/97 as regards the use of staging points Welfare of animals during transport (details for horses, pigs, sheep and cattle). *Report from the Scientific Committee on Animal Health and Animal Welfare*, adopted on 11 March 2002.

http://europa.eu.int/comm/food/fs/sc/scah/out71_en.pdf

Opinion of the Scientific Panel on Animal Health and Welfare on a request from the Commission related to the welfare of animals during transport. Adopted 30 March 2004. http://www.efsa.eu.int/science/ahaw/ahaw_opinions/424/opinion_ahaw_01_atrans_ej_44_en1.pdf

Opinion of the Scientific Panel on Animal Health and Welfare on a request from the Commission related to welfare aspects of the main systems of stunning and killing the main commercial species of animals. (Question N° EFSA-Q-2003-093). Adopted by the AHAW Panel on the 15th of June 2004.

http://www.efsa.eu.int/science/ahaw/ahaw_opinions/495_en.html

5.1.2 *European Food Safety Authority (EFSA)*

The European Food Safety Authority (EFSA) Scientific Panel on Animal Health and Welfare was asked by the Commission services to report on the welfare aspects of the main systems of stunning and killing in the main commercial species of animals with consideration of Directive 93/119/EC. Species referred to in the Opinion (ESFA, 2004)³ are: cattle, sheep, pigs, poultry, horses and farmed fish. Welfare aspects of the systems for stunning other species, such as rabbits, deer, ratites or goats, have not been included in the opinion. Procedures appropriate to cattle, sheep, pigs, chickens, turkeys, farmed fish and horses and their related minimum requirements such that unconsciousness and insensibility are induced and poor welfare is minimised, are recommended.

5.1.3 *Council of Europe*

The European Union has adopted five Conventions drawn up by the Council of Europe (European Treaty Series: ETS) to control the use of animals by humans. The Conventions relate to animal welfare for animals handled during international transport (ETS 65, 1968), animals kept for farming purposes (ETS 87, 1976), animals for slaughter (ETS 102, 1979), animals kept as pet animals (ETS 125, 1987), and vertebrate animals used for experimental and other scientific purposes (ETS 123, 1986).

European Convention for the Protection of Animals During International Transport

The Convention sets compulsory norms for space, ventilation and hygiene, transportation means, food and water, loading and unloading of animals and veterinary assistance for the international transport of animals.

Article 46 of the Convention applies specifically to cold-blooded animals including fishes. It states "cold-blooded animals shall be transported in such containers, under such conditions, in particular with regard to space, ventilation and temperature, and with such supply of water and oxygen as are considered appropriate for the species. They shall be transported to their destination as soon as possible".

European Convention for the Protection of Animals Kept for Farming Purposes

This convention is a "framework convention", which gives principles for the keeping, care and housing of animals, in particular in intensive breeding systems. The Convention applies to animals bred or kept for the production of food, wool, skin or fur or for other farming purposes. It concerns in particular animals in intensive stock-farming systems. In this Convention no special attention is paid to the welfare of fishes. Fish farmed for food are included within the scope of the EU Directive

³ ESFA, 2004. Summary of Opinion of the Scientific Panel on Animal Health and Welfare (AHAW) on a request from the Commission related to welfare aspects of the main systems of stunning and killing the main commercial species of animals (Question number: EFSA-Q-2003-093). http://www.efsa.eu.int/science/ahaw/ahaw_opinions/495_en.html

concerning the protection of animals kept for farming purposes (Council Directive 98/58/EC). There is, however no specific EU regulation relating to the welfare of fish and the Annex to the Directive – which contains detailed provisions – is not applicable to fish. In addition the Council of Europe is developing a recommendation on the welfare of farmed fish. This recommendation is to be adopted under the Convention on the Protection of Animals Kept for Farming Purposes and, in due course, form the basis of detailed EU legislation but this is a long term prospect.

European Convention for the Protection of Animals for Slaughter

The main purpose of the Convention is to help harmonise methods of slaughter in Europe and make them more humane. The first set of provisions impose a number of obligations concerning the treatment of animals in slaughterhouses: use of suitable equipment for unloading animals; no brutalising or ill-treatment of animals, and in particular no striking of sensitive parts of their bodies; lairaging and care for animals which are not slaughtered immediately on arrival; provision of the requisite facilities at slaughterhouses. As regards the slaughter operation, the Convention stipulates that all animals must be stunned before being bled. Stunning of large animals must be effected by means of a pistol (an instrument which administers a blow or penetrates at the level of the brain), by electro-narcosis or by gas. The use of a poleaxe, hammer or puntilla is prohibited by the Convention. Furthermore, large animals must neither be suspended nor have their movements restricted before being stunned. Where there are exceptions to these rules (ritual slaughter, emergency slaughter, slaughter of poultry and rabbits, etc.), slaughter must be effected in such a way as to spare the animals any unnecessary suffering. These last provisions apply equally to slaughter in places other than slaughterhouses. This Convention only applies to the movement, restraint, stunning and slaughter of domestic solipeds, ruminants, pigs, rabbits and poultry. Therefore it does not apply to fishes.

European Convention on the Protection of Pet Animals

The Convention aims essentially at assuring the welfare of animals, and in particular, of pet animals kept for private enjoyment and companionship. Animals belonging to threatened species, and protected by other Conventions, in particular, the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES), opened for signature in Washington, on 3 March 1973, and the Convention on the Conservation of European Wildlife and Natural Habitats (ETS No. 104), opened for signature by the member States of the Council of Europe and non-member States which have participated in its elaboration and by the European Economic Community, in Bern, on 19 September 1979, are thus excluded from the provisions of this text. In this Convention no special attention is paid to the welfare of fish. Since this Convention only applies to pet animals it is beyond the scope of this report and will therefore not be discussed any further.

European Convention for the Protection of Vertebrate Animals Used for Experimentation and other Scientific Purposes

The Convention is designed primarily to reduce both the number of experiments and the number of animals used for such purposes. It encourages Parties not to experiment on animals except where there is no alternative. All research into alternative methods should be encouraged. Animals to be experimented on should be selected on the basis of clearly established quantitative criteria and must be well cared for and spared avoidable suffering whenever possible. To this end, the Convention lays down a number of principles that are to be considered only as a starting point. The Parties meet regularly to examine the application of the Convention and, if appropriate, to extend or strengthen its provisions. In Appendix A of this Convention, species specific provisions for fish are given. Here general considerations for water quality, health, housing and enrichment, feeding, handling and humane killing are provided. Specific recommendations are listed for rainbow trout, Atlantic salmon, tilapia, zebrafish, seabass, Atlantic halibut, cod, turbot and African walking catfish.

5.1.4 World Organisation for Animal Health (OIE)

In May 2002, the OIE has adopted a Resolution on Animal Welfare. The 167 member nations of the OIE have accepted to start the development of policies and guiding principles to provide a sound foundation from which to elaborate specific recommendations and standards. In developing these policies, the OIE will give priority to animal welfare issues regarding animals used in agriculture and aquaculture. In February 2004, OIE organized a global conference on animal welfare: an OIE

initiative. The proceedings of the conference are published and have contributions about pain, fear and distress (Duncan, 2004), injuries and diseases (Algers, 2004) and animal welfare issues relating to aquaculture (Håstein, 2004).
(http://europa.eu.int/comm/food/animal/welfare/international/2003_2073_31_en.pdf).

5.2 National legislation and regulation

The following overview gives the data on legislation and regulation in various countries in Europe. A part of the information was obtained by the questionnaire that was sent to the EIFAC national correspondents in March 2003. Also information was obtained from participants of the EIFAC workshop in Utrecht (24-26 March 2004) and from various sources on the internet. Chapter 6 (Legislation) in Thorsteinsson (2002), gives an overview of the very variable legal position of fish tagging within Europe.

5.2.1 Austria

Regulations concerning fish welfare can be found in the 'Fischereigesetz'. Different Austrian states each have their own 'Fischereigesetz', but they all provide some regulations concerning the protection of individual fish as well as the fish population.

5.2.2 Finland

Finnish requirements concerning animal welfare are laid down in the Act on Animal Protection (1996), Statute on Animal Protection (1996), Statute on the Transport of Animals (1996) and Statute on Animal Experimentation (1985). More detailed provisions on requirements for the keeping, care, treatment and handling of animals have been laid down in several Decisions issued by the Ministry of Agriculture and Forestry. The Decisions concern the keeping of different animal species and activities and measures involving animals.

The objective of the Finnish Act on Animal Protection is to protect animals from suffering, pain and distress in the best manner possible and to promote the welfare and good treatment of animals. The Act concerns all animals, both wild animals and animals kept by humans, such as pets, domestic animals and farmed animals. It provides general regulations concerning general provisions, the keeping of animals, the transport of animals, import of animals, experiments with animals, the slaughter of animals. The Act does not provide specific regulations on fish.

5.2.3 France

The present animal protection legislation on animal experimentation in France (the decree of 19 October 1987 and the three ministerial orders of 19 April 1988) incorporates the European Directive 86/609, while maintaining all provisions of the previous national laws. The regulations apply only to living vertebrate animals. Experiments can only be performed after written justification.

Animal experimentation can only be performed under the responsibility of an individual holding a personal licence issued by the Ministry of Agriculture. The authorisation specifies the area of activity, the animal species utilised and the experimental protocols. Applicants must have a minimum education in the relevant biology and training in animal experimentation. In addition, specific training is required for any surgical procedures. The training in animal experimentation must be approved by the Ministry of Agriculture.

The facilities where animal experiments take place must be approved jointly by the Ministry of Agriculture and the ministries with administrative responsibility over the institution conducting or funding the experimentation. These facilities must be staffed with an adequate number of qualified personnel with appropriate training in accordance with their responsibilities in category I (persons responsible for the experiment), II (personnel participating in the experiment) or III (personnel in charge of caring for and husbandry of the animals) as defined in the regulations. In addition, the facilities must maintain amongst other items, a register of the animals obtained and sent out, visits to the animal units, animal housing and controlled environments.

The approval of the facilities includes details of the type of experimentation, the animal species to be used, protocols to be applied and requires statistics on animal numbers to be collected. To maintain wild animals or animals of wild origin requires a facility certificate of the Ministry of the Environment. The animals must be obtained from identified establishments. Cats, dogs and non-human primates must be individually identified by permanent means; the identification number of cats and dogs is tattooed and registered in a national file. The control of these regulations is the responsibility of the Ministry of Agriculture's Departmental Veterinary Services. Depending on the seriousness of any offences, sanctions can be administrative or through the courts.

The Commission Nationale de l'Experimentation Animale, presided over by a state counsellor, is an advisory body to the Ministries charged with giving opinions or making proposals on all aspects of the present legislation. This commission is also consulted regarding training and the development of 'alternative' methods.

5.2.4 Germany

In Germany, legislation to protect farmed eel at slaughter came into force in April 1999 (TierSchlV, 1997; 1999).

The German Animal Welfare Act dates back to 1934: This law was revised in 1972, again in 1986, 1993, 1997, 1998, 2000 and 2003. It covers all species of animals. Since 1 August 2002, animal welfare is defined as a national objective. The aim of the German Animal Welfare Act is the overall protection of animals as creatures under human responsibility. Nobody is allowed to address any pain, suffering or damage to an animal without a sound reason. The act provides regulations for animal husbandry, killing, operations on animals, experiments on animals, animal breeding and import, movement and keeping of animals. The sections applying to experimentation cover both vertebrates and invertebrates. However, licences are only required for the use of vertebrates. Reports have to be submitted on projects using invertebrates, although the number used does not have to be recorded. No application has to be made to kill animals for in vitro experimentation or for other reasons and this does not have to be reported. However, there are plans to start to record the numbers of animals killed for such purposes. In this act no special attention is paid to the welfare of fish. Additional acts are available on transportation and slaughter of animals. Both provide specific regulations on fish.

Further regulations concerning fish welfare can be found in the fisheries legislation. The German Länder each have their own "Fischereigesetz", but they all provide some regulations concerning the protection of individual fish as well as the fish population. Typical examples for restrictions are the use of live bait fish, keep nets, fishing gear, and the practice of "put and take" and "catch and release". Angling contests are in most cases forbidden.

Since 2004, a procedure for testing the toxicity of waste water has been changed: instead of young adults of the fish species "Goldorfe" (*ide*; *Leuciscus ide* L.), now only the use of fish eggs are allowed for the tests.

5.2.5 Ireland

National legislation

The Cruelty to Animals Act 1876

EC (Amendment to The Cruelty to Animals Acts of 1976)

Animal Remedies Act 1996

www.irishstatutebook.ie/ZZSI17Y1994.html

National regulations

Regulations of 1994, Statutory Instruments number 17 of 1994.

Ministry involved

Department of Communications, Marine and Natural Resources Health

Department of Agriculture

5.2.6 *Netherlands*

On 23 September 1992 the Animal Health and Welfare Act came into force in the Netherlands. The Act is based on the 'no, unless – principle', indicating that no handling of animals is allowed unless stated otherwise. It provides general rules concerning animal health, animal welfare, transportation, slaughter and housing. Furthermore, it provides regulations on biotechnology. In this act no special attention is paid to the welfare of fish.

On 5 February 1997 a revised Experiments on Animals Act came into force in the Netherlands. The main change introduced by the revised Act is to require that research plans must be approved by an ethical review committee, which has to consider the benefit to come from the experiments and whether this justifies the distress caused to the animals to be used. No maximum duration for a research plan is stipulated. Committees are required to give a decision on an application within three months and, if they do not approve the research plan, the applicant can appeal to the central animal review committee established under the previous legislation.

The structure of these committees is defined in the new Act, which requires that they have at least seven members made up from equal numbers of experts in animal experiments, alternative methods, animal welfare, and ethical assessment. At least two members must not be conducting animal experiments and at least three members, including the chairperson, must not be employed by a scientific institution applying to the committee. In this act, no special attention is paid to the welfare of fishes.

A third Act available for regulating fish welfare is the Fisheries Decree. There is scope in the Fisheries Decree for welfare regulations concerning the treatment of fishes that are not kept. This mainly concerns the methods used to catch the fishes. In connection with this, two measures have been taken:

- Live bait-fish may not be used for fishing since 1998;
- The Minister of Agriculture, Nature and Food Quality can forbid the use of devices for stunning, wounding or killing fish.

5.2.7 *Norway*

Animal welfare in Norway is regulated by the Animal Welfare Act of 20 December 1974. This Act applies to live mammals, birds, toads, frogs, salamanders (newts), reptiles, fish, and crustaceans. It provides general regulations concerning general provision on how animals should be treated, special provision concerning livestock and domestic reindeer, showing of animals, animal boarding, accommodation, trade in and hiring out of animals and animal use in teaching and research. Animal experimentation is also regulated through the Animal Welfare Act. Furthermore it provides special provisions concerning fish and crustacean. Chapter 3 of the Animal Welfare Act states that it is forbidden to suspend live fish on a rod, line, or hook or the like, which is stuck into or through the body of the fish. Furthermore, it is forbidden to keep fish in a net-cage in such a way that they are at risk of suffering needlessly and to place live fish or crustacean in a shop window.

5.2.8 *Poland*

Animal welfare in Poland is regulated by the Polish Animal Protection Act of 1997. This Act applies to domestic animals, farm animals, animals used for the purposes of entertainment, shows, films and sports and for special purposes, animals used in experiments, animals kept in zoological gardens, free living animals (wild) and animals not belonging to the domestic fauna. It provides general regulations concerning general provision on how animals should be treated, domestic animals, farm animals, animals used for the purposes of entertainment, shows, films and sports, animals in zoological gardens, animals living at large (wild), transport of animals, operations and animals and the killing of animals. Animal experimentation is also regulated through the Animal Protection Act. No specific regulations on fish are provided in this Act.

5.2.9 Sweden

The Swedish system of regulating animal welfare comes under a general law, the Animal Protection Act and the Animal Protection Ordinance, both of 1988. The Acts apply to the care and treatment of domestic animals. Furthermore it applies to animals kept in captivity or used for scientific research or education, the diagnosis of disease, the manufacture of drugs or chemical products or other similar purposes. The Acts provide general rules concerning animal management and treatment, surgical procedures, slaughter, transportation, use of animals for scientific purposes, sporting events and exhibitions involving animals, etc. The Acts do not provide specific regulations on fishes.

All animal research projects require ethical committee approval, including feeding studies, experiments under terminal anaesthesia and the killing of animals to remove tissues for use in *in vitro* biomedical research.

5.2.10 Switzerland

The Swiss system of regulating animal welfare comes under the Act on Animal Protection (1978) and the Animal Protection Ordinance (1981). The Acts apply to vertebrates only. They provide general regulations concerning general provisions, the keeping of animals, the trade of animals and use of animals for advertising purposes, the transport of animals, surgical operations performed on live animals, experiments with animals, the slaughter of animals, etc. The Acts do not provide specific regulations on fish.

Further regulations concerning fish welfare can be found in the “Bündesgesetz über die Fischerei” (1991) and the “Verordnung zum the Bündesgesetz über die Fischerei” (1993). It provides regulations to protect caught fish against unnecessary damage and injury and it bans the use of live baitfish.

5.2.11 United Kingdom

In the United Kingdom animal welfare is covered by four types of legislation:

1. Legislation to protect animals from cruelty

- The Protection of Animals Acts 1911-1988
- The Veterinary Surgeons Act 1966
- The Abandonment of Animals Act 1960
- The Animals (Cruel Poisons) Act 1962
- The Docking and Nicking of Horses Act 1949

2. Legislation to protect the welfare of animals kept in commercial enterprises

- The Dogs Act 1973
- The Breeding of Dogs Act 1973 and 1991
- The Breeding and Sale of Dogs (Welfare) Act 1999
- The Animal Boarding Establishments Act 1963
- The Riding Establishments Acts 1964 and 1970
- The Performing Animals (regulation) Act 1925
- Agriculture (Miscellaneous Provisions) Act 1968
- The Zoo Licensing Act 1981

3. Legislation to protect society from the nuisance of animals

- The Animals Act 1971
- The Dangerous Dogs Act 1989
- The Dangerous Wild Animals Act 1976

4. Legislation to protect the welfare of animals involved in research

- Animals (Scientific Procedures) Act 1986

Considering the scope of this report, important legislation in the United Kingdom to be discussed is the Protection of Animals Act 1911, the Agriculture (Miscellaneous Provisions) Act 1968 and the Animals (Scientific Procedures) Act 1986. The main and most important offences of cruelty to captive and domestic animals are set out in the Protection of Animals Act 1911 with respect England and Wales. Similar legislation covers Scotland and slightly different legislation is in force in Northern Ireland. Although the 1911 Act has been amended on a number of occasions, its most significant language and features have remained untouched. Even though the 1911 Act is now over 90 years old and was itself considered a consolidating statute that drew on a number of nineteenth century statutes, it remains an imperfect, but nevertheless reasonably simple and effective, statute for the prosecution of persons who inflict unnecessary suffering on captive and domestic animals.

The Agriculture (Miscellaneous Provisions) Act 1968 is sometimes used in welfare cases, instead of the Protection of Animals Act 1911. On occasions, it is used to support cases brought under the 1911 Act. It is particularly useful where large numbers of animals are involved, as the offences are worded to cover more than one animal. It can only be used where the offence occurred on agricultural land. The offences provide for two levels of suffering, the more serious offence being “causing unnecessary pain” and the less serious, “causing unnecessary distress”

The British system for regulating the use of animals in experiments is defined by the Animals (Scientific Procedures) Act. The main principle of this law is the requirement for researchers to obtain personal and project licences and a certificate for the establishment where the animal experimentation is conducted. The legislation covers all vertebrate animals and has recently been extended to include the octopus.

National legislation United Kingdom

In England and Wales, the Environment Agency makes byelaws under Salmon and Freshwater Fisheries Act 1975 to control angling practice e.g. use of knotless netting for nets.

In Scotland, regulations governing fishing are administered by local fisheries boards and regulations governing the ways in which a fish can be taken are laid down in the Salmon and Freshwater Fisheries (Protection) (Scotland) Act 1951.

Legislation pertaining to the welfare of farmed fish:

- The Protection of Animals (Scotland) Act 1912 – prevents any unnecessary suffering to any domestic or captive animal.
- The Agriculture (Miscellaneous Provisions) Act 1968 – prevents unnecessary pain or unnecessary distress to any livestock (any creature kept for the production of food).
- The Welfare of Animals (Slaughter and Killing) Regulations 1995 –implements Council Directive 03/119/EC and applies to the movement, lairaging, restraint, stunning, slaughter and killing of animals bred and kept for the production of meat, skin, fur or other products.
- The Welfare of Animals (Transport) Order 1997 – this order requires that all animals, including fish, are transported in a way that does not, and is not likely to, cause injury or unnecessary suffering.
- The Welfare of Farmed Animals (Scotland) Regulations 2000 –implements Council Directive 98/58/EC and requires reasonable steps to be taken to ensure animals kept for farming purposes are not caused any unnecessary pain, suffering or injury.

National regulations

The Department for Environment, Food and Rural Affairs (DEFRA), in partnership with the Scottish Executive and the Welsh Assembly Government, are working to develop an all-inclusive and comprehensive long-term Animal Health and Welfare strategy for Great Britain. This project will be closely linked to the development of the Northern Ireland and Republic of Ireland “All Island” animal health programmes. (Currently, a new Animal Welfare Bill is progressing through the British Parliament (see <http://www.defra.gov.uk/animalh/welfare/bill>), but note that although fish are referred, to in the context of fish being kept, angling practices are excluded.)

Ministry involved

Fisheries Minister.

Departments involved

Department for Environment, Food and Rural Affairs (DEFRA), Nobel House, 17 Smith Square, London SW1P 3JR, United Kingdom. <http://www.defra.gov.uk/>

5.2.12 Denmark

The legislation on animal welfare is under the Ministry of Justice and is governed by the law for animal protection from 1991. This law applies mainly to pet and farm animals, and nothing is mentioned about fish. The use of vertebrate animals for experiments is ruled by the law on use of experimental animals from 1993. All vertebrate animal research projects that may cause pain, suffering, fear or permanent harm require approval from the ethical committee. For the use of fish in experiments, only invasive methods like surgical implantation of electronic tags require special permission, where the applicant must have attended a training course in handling of experimental animals. Furthermore veterinarian supervision is required. Regarding fish welfare in aquaculture, commercial fisheries and recreational fisheries, the only well-known rule is the prohibition of killing eels with salt or ammonia.

CHAPTER 6 CURRENT CODES AND GUIDELINES

6.1 Aquaculture

Several guidelines covering aspects of welfare of fishes in aquaculture are present today throughout the world:

- The Farm Animal Welfare Council (FAWC) in the United Kingdom has published the Report on the Welfare of Farmed Fish (FAWC, 1996). It offers guidelines about the farming of salmon (*Salmo salar*) and trout (*Oncorhynchus mykiss* and *Salmo trutta*). It also presents brief comments on carp (*Cyprinus carpio*). Furthermore, the welfare of species of wrasse used for parasite control during salmon farming (*Ctenolabrus rupestris*, *Centrolabrus exoletus* and *Crenilabrus melops*) is also discussed.
- The British Columbia (Canada) Salmon Farmers Association has also produced a Draft Code of Practice for the salmon farming industry. Goal of this code of practice is to protect the safety and well being of employees, the natural environment and salmon stocks.
- The Federation of European Aquaculture Producers (FEAP) has published a Code of Conduct for European Aquaculture, prepared to promote the responsible development and management of a viable European aquaculture sector in order to assure a high standard of quality food production while respecting environmental considerations and consumers' demands.
- The Holmenkollen Guidelines for Sustainable Aquaculture 1998 call for "standards and practices which embody ethical principles for ensuring health and welfare of fish and shellfish and for slaughter practices" It provides some general policy recommendations and a statement of the ethical responsibilities of the industry.
- FAO Technical Guidelines for Responsible Fisheries No. 6: Inland Fisheries (FAO 1997).
- FAO Technical Guidelines for Responsible Fisheries No. 5.: Aquaculture Development (FAO 1997). Specifically: contains references and recommendations regarding:
 - stress management (p.28; under CCRF Article 9.4.2)
 - humane killing of fish (p.31; under CCRF Article 9.4.6)
 - <ftp://ftp.fao.org/docrep/fao/003/W4493e/W4493e00.pdf>.

6.2 Commercial fisheries

Codes of conduct concerned with the welfare of fishes in commercial fisheries are rare. The FAO Fisheries Department has published a Code of Conduct for Responsible Fisheries (CCRF).

<http://www.fao.org/fi/agreem/codecond/ficonde.asp>

<ftp://ftp.fao.org/docrep/fao/005/v9878e/V9878E00.pdf>

The CCRF provides a specific reference and recommendation to "the use of fishing gear and practices that increase survival rates of escaping fish should be promoted." CCRF Article 8.4.5 states that: "States, with relevant groups from industry, should encourage the development and implementation of technologies and operational methods that reduce discards. The use of fishing gear and practices that lead to the discarding of catch should be discouraged and the use of fishing gear and practices that increase survival rates of escaping fish should be promoted."

6.3 Angling

Several guidelines on the welfare of fishes in angling are published:

- Eleven national member associations of Recfish Australia have published the National Code of Practice for Recreational and Sport Fishing. The Code addresses four main areas of fishing responsibility. These include looking after our fisheries, protecting the environment, treating fish humanely and respecting the rights of others.
- The National Angling Alliance United Kingdom has published a Code of Conduct for Coarse Anglers. The Code was designed to complement the existing legislative framework and provides a commonsense guide to values and behaviour to which anglers should aspire.
- Central Fisheries Board Ireland Leaflet No. 1. *Catch and Release for Atlantic Salmon*.
- Several other small angling associations have produced general, local codes.

6.4 Ornamental fish

The ornamental fish sector has instituted a code of conduct that addresses, among other things, the welfare of aquarium fish (Davenport, 1993). Based on the Dutch Legislation on Animal Health and Welfare the Council of Animal Affairs has produced a document with a “positive list” based on welfare criteria of fish taxa which can be kept in aquaria. The Dutch Ministry of Agriculture, Nature Conservation and Food Quality, however, has not adopted the list because it is not feasible to enforce the regulation.

CHAPTER 7 CITED REFERENCES

- Ainslie, B.J., Post, J.R. & Paul, A.J., 1998. Effects of pulsed and continuous DC electrofishing on juvenile rainbow trout. *North American Journal of Fisheries Management*, 18, 905-918.
- Alabaster, J. S. & Lloyd, R., 1980. *Water Quality Criteria for Freshwater Fish*. London: Butterworth & Co Ltd.
- Alanära, A. & Brännäs, E., 1996. Dominance in demand-feeding behaviour in Arctic charr and rainbow trout: the effect of stocking density. *Journal of Fish Biology*, 48, 242-254.
- Algers, B., 2004. Injury and disease. In Proceedings Global conference on animal welfare: an OIE initiative. Paris, 23-25 February 2004: 176-185.
- Anderson, W.G., Booth, R., Beddow, T.A., McKinley, R.S., Finstad, B., Økland, F. & Scruton, D., 1998. Remote monitoring of heart rate as a measure of recovery in angled Atlantic salmon, *Salmo salar* (L.). *Hydrobiologica*, 371/372, 233-240.
- Anonymous. 1979. *The welfare of the food animals: An assessment of developments in the livestock industry and recommendations for the most practical way in which animal welfare requirements can be met*. Proceedings of a Symposium, The Universities Federation for Animal Welfare, September 28-29, 1978, p. 68.
- Anonymous, 1993. Report of Activities 1992 UK Scotland. ICES Working Group on Fishing Technology and Fish Behaviour Working Group, Gothenburg, April 1993. (Unpublished).
- Anonymous, 2002. Marine Harvest wins welfare award. *Fish Farming Today*, October 2002.
- Ashton, N., Brown, N. & Easty, D., 1969. Trematode cataract in freshwater fish. *Journal of Small Animal Practice*, 10, 471-478.
- Azam K., Mackie I.M. & Smith J., 1989. The effect of slaughter method on the quality of rainbow trout (*Salmo gairdneri*) during storage on ice. *International Journal of Food Science and Technology*, 24, 69-79.
- Bandeen, J. & Leatherland, J.F., 1997. Transportation and handling stress of white suckers raised in cages. *Aquaculture International*, 5, 385-396.
- Barahona-Fernandes, M.H., 1979. Some effects of lighting intensity and photoperiod on the seabass larvae (*Dicentrarchus labrax* (L.)) reared at the Centre Oceanologique de Bretagne. *Aquaculture*, 17, 311-321.
- Barnett, C.W. & Pankhurst, N.W., 1998. The effects of common laboratory and husbandary practices on the stress response of greenback flounder (*Rhombosolea tapirina*, Gunther 1862). *Aquaculture*, 162, 313-329.
- Barton, B.A., 2000. Salmonid fishes differ in their cortisol and glucose responses to handling and transport stress. *North American Journal of Aquaculture*, 62, 12-18.
- Barwick, D.H., 1985. Stocking and hooking mortality of planted rainbow trout in Jocassee Reservoir, North Carolina. *North American Journal of Fisheries Management*, 5, 580-583.
- Berg T., Erikson U. & Nordtvedt T.S., 1997. Rigor mortis assessment of Atlantic salmon (*Salmo salar*) and effects of stress. *Journal of Food Science*, 62, 439-446.
- Berka, R., 1986. The transport of live fish – a review. *FAO Technical paper* 48.
- Bermond, B., 1997. The myth of animal suffering. In: M. Dol, S. Kasanmoentalib, S. Lijmbach, E. Rivas & R. van den Bos (eds.). *Animal consciousness and animal ethics*. (pp. 125 – 143) Assen: Van Gorcum.
- Bettoli, P.W. & Osborne, R.S., 1998. Hooking mortality and behavior of striped bass following catch and release angling. *North American Journal of Fisheries Management*, 18, 609-615.
- Beukema, J.J., 1970a. Acquired hook-avoidance in the pike *Esox lucius* L. fished with artificial and natural baits. *Journal of Fish Biology*, 2, 155-160.
- Beukema, J.J., 1970b. Angling experiments with carp (*Cyprinus carpio* L.) II. Decreasing catchability through one-trial learning. *Netherlands Journal of Zoology*, 20, 81-92.
- Bjerkås, E. & Bjørnstad, E., 1999. Is there a connection between rapid fluctuation in water temperature and cataract development in the atlantic salmon (*Salmo salar* L.)? *Bulletin of the European*

- Association of Fish Pathologist*, 19, 166-169.
- Bjerkås, E., Bjørnstad, E., Breck, O. & Waagbø, R., 2001. Water temperature regimes affect cataract development in smolting Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases*, 24, 281-291.
- Bjerkås, E., Wall, A.E., & Prapas, A., 2000. Screening of farmed seabass (*Dicentrarchus labrax* L) and sea bream (*Sparus aurata* L.) for cataract. *Bulletin of the European Association of Fish Pathologist*, 20, 180-185.
- Bolla, S. & Holmefjord, I., 1988. Effect of temperature and light on development of Atlantic halibut larvae. *Aquaculture*, 74, 355-358.
- Bosakowski, T. & Wagner, E. J., 1994. A survey of trout fin erosion, water quality and rearing conditions at state fish hatcheries in Utah. *Journal of the World Aquaculture Society* 25, 308-316.
- Boyd, C. E. , 1982. *Water Quality Management for Pond Fish Culture*. Oxford: Elsevier.
- Braithwaite, V.A., 2004. Fish Pain Perception. Workshop Fish Welfare in Aquaculture, Painter's Lodge, Campbell River, BC, September 27, 2004.
http://www.aquanet.ca/English/research/forum_welfare.php
- Braithwaite, V.A. & Huntingford, F.A., 2004. Fish and welfare: do fish have the capacity for pain perception and suffering? *Animal welfare* 13 Supplement: 87-92.
- Bromage, N, Porter, M & Randall, C., 2001. The environmental regulation of maturation in farmed finfish with special reference to the role of photoperiod and melatonin. *Aquaculture* 197, 63-98
- Broom D. M. 2001. Evolution of pain. Royal Society of Medicine International Congress and Symposium Series, 246,17-26.
- Buchanan, S., Farrell, A.P., Fraser, J., Gallagher, P., Joy, R. & Routledge, R., 2002. Reducing gill-net mortality of incidentally caught coho salmon. *North American Journal of Fisheries Management*, 22, 1270-1275.
- Byrne, H., 2002. *The welfare and quality of farmed salmonids at harvest*. Ph.D. thesis, The Robert Gordon University Aberdeen, Scotland, 269 pp.
- Caddy, J.F., 1973. Underwater observations of tracks of dredges and trawls and some effects of dredging on a scallop ground. *Journal of the Fisheries Research Board of Canada*, 30, 173-180.
- Carragher, J.F. & Pankhurst, N.W., 1991. Stress and reproduction in a commercially important marine fish, *Pagrus auratus* (Sparidae). *Reproductive Physiology of Fish*, pp. 253-255. Fish Symposium 91, Sheffield, England.
- Chandoo, K.P., Duncan, I.J.H. & Moccia R.D., 2004. Can fish suffer?: perspectives on sentience, pain, fear and stress. *Applied Animal Behaviour Science*, 86, 225-250.
- Cho, G.K., Heath, J.W. & Heath, D.D., 2002. Electroshocking influences Chinook salmon egg survival and juvenile physiology and immunology. *Transactions of the American Fisheries Society*, 131, 224-233.
- Chopin, F.S. & Arimoto, T., 1995. The condition of fish escaping from fishing gears-a review. *Fisheries Research* 21, 315-327.
- Chopin, F.S., Arimoto, T. & Inoue, Y., 1996. A comparison of the stress response and mortality of sea bream *Pagrus major* captured by hook and line and trammel net. *Fisheries-Research*, 28, 277-289.
- Clapp, D.F. & Clark Jr., R.D., 1989. Hooking mortality of smallmouth bass caught on live minnows and artificial spinners. *North American Journal of Fisheries Management*, 9, 81-85.
- Clearwater, S.J. & Pankhurst, N.W., 1997. The response to capture and confinement stress of plasma cortisol, plasma sex steroids and vitellogenic oocytes in the marine teleost, red gurnard. *Journal of Biology*, 50, 429-441.
- Cooke, S.J. & Cowx, I.G., 2006. Contrasting recreational and commercial fishing: searching for common issues to promote unified conservation of fisheries resources and aquatic environments. *Biological Conservation*, 128, 93-108.
- Cooke, S.J. & Hogle, W.J., 2000. Effects of retention gear on the injury and short-term mortality of adult smallmouth bass. *North American Journal of Fisheries Management*, 20, 1033-1039.
- Cooke, S.J., Philipp, D.P., Dunmall, K.M. & Schreer, J.F., 2001. The influence of terminal tackle on injury, handling time,

- and cardiac disturbance of rock bass. *North American Journal of Fisheries Management*, 21, 333-342.
- Cowx, I & Lamarque, P. (1990) *Fishing with electricity*, Fishing News Books, Oxford.
- Cowx, I.G. & Gerdeaux, D., 2004. The effects of fisheries management practices on freshwater ecosystems *Fisheries Management and Ecology*, 11, 145-152.
- Cullen, A.P., Monteith-McMaster, C.A. & Sivak, J.G., 1994. Lenticular changes in rainbow trout following chronic exposure to UV radiation. *Current Eye Research*, 13, 731-737.
- Davenport, K., 1993. OFI (UK) code of conduct: the way ahead. *Aquarist and Pondkeeper* (February 1993): 17-18.
- Davenport, J., Baras, E., Fabi, G. & Jonsson, G., 2002. Chapters 6 (Legislation) & 7 (Fish welfare and health in relation to tagging) In: Thorsteinsson V. (coord.) 2002. *Tagging Methods for Stock Assessment and Research in Fisheries. Report of Concerted Action FAIR CT.96.1394 (CATAG)*. Reykjavik. Marine Research Institute Technical Report No. 79. pp 183.
- Davidson, G.W., Thorarensen, H.T., Lokman, M. & Davie, P.S., 1997. Stress of capture and captivity in kahawai *Arripis trutta* (Bloch and Schneider) (Perciformes: Arripidae). *Comparative Biochemistry and Physiology*, 118A, 1405-1410.
- Davis, K.B. & Parker, N.C., 1990. Physiological stress in striped bass: Effect of acclimation temperature. *Aquaculture*, 91, 349-358.
- DeAlteris, J.T. & Reifsteck, D.M., 1993. Escapement and survival of fish from the codend of a demersal trawl. ICES Marine Science Symposium, 11-13 June 1992, Bergen. ICES Secretariat, Copenhagen, 196, 128-131.
- Dieterman, D.J., Baird, M.S. & Galat, D.L., 2000. Mortality of paddlefish in hoop nets in the lower Missouri river, Missouri. *North American Journal of Fisheries Management*, 20, 226-230.
- Dolan, C.R., Miranda, L.E. & Henry, T.B., 2002. Electrofishing for crappies: electrical settings influence immobilization efficiency, injury, and mortality. *North American Journal of Fisheries Management*, 22, 1442-1451.
- Donaldson, E. M. , 1981. The pituitary-interrenal axis as an indicator of stress in fish. In *Stress and Fish* (Pickering, A. D., ed.), pp. 11-47. London: Academic Press.
- Dotson, T., 1982. Mortalities in trout caused by gear type and angler-induced stress. *North American Journal of Fisheries Management*, 2, 60-65.
- Downing, G. & Litvak, M.K., 1999. The influence of light intensity on growth of larval haddock. *North American Journal of Aquaculture*, 61, 135-140.
- Duncan, I.J.H., 2004. Pain, fear and distress. In Proceedings Global conference on animal welfare: an OIE initiative. Paris, 23-25 February 2004: 160-169.
- Duncan, I.J.H. & Fraser, D., 1997. Understanding animal welfare. *Animal Welfare* (ed. by M.C. Appleby & B.O. Hughes), pp. 19-31. CAB International. University Press, Cambridge.
- Dunning, D.J., Ross, Q.E., Mattson, M.T. & Geoghegan, P., 1989. Handling and capture mortality in trawls and seines. *North American Journal of Fisheries Management*, 9, 171-176.
- Duray, M. & Kohno, H., 1988. Effects of continuous lighting on growth and survival of first-feeding larval rabbitfish *Siganus guttatus*. *Aquaculture*, 72, 73-79.
- Dwyer, W.P., Shepard, B.B. & White, R.G., 2001. Effect of backpack electroshock on westslope cutthroat trout injury and growth 110 and 250 days posttreatment. *North American Journal of Fisheries Management*, 21, 646-650.
- Ehrensing, R.H., Michell, G.F. & Kastin, A.J., 1982. Similar antagonism of morphine analgesia by MIF-1 and naloxone in *Carassius auratus*. *Pharmacology Biochemistry and Behavior*, 17, 757-761.
- Einen, O., 2001. Live-chilling and crowding stress before slaughter of Atlantic salmon (*Salmo salar*). *Aquaculture*, 192, 265-280.
- Ejike, C. & Schreck, C.B., 1980. Stress and social hierarchy rank in coho salmon. *Transactions of the American Fisheries Society*, 109, 423-426.
- Ellis, T., North, B., Scott, A.P., Bromage, N.R., Porter, M. & Gadd, D., 2002. The relationships between stocking density and welfare in farmed rainbow trout. *Journal of Fish Biology* 61: 493-531.
- Erickson, H.S., 2003. *Information Resources on Fish Welfare: 1970-2003*. AWIC

- Resource Series No. 20. U.S. Dept. of Agriculture, National Agricultural Library, Animal Welfare Information Center. Beltsville, MD. 436 pp.
- Erikson, U., Sigholt, T. & Seland, A., 1997. Handling stress and water quality during live transportation and slaughter of Atlantic salmon (*Salmo salar*). *Aquaculture*, 149, 243-252.
- Falk, M.R., Gillmann, D.V. & Dahlke, L.W., 1974. Comparison of mortality between barbed and barbless hooked lake trout. Technical Report Series CEN/T-74-1. Department of Environmental Fisheries and Marine Services, Winnipeg, Manitoba.
- Ferguson, R.A. & Tufts, B.L., 1992. Physiological effects of brief air exposure in exhaustively exercised rainbow trout (*Oncorhynchus mykiss*): implications for "catch and release" fisheries. *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 1157-1162.
- Fraser, D., Weary, D.M., Pajor, E.A. & Milligan, B.N., 1997. A scientific conception of animal welfare that reflects ethical concerns. *Animal Welfare*, 6, 187-205.
- Fraser, P.J., Duncan, G. & Tomlinson, J., 1990. Nuvar and cataracts in Atlantic salmon (*Salmo salar*). *Experimental Eye Research*, 50, 443-447.
- Fritz, K.R. & Johnson, D.L., 1987. Survival of freshwater drums released from Lake Erie commercial shore seines. *North American Journal of Fisheries Management*, 7, 293-298.
- FSBI, 2002. *Fish Welfare*. Briefing Paper 2. Fisheries Society of the British Isles, Granta Information Systems, 82A High Street, Sawston, Cambridge CB2 4H, 25 pp.
- Gallman, E.A., Isely, J.J., Tomasso, J.R. & Smith, T.I.J., 1999. Short-term physiological responses of wild and hatchery-produced red drum during angling. *North American Journal of Fisheries Management*, 19, 833-836.
- Garcia, L.M.B., Hilomen-Garcia, G.V. & Emata, A.C., 2000. Survival of captive milkfish *Chanos chanos* Forsskal broodstock subjected to handling and transport. *Aquaculture Research*, 31, 575-583.
- Garcia-Garbi, N., Inglis, V. & Turnbull, J.F., 1998. Assessment of phagocytosis by fluorochrome acridine orange. *Methodology in Fish Disease Research*, 91-96. Ed. A. Barnes. Fish Research Services, Aberdeen, Scotland.
- Greaves, K. & Tuene, S., 2001. The form and context of aggressive behaviour in farmed Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 193, 139-147.
- Greaves, K., 2001. Manipulating aggression among juvenile Atlantic halibut (*Hippoglossus hippoglossus*) in culture conditions. PhD. Thesis. University of Glasgow.
- Grutter, A.S. & Pankhurst, N.W., 2000. The effects of capture, handling, confinement and ectoparasite load on plasma levels of cortisol, glucose and lactate in the coral reef fish *Hemigymnus melapterus*. *Journal of Fish Biology*, 57, 391-401.
- Gustaveson, A.W., Wydoski R.S. & Wedemeyer, G.A., 1991. Physiological response of largemouth bass to angling stress. *Transactions of the American Fisheries Society*, 120, 629-636.
- Habera, J.W., Strange, R.J. & Saxton, A.M., 1999. AC electrofishing injury of large brown trout in low-conductivity streams. *North American Journal of Fisheries Management*, 19, 120-126.
- Hargis, W.J., 1991. Disorders of the eye in finfish. *Annual Reviews of Fish Diseases*, 1, 95-117.
- Hartley, R.A. & Moring, J.R., 1991. Initial and delayed mortality of largemouth bass and smallmouth basses due to tournaments. In: *Warmwater Fisheries Symposium 1*, pp 269-272. Gen. Tech. Rep. RM-207. Fort Collins, CO: U.S. Department of Agriculture, Forestry Service.
- Hartley, R.A. & Moring, J.R., 1993. Observations of black bass (Centrarchidae) confined during angling tournaments: A cautionary note concerning dissolved oxygen. *Aquaculture Fisheries Management*, 24, 575-579.
- Håstein, T., 2004. Animal welfare issues relating to aquaculture. In proceedings Global conference on animal welfare: an OIE initiative: 212-220.
- Hay, D.E., Cooke, K.D. & Gissing, C.V., 1986. Experimental studies of Pacific

- herring gillnets. *Fisheries Research*, 4, 191-211.
- Hegen, H.E., Saul, G.E. & Matlock, G.C., 1987. Survival of hook-caught spotted seatrout. *Proc. 38th Annu. Conf. Southeast. Assoc. Fish Wildl. Agencies*, 38, 488-494.
- Hopkins, T. E. & Cech, J.J., 1992. Physiological effects of capturing striped bass in gill nets and fyke traps *Transactions American Fisheries Society* 121, 819-822.
- Hulbert, P.J. & Engstrom-Heg, R.E., 1980. Hooking mortality of worm-caught hatchery brown trout. *New York Fish and Game Journal*, 27, 1-10.
- Hunsaker, D., Marnell, L.F. & Sharpe, P., 1970. Hooking mortality of Yellowstone cutthroat trout. *The Progressive Fish Culturist*, 32, 231-235 .
- Israeli-Weinstein, D. & Kimmel, E., 1998. Behavioral response of carp (*Cyprinus carpio*) to ammonia stress. *Aquaculture*, 165, 81-93.
- Iversen, M., Finstad B. & Nilssen, K.J., 1998. Recovery from loading and transport stress in Atlantic salmon (*Salmo salar* L.) smolts. *Aquaculture*, 168, 387-394.
- Jepsen, N. & Aarestrup, K., 1999. A comparison of the growth of radio-tagged and dye-marked pike. *Journal of Fish Biology* 55, 880-883.
- Jepsen, N. & Berg, S., 2002. The use of winter refuges by roach tagged with miniature radio transmitters. *Hydrobiologia* 483, 167-173.
- Jørgensen, E.H., Christiansen, J.S. & Jobling, M., 1993. Effects of stocking density on food intake, growth performance and oxygen consumption in Arctic charr (*Salvelinus alpinus*). *Aquaculture* 110, 191-204.
- Juell, J-E & Fosseidengen, J.E., 2004. Use of artificial light to control swimming depth and fish density of Atlantic salmon (*Salmo salar*) in production cages. *Aquaculture* 223, 269-282.
- Kaimmer, S.M. & Trumble, R.J., 1998. Injury, condition, and mortality of Pacific halibut bycatch following careful release by Pacific cod and sablefish longline fisheries. *Fisheries research*, 38, 131-144.
- Keefe, M. & Whitesel, T.A., 2000. Induced mortality and sublethal injuries in embryonic brook trout from pulsed DC electroshocking. *North American Journal of Fisheries Management*, 20, 320-327.
- Kestin, S.C., Wotton, S.B. & Adams, S., 1995. The effect of CO₂, concussion or electrical stunning of rainbow trout on fish welfare. In: *Quality in aquaculture* (eds. K. Scheire, L. Apselagh and H. Jonkeere). Int. Conf. Aquaculture Eur. '95, Trondheim, Norway, pp 380-381.
- Kincaid, H.L., 1989. Nuclear cataract frequency in lake trout recovered from Lake Ontario in 1988. Atlantic Salmon Workshop, Atlantic Salmon Federation Research Center, 18-28.
- Laidley, C.W. & Leatherland, J.F., 1988. Cohort sampling, anaesthesia and stocking-density effects on plasma cortisol, thyroid hormone, metabolite and ion levels in rainbow trout, *Salmo gairdneri* Richardson. *Journal of Fish Biology*, 33, 73-88.
- Lambooij, E., Van de Vis, J.W., Kuhlmann, H., Münkner, W., Oehlenschläger, J., Kloosterboer, R.J., & Pieterse, C., 2002. A feasible method for humane slaughter of eel (*Anguilla anguilla* L.): electrical stunning in fresh water prior to gutting. *Aquaculture Research*, 33, 643-652.
- Lockwood, S.J., Pawson, M.G. & Eaton, D.R., 1983. The effects of crowding on mackerel (*Scomber scombrus* L.) – physical condition and mortality. *Fisheries Research*, 2, 129-147.
- Loftus, A.J., Taylor, W.W. & Keller, M., 1988. An evaluation of lake trout (*Salvelinus namaycush*) hooking mortality in the upper Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 45, 1473-1479.
- Lowe, T.E. & Wells, R.M.G., 1996. Primary and secondary stress responses to line capture in the blue mao mao. *Journal of Fish Biology*, 49, 287-300.
- Main, J. & Sangster, G.I., 1990. An assesment of the scale damage to and survival rates of young gadoid fish escaping from the cod-end of a demersal trawl. *Scottish Fisheries Research Report* No. 46.
- Matlock, G.C., McEachron, L.W., Dailey, J.A., Unger, P.A. & Chai, P., 1993. Spotted seatrout caught on single-barb and treble hooks. *North American Journal of Fisheries Management*, 13, 186-189.
- Mazur, C.F. & Iwama, G.K., 1993. Handling and crowding stress reduces number of

- plaque-forming cells in Atlantic salmon. *Animal Health*, 5, 98-101.
- Mazur, C.F. & Iwama, G.K., 1993. Effect of handling and stocking density on hematocrit, plasma cortisol and survival in wild and hatchery-reared chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture*, 112, 291-299.
- McLoughlin, R.J., Young, P.C. & Parslow, J., 1991. The Australian scallop dredge: estimates of catching efficiency and associated indirect fishing mortality. *Fisheries Research*, 11, 1-24.
- Meade, J. W., 1985. Allowable ammonia for fish culture. *Progressive Fish-Culturist* 47, 135-145.
- Milne, D.J. & Ball, E.A.R., 1956. The mortality of small salmon when caught by trolling and tagged or released untagged. *Fish. Res. Board Can. Prog. Rep.*, 106, 10-13.
- Moller, D., 2001. Process for slaughtering fish. *European Patent Specification WO 2000/038530 (06.07.2000 Gazette 2000/27)* 11 p.
- Montero, D., Izquierdo, M.S., Tort, L., Robaina, L. & Vergara, J.M., 1999. High stocking density produces crowding stress altering some physiological and biochemical parameters in gilthead seabream, *Sparus aurata*, juveniles. *Fish Physiology and Biochemistry*, 20, 53-60.
- Morzel M., Sohier D. & Van de Vis J.W., 2003. Evaluation of slaughtering methods of turbot with respect to animal protection and flesh quality. *Journal of the Science of Food and Agriculture*, 82, 19-28.
- Morzel, M. & Van de Vis, J.W., 2003. Effect of the slaughter method on the quality of raw and smoked eels (*Anguilla anguilla* L.). *Aquaculture Research*, 34, 1-11.
- Nelson, J.D., Waiwood, K.G. & Smith, S.J., 1989. Survival of Atlantic halibut (*Hippoglossus hippoglossus*) caught by longline and otter trawl gear. *Canadian Journal of Fisheries and Aquatic Sciences*, 46, 887-897.
- Nelson, K.L., 1998. Catch-and-release mortality of striped bass in the Roanoke River, North Carolina. *North American Journal of Fisheries Management*, 18, 25-30.
- Nickum, J.G., H.L. Bart, P.R. Bowser, I.E. Greer, C. Hubbs, J.A. Jenkins, J.R. MacMillan, J.W. Rachlin, J.D. Rose, P.W. Sorensen & J.R. Tomasso, 2004. Guidelines for the use of fishes in research. American Fisheries Society; American Institute of fisheries research biologists; American Society of Ichthyologists and Herpetologists.
- North, B.P., Turnbull, J.F., Ellis, T., Porter, M.J., Migaud, H., Bron, J. & Bromage, N.R., *Aquaculture* in press. The impact of stocking density on the welfare of rainbow trout (*Oncorhynchus mykiss*).
- Nuhfer, A.J. & Alexander, G.R., 1992. Hooking mortality of trophy-sized wild brook trout caught on artificial lures. *North American Journal of Fisheries Management*, 12, 634-644.
- Olla, B.L., Davis, M.W. & Schreck, C.B., 1992. Comparison of predator avoidance capabilities with corticosteroid levels induced by stress in juvenile coho salmon. *Transactions of the American Fisheries Society*, 121, 544-547.
- Olla, B.L., Davis, M.W. & Schreck, C.B., 1997. Effects of simulated trawling on sablefish and walleye Pollock: the role of light intensity, net velocity and towing duration. *Journal of Fish Biology*, 50, 1181-1194.
- Ortega, C., Muzquiz, J. L., Fernandez, A., Ruiz, I., de Blas, I., Simon, M. C. & Alonso, J. L., 1996. Water quality parameters associated with *Aeromonas* spp.-affected hatcheries. *Veterinary Research* 27, 553-560.
- Ortuño, J., Esteban, M.A. & Meseguer, J., 2001. Effects of short-term crowding stress on the gilthead seabream (*Sparus aurata* L.) innate immune response. *Fish & Shellfish Immunology*, 11, 187-197.
- Ott Jr., R.A. & Storey, K.W., 1993. Channel catfish hooking mortality. *Proc. 45th Annu. Conf. Southeast Assoc. Fish Wildl. Agencies*, 45, 399-406.
- Øverli, Ø., Olsen, R.E., Løvik, F. & Ringø, E., 1999. Dominance hierarchies in arctic charr, *Salvelinus alpinus* L.: differential cortisol profiles of dominant and subordinate individuals after handling stress. *Aquaculture Research*, 30, 259-264.
- Pálsson, O.K., Einarsson, H.A. & Björnsson, H., 2003. Survival experiments of undersized cod in a hand-line fishery at Iceland. *Fisheries Research*, 61, 73-86.

- Pankhurst, N.W. & Dedual, M., 1994. Effects of capture and recovery on plasma levels of cortisol, lactate and gonadal steroids in a natural population of rainbow trout. *Journal of Fish Biology* 45, 1013-1025 .
- Parker, R.R., Black, E.C., Black & Larkin, P.A., 1959. Fatigue and mortality in troll-caught Pacific salmon (*Oncorhynchus*). *Journal of the Fisheries Research Board of Canada*, 16, 429-448.
- Paterson, B.D., Rimmer, M.A., Meikle, G.M. & Semmens, G.L., 2003. Physiological responses of the Asian seabass *Lates calcarifer* to water quality deterioration during simulated live transports: acidosis, red-cell swelling, and levels of ions and ammonia in the plasma. *Aquaculture*, 218, 717-728.
- Payer, R.D., Pierce, R.B. & Pereira, D.L., 1989. Hooking mortality of walleyes caught on live and artificial baits. *North American Journal of Fisheries Management*, 9, 188-192.
- Peters, G. Klinger, H. & Delventhal, H. 1980. Diagnosis of stress in the European eel. Paper presented at the ICES/Special Meeting on diseases of commercially important marine fish and shellfish, no 27
- Peters, G., Delventhal, H. & Klinger, H., 1980. Physiological and morphological effects of social stress in the eel (*Anguilla anguilla* L.). *Archives Fischerei Wissenschaft*, 30, 157-180.
- Peters, G., Faisal, M., Lang, T. & Ahmed, I., 1988. Stress caused by social interactions and its effect on susceptibility to *Aeromonas hydrophila* infection in rainbow trout *Salmo gairdneri*. *Diseases of Aquatic Organisms*, 4, 83-89.
- Pickering, A.D. & Stewart, A., 1984. Acclimation of the interrenal tissue of the brown trout, *Salmo trutta* L., to chronic crowding stress. *Journal of Fish Biology*, 24, 731-740.
- Pickering, A. D., Pottinger, T. G., Sumpter, J. P., Carragher, J. F. & Le Bail, P. Y., 1991. Effects of acute and chronic stress on the levels of circulating growth hormone in the rainbow trout, *Oncorhynchus mykiss*. *General and Comparative Endocrinology* 83, 86-93.
- Pickering, A.D., 1998. Stress responses of farmed fish. In *Biology of Farmed Fish*. (Black, K.D., Pickering, A.D., eds), Sheffield Academy Press, Sheffield, 222-225.
- Plumb, J.A. 1994. *Health Maintenance of Cultured Fishes: Principal Microbial Diseases*. CRC Press, Inc.
- Plumb, J.A., Grizzle, J.M. & Rodgers, W.A., 1988. Survival of caught and released largemouth bass after containment in live wells. *North American Journal of Fisheries Management*, 8, 325-328.
- Poli, B.M., Zampacavello, G., Lurzan, F., de Francesco, M., Mosconi, G. & Parisi, G., 2002. Biochemical stress indicators changes in seabass as influenced by slaughter methods. Pp 429-430. In: *Seafarming Today and Tomorrow*. Compiled: B. Basurco & M. Saroglia. *European Aquaculture Society Special Publication* 32.
- Pottinger, T.G. & Pickering, A.D., 1992. The influence of social interaction on the acclimation of rainbow trout, *Oncorhynchus mykiss* (Walbaum) to chronic stress. *Journal of Fish Biology*, 41, 435-447.
- Pottinger, T.G. 1997. *Effects of retention of fish in keepnets*. Ambleside: Institute of Freshwater Ecology, Environment Agency, 36p.
- Pottinger, T.G., 1997. Changes in water quality within anglers' keepnets during the confinement of fish. *Fisheries Management and Ecology*, 4, 341-354.
- Pottinger, T.G., 1998. Changes in blood cortisol, glucose and lactate in carp retained in anglers' keepnets. *Journal of Fish Biology*, 53, 728-742.
- Procarione, L.S., Barry, T.B. & Malison, J.A., 1999. Effects of high rearing densities and loading rates on the growth and stress responses of juvenile rainbow trout. *North American Journal of Aquaculture*, 61, 91-96.
- Raat, A.J.P., 1985. Analysis of angling vulnerability of common carp, *Cyprinus carpio* L., in catch-and-release angling in ponds. *Aquaculture and Fisheries Management*, 16, 171-187.
- Raat, A.J.P., Klein Breteler, J.G.P. & Jansen, S.A.W., 1997. Effects on growth and survival of retention of rod-caught cyprinids in large keepnets. *Fisheries Management and Ecology*, 4, 355-368.

- Raat, A.J.P. & Van den Bos, R. 1999 (eds.) *Welzijn van Vissen*. Tilburg University Press, Tilburh, 211 pp.
- Roach, S.M., 1999. Influence of electrofishing on the mortality of arctic grayling eggs. *North American Journal of Fisheries Management*, 19, 923-929.
- Robb D.H.F., Kestin S.C. & Warriss P.D., 2000. Muscle activity at slaughter: I. Changes in flesh colour and gaping in rainbow trout. *Aquaculture*, 182, 261-269.
- Robb D.H.F., Wotton S.B., McKinstry J.L., Sørensen N.K. & Kestin S.C., 2000. Commercial slaughter methods used on Atlantic salmon: determination of the onset of brain failure by electroencephalography. *Veterinary Record*, 147, 298-303.
- Robb, D.F.H. & Kestin, S.C., 2002. Methods used to kill fish: field observations and literature reviewed. *Animal Welfare*, 11, 269-282.
- Robb, D.F.H. & Roth, B., 2002. Brain activity of Atlantic salmon (*Salmo salar*) following electrical stunning using various field strengths and pulse durations. *Aquaculture*, 216, 363-369.
- Robertson, L., Thomas, P. & Arnold, C.R., 1988. Plasma cortisol and secondary stress responses of cultured red drum (*Sciaenops ocellatus*) to several transportation procedures. *Aquaculture*, 68, 115-130.
- Rose, J.D., 2002. The Neurobehavioral Nature of Fishes and the Question of Awareness and Pain. *Reviews in Fisheries Science*, 10, 1-38.
- Rose, J.D., 2003. A critique of the paper: "Do fish have nociceptors: Evidence for the evolution of a vertebrate sensory system" published in Proceedings of the Royal Society, 2003 by Sneddon, Braithwaite and Gentle. Web site: http://uwadmnweb.uwyo.edu/Zoology/faculty/Rose/recent_publications.htm.
- Rosenthal, H., Hansen, P. D., Peters, G. & Hofmann, R. , 1984. Water quality in relation to simulated stocking density and its effect on rainbow trout. ICES-CM-1984/F:19, 13 pp.
- Roth, B., Moeller, D., Veland, J.O. & Slinde, E., 2002. The effects of stunning methods on rigor mortis and texture properties of Atlantic salmon (*Salmo salar*). *Journal of Food Science*, 67, 1462-1466.
- Rotllant, J. & Tort, L., 1997. Cortisol and glucose responses after acute stress by net handling in the sparid red porgy previously subjected to crowding stress. *Journal of Fish Biology*, 51, 21-28.
- Rouger, Y., Aubin, J., Breton, B., Fauconneau, B., Fosticr, A., Le Bail, P., Loir, M., Prunet, P. & Maisse, G. 1998. Response of rainbow trout (*Oncorhynchus mykiss*) to transport stress. *Bulletin Français de la Pêche et de la Pisciculture*, 350-51, 511-519.
- Ruane, N.M., 2002. Stress and stress disorders in a teleost fish, the common carp *Cyprinus carpio* L. PhD thesis, Fish Culture & Fisheries Group, Wageningen Institute for Animal Sciences, Wageningen University, The Netherlands.
- Sandodden, R., Finstad, B. & Iversen, M., 2001. Transport stress in Atlantic salmon (*Salmo salar*): anaesthesia and recovery. *Aquaculture Research*, 32, 87-90.
- Sangster, G.I. & Lehmann, K., 1993. Assesment of the survival of fish escaping from commercial fishing gears. ICES Fish Capture Committee CM 1993, B: 2, 10 pp.
- Sangster, G.I., Lehmann, K., Breen, M., 1996. Commercial fishing experiments to assess the survival of haddock and whiting after escape from four sizes of diamond mesh cod-ends. *Fisheries Research*, 25, 323-345.
- Scapigliati, G., Scalia, D., Marras, A., Meloni, S. & Mazzini, M., 1999. Immunoglobulin levels in the teleost seabass *Dicentrarchus labrax* (L.) in relation to age, season, and water oxygenation. *Aquaculture*, 174, 207-212.
- Schill, D.J. & Elle, F.S., 2000. Healing of electroshock-induced hemorrhages in hatchery rainbow trout. *North American Journal of Fisheries Management*, 20, 730-736.
- Schill, D.J., Griffith, J.S. & Gresswell, R.E., 1986. Hooking mortality of cutthroat trout in a catch and release segment of the Yellowstone river, Yellowstone National Park. *North American Journal of Fisheries Management*, 6, 226-232.
- Schreck, C.B., 1990. Physiological, behavioral, and performance indicators of stress. *American Fisheries Society Symposium*, 8, 29-37.
- Schreck, C.B., Contreras-Sanchez, W. & Fitzpatrick, M.S., 2001. Effects of stress

- on fish reproduction, gamete quality and progeny. *Aquaculture*, 197, 3-24.
- Schreck, C.B., Jonsson, L., Feist, G., Reno, P., 1995. Conditioning improves performance of juvenile chinook salmon, *Oncorhynchus tshawytscha*, to transportation stress. *Aquaculture*, 135, 99-110.
- Schreer, J.F., Cooke, S.J. & McKinley, R.S., 2001. Cardiac response to variable forced exercise at different temperatures: An angling simulation for smallmouth bass. *Transactions of the American Fisheries Society*, 130, 783-795.
- Schwalme, K. & Mackay, W.C., 1985. The influence of angling-induced exercise on the carbohydrate metabolism of northern pike (*Esox lucius*). *Journal of Comparative Physiology B*, 156, 67-75.
- Shepherd, J. & Bromage, N. R., 1988. *Intensive Fish Farming*. Oxford: BSP Professional Books.
- Sherwin, C., 2001. Can invertebrates suffer? Or, how robust is argument-by analogy? *Animal Welfare*, 10, S103-118.
- Shingles, A., McKenzie, D.J., Taylor, E.W., Moretti, A., Butler, P.J., Ceradini, S., 2001. The effects of sublethal ammonia on swimming performance in rainbow trout, *Oncorhynchus mykiss*. *Journal of Experimental Biology*. 204: 2691-2698.
- Sigurgisladottir S., Sigurdardottir M.S., Ingvarsdottir H., Torrisen O.J. & Hafsteinsson H., 2001. Microstructure and texture of fresh and smoked Atlantic salmon, *Salmo salar* L., fillets from fish reared and slaughtered under different conditions. *Aquaculture Research* 32, 1-10.
- Skjervold, P.O., Fjæra, S.V., Østby, P.B. & Einen, O., 2001. Live-chilling and crowding stress before slaughter of Atlantic salmon (*Salmo salar*). *Aquaculture*, 192, 265-280.
- Sloman, K.A., Motherwell, G., O'Connor, K.I. & Taylor, A.C., 2000. The effects of social stress on the Standard Metabolic Rate (SMR) of brown trout, *Salmo trutta*. *Fish Physiology and Biochemistry*, 23, 49-53.
- Smith, E.M. & Howell, P.T., 1987. The effects of bottom trawling on American lobsters *Homarus americanus*, in Long Island South. *Fisheries Bulletin*, 85, 737-744.
- Sneddon, L.U., 2003. The evidence for pain in fish: the use of morphine as an analgesic. *Applied Animal Behaviour Science*, 83, 153-162.
- Sneddon, L.U., Braithwaite, V.A. & Gentle, M.J., 2003. Do fishes have nociceptors? Evidence for the evolution of a vertebrate sensory system. *Proceedings: Biological Sciences (The Royal Society)* 270, 1520, 1115-1121.
- Snieszko, S.F., 1974. The effects of environmental stress on outbreaks of infectious diseases of fishes. *Journal of Fish Biology*, 6, 197-208.
- Soderberg, R. W., 1985. Histopathology of rainbow trout, *Salmo gairdneri* Richardson, exposed to diurnally fluctuating un-ionized ammonia levels in static-water ponds. *Journal of Fish Diseases* 8, 57-64.
- Soderberg, R. W., Flynn, J. B. & Schmittou, H. R. , 1983. Effects of ammonia on growth and survival of rainbow trout in intensive static-water culture. *Transactions of the American Fisheries Society* 112, 448-451.
- Soderberg, R. W., McGee, M. V., Grizzle, J. M. & Boyd, C. E., 1984. Comparative histology of rainbow trout and channel catfish grown in intensive static water aquaculture. *Progressive Fish-Culturist* 46, 195-200.
- Soldal, A.V. & Isaksen, B., 1993. Survival of cod (*Gadus gadus*) and haddock (*Melanogrammus aeglefinus*) escaping from a Danish seine at the sea surface. ICES Fishing Technology and Fish Behaviour Working Group, Gothenburg, April 1993.
- Soldal, A.V., Isaksen, B., Marteinson, E.J. & Engås, A., 1991. Scale damage and survival of cod and haddock escaping from a demersal trawl. ICES Fish Capture Committee C.M. 1991/B:44 12 pp.
- Spooler, H., De Rosa, G., Hörning, B., Wailblinger, S. & Wemelsfelder, F., 2003. Integrating parameters to assess on-farm welfare. *Animal Welfare* 12: 529-534.
- Spruijt, B.M., 1999. Hebben vissen een gevoel? Een gevoelig onderwerp! [Have fishes a feeling? A sensitive subject!]. Chapter 16 in: Raat & Van den Bos (eds). *Welzijn van Vissen*: 91-99.
- Stangeland K., Hoie, S. & Taksdal, T., 1996. Experimental induction of infectious pancreatic necrosis in Atlantic salmon

- (*Salmo salar* L.) postsmolts. *Journal of Fish Diseases*, 19, 323-327.
- Stevens, B.G., 1990. Survival of King and Tanner crabs captured by commercial sole trawlers. *Fisheries Bulletin*, 88, 731-744.
- Suuronen, P., Erickson, D.L. & Orrensalo, A., 1996. Mortality of herring escaping from pelagic trawl codends. *Fisheries Research*, 25, 305-321.
- Suuronen, P., Lehtonen, E., Tschernij, V. & Orrensalo, A., 1993. Survival of Baltic herring (*Clupea harengus* L.) escaping from a trawl codend and through a rigid sorting grid. ICES C.M. 1993/B:14.
- Swift, D. J. , 1981. Changes in selected blood component concentrations of rainbow trout, *Salmo gairdneri* Richardson, exposed to hypoxia or sublethal concentrations of phenol or ammonia. *Journal of Fish Biology* 19, 45-61.
- Symons, P.E.K., 1970. The possible role of social and territorial behaviour of Atlantic salmon parr in the production of smolts. *Tech. Rep. Fish. Res. Board Can.* 206, 1-25.
- Tandler, A. & Helps, S., 1985. The effects of photoperiod and water exchange rate on growth and survival of gilthead sea bream (*Sparus aurata*, Linnaeus: Sparidae) from hatching to metamorphosis in mass rearing systems. *Aquaculture*, 48, 71-82.
- Thompson, R.B. & Hunter, C.J., 1973. Viability of adult sockeye salmon that disentangle from gillnets. *International North Pacific Fisheries Commission Annual Report*, 1971, 107-109.
- Thompson, R.B., Hunter, C.J. & Patten, B.G., 1971. Studies of live and dead salmon that unmesh from gillnets. *International North Pacific Fisheries Commission Annual Report*, 1969, 108-112.
- Thorstad, E.B., Næsje, T.F., Fiske, P. & Finstad, B., 2003. Effects of hook and release on Atlantic salmon in the River Alta, northern Norway. *Fisheries Research*, 60, 293-307.
- Thorsteinsson, V. 2002. Tagging methods for stock assessment and research in Fisheries. *Report of a Concerted Action FAIR CT.96.1394 (CATAG)*. Reykjavik. Marine Research Institute Technical Report No. 79. pp 183.
- Titus, R.G. & Vanicek, C.D., 1988. Comparative hooking mortality of lure caught Lahontan cutthroat trout at Heenan Lake, California. *California Fish and Game*, 74, 218-225.
- Tort, L., Sunyer, J.O., Gómez, E. & Molinero, A., 1996. Crowding stress induces changes in serum haemolytic and agglutinating activity in the gilthead sea bream *Sparus aurata*. *Veterinary Immunology and Immunopathology*, 51, 179-188.
- Turnbull, J., Bell, A., Adams, C., Bron, J. & Huntingford, F., 2005. Stocking density and welfare of cage farmed Atlantic salmon: application of multivariate analysis. *Aquaculture* 243: 121-132.
- Van de Vis, J.W., 2001. is the co-ordinator of PCT application (PCT/NL01/00412), entitled "*Method and device for stunning and killing of aquatic animals*", emanating from the EU project on optimisation of harvest procedures of farmed fish contract FAIR CT-97-3127. The patent application was published in December 2001 (WO 01/95732 A1).
- Van de Vis, J.W., Kestin, S.C., Robb, D.F.H., Oehlenschläger, J., Lambooi, E., Münkner, W., Kuhlmann, H., Münkner, W. Kloosterboer, R.J , Tejada, M., Huidobro, A., Tejada, M., Kloosterboer, R.J., Otterå, H., Roth, B., Sørensen, N.K., Aske., L. Byrne, H. & Nesvadba, P., 2003a. Is humane slaughter of fish possible for industry? *Aquaculture Research*, 34, 211-220.
- Van de Vis, J.W., Kloosterboer, R.J., Gerritzen., M.A. and Lambooi, E., 2003b. Development of a humane slaughter method for farmed African catfish (*Clarias gariepinus*). In *Proceedings of the First Joint Trans Atlantic Fisheries Technology Conference 10-14 June 2003, Reykjavik, Iceland*, Icelandic Fisheries Laboratories, Reykjavik, Iceland, pp 390-391, ISBN 9979-74-005-1.
- Vazzana, M., Cammarata, M., Cooper, E.L. & Parrinello, N., 2002. Confinement stress in seabass (*Dicentrarchus labrax*) depresses peritoneal leukocyte cytotoxicity. *Aquaculture*, 210, 231-243.
- Vijayan, M.M. & Leatherland, J.F., 1988. Effect of stocking density on the growth and stress-response in brook charr, *Salvelinus fontinalis*. *Aquaculture*, 75, 159-170.
- Vijayan, M.M., Ballamtyne, J.S. & Leatherland, J.F., 1990. High stocking density alters the energy metabolism of

- brook charr, *Salvelinus fontinalis*. *Aquaculture*, 88, 371-381.
- Vijayan, M.M., Pereira, C.E., Grau, G. & Iwama, G.K., 1997. Metabolic responses associated with confinement stress in tilapia: the role of cortisol. *Comparative Biochemistry and Physiology*, 116, 89-95.
- Vincent-Lang, D., Alexandersdottir, M. & McBride, D., 1993. Mortality of coho salmon caught and released using sport tackle in the Little Susitna River, Alaska. *Fisheries Research*, 15, 339-356.
- Waagbø, R., Bjerås, E., Sveier, H., Breck, O., Bjerås, I., Bjørnstad, E. & Maage, A., 1996. Nutritional status assessed in groups of smelting Atlantic salmon, *Salmo salar* L., developing cataracts. *Journal of Fish Diseases*, 19, 365-373.
- Waagbø, R., Sveier, H., Breck, O., Bjørnstad, E., Maage, A., & Bjerås, E., 1998. Cataract Formation in Smolting Atlantic Salmon, *Salmo salar*, Fed Low and High Energy Diets. *Bulletin of the European Association of Fish Pathologist*, 18, 201-205.
- Wagner, E.J., Bosakowski, T. & Intelmann, S., 1997. Combined effects of temperature and high pH on mortality and the stress response of rainbow trout after stocking. *Transactions of the American Fisheries Society*, 126, 995-998.
- Waring, C.P., Stagg, R.M., Poxton, M.G., 1996. Physiological responses to handling in the turbot. *Journal of Fish Biology*, 48, 161-173.
- Warner, K., 1979. Mortality of landlocked Atlantic salmon (*Salmo salar* L.) hooked on flies and worms in the river nursery area. *Transactions of the American Fisheries Society*, 107, 518-522.
- Wassenberg, T.J. & Hill, B.J., 1989. The effect of trawling and subsequent handling on the survival rates of the by-catch of prawn trawlers in Moreton Bay, Australia. *Fisheries Research*, 7, 99-110.
- Wedemeyer, G.A., 1996. *Physiology of fish in intensive culture systems*. Chapman and Hall, New York, 272 pp.
- Wendelaar Bonga, S.E., 1997. The stress response in fish. *Physiological Reviews* 77, 591-625.
- Wertheimer, A., 1988. Hooking mortality of chinook salmon released by commercial trollers. *North American Journal of Fisheries Management*, 8, 346-355.
- Weyts, F.A.A., Cohen, N., Flik, G. & Verburg-Kemenade, B.M.L., 1999. Interactions between the immune system and the hypothalamus-pituitary-interrenal axis in fish. *Fish & Shellfish Immunology*, 9, 1-20.
- Wicks, B.J., Tang, Q., Joensen, R., Randall, D.J. 2002. Swimming and ammonia toxicity in salmonids: the effect of sub lethal ammonia exposure on the swimming performance of coho salmon and the acute toxicity of ammonia in swimming and resting rainbow trout. *Aquatic Toxicology*. 59: 55-69.
- Winfree, R.A., Kindschi, G.A. & Shaw, H.T., 1998. Elevated water temperature, crowding, and food deprivation accelerate fin erosion in juvenile steelhead. *The Progressive Fish-Culturist*, 60, 192-199.
- Wolffrom, T., 2004. Farmed fish and welfare. European Commission, Directorate-General For Fisheries, Research and Scientific Analysis Unit (A4). 39 p. http://europa.eu.int/comm/fisheries/doc_et_publ/liste_publi/farmedfish_en.pdf
- Wydoski, R.S., Wedemeyer, G.A. & Nelson, N.C., 1976. Physiological response to hooking stress in hatchery and wild rainbow trout (*Salmo gairdneri*). *Transactions of the American Fisheries Society*, 105, 601-606.
- Yin, Z., Lam, T.J. & Sin, Y.M., 1995. The effects of crowding stress on the non-specific immune response in fancy carp (*Cyprinus carpio* L.). *Fish and Shellfish Immunology*, 5, 519-529.

CHAPTER 8 AVAILABLE INFORMATION ON THE INTERNET

8.1 Angling organizations

<http://www.anglersnet.co.uk/saa/index.htm>

Web site of the Specialist Anglers' Alliance (SAA). The SAA is an angling/conservation/political body, which is the umbrella organization representing specialist-angling groups, which are in SAA membership. The National Angling Alliance Code of Conduct can be found on their web site.

<http://www.anglingfoundation.com/>

The Angling Foundation of the United Kingdom provides a code for retaining nets.

<http://www.recfishoz.com/>

Web site of Recfish Australia, an organization with a mission to represent the interests of recreational and sport fishers at a national level to ensure quality of fishing in Australia. The national code of practice for recreational and sport fishing 2001 can be found on this web site.

<http://www.ssaa.org.au/fish.html>

A position statement of the Sporting Shooters Association of Australia, regarding animal welfare aspects of recreational fishing.

<http://www.anglersnet.co.uk/saa/naa.htm>

Sport fisheries organization: National Angling Alliance, Fishmongers' Chambers, 110 Upper Thames Street, London EC4R 3TW, United Kingdom.

<http://www.fisheries.co.uk/ccfa/index.htm>

Commercial fisheries organization: Commercial Coarse Fisheries Association, C/o Tingrith Fishery, Tingrith, Bedfordshire MK17 9EW, United Kingdom.

<http://www.cfb.ie>

Irish Central Fisheries Board

<http://www.vdsf.de>

Largest angler's organization in Germany, also providing a code of angling practice in concordance with animal welfare.

<http://www.anglerverband.com/frame.htm>

The web site of the German anglers association. It contains a code of honour ("Ehrenkodex") how to deal with fishes.

8.2 Animal rights organizations

<http://www.animalfreedom.org/>

Animal Freedom provides information about the impact of angling on the fish and the surrounding environment.

<http://www.animalfrontline.nl/>

Animal Frontline is against recreational fisheries, ornamental fish keeping and commercial sea fisheries.

<http://www.anti-angling.com/>

Web site of the Campaign for the Abolition of Angling, dedicated to banning angling. Details can be found regarding campaigns, a comprehensive fact sheet and the scientific evidence regarding fish and pain.

<http://www.dierenbescherming.nl/>:

The “Dierenbescherming” is a Dutch organisation that provides information about angling and the results of angling on the welfare of fish. The “Dierenbescherming” are against all forms of angling.

<http://www.nofishing.net/>

A site linked to PETA, which provides information against recreational and commercial fisheries and the consumption of fish.

<http://www.pisces.demon.co.uk/welfare.html>

The report “The Welfare of Fish and Aquatic Invertebrates” by the Submission of the Australian and New Zealand Federation of Animal Societies.

<http://www.rspca.org.uk/>

RSPCA Web site of the RSPCA (the Royal Society for the Prevention of Cruelty to Animals). The RSPCA focuses on promoting kindness and preventing cruelty to animals.

<http://www.veganisme.non-profit.nl/>

The “Nederlandse Vereniging voor Veganisme” provides information against the bio-industry.

<http://www.vissenbescherming.nl/>

The “Vissenbescherming” is a Dutch organisation that provides information about keeping goldfish in small bowls, commercial sea fisheries, eel fisheries and recreational fisheries. The “Vissenbescherming” would like to see a ban on goldfish bowls and recreational fisheries. Furthermore, they strive for improved slaughter methods for the commercial sea fisheries and eel fisheries.

8.3 Animal welfare organizations

<http://www.awfc.ca/home.html>

Web site of the Animal Welfare Foundation of Canada. The report “A Report on the Animal Welfare Aspects of Fish Farming” can be found on this Web site.

<http://www.ciwf.co.uk/>

Web site of “Compassion in World Farming”. Several reports dealing with the welfare of fish in aquaculture can be found on this web site.

<http://www.fawc.org.uk/>

Web site of the Farm Animal Welfare Council (FAWC) (5th Floor, 1A Page Street, London SW1P 4PQ, United Kingdom), an independent advisory body established by the Government in 1979. Its terms of reference are to keep under review the welfare of farm animals on agricultural land, at market, in transit and at the place of slaughter; and to advise the Government of any legislative or other changes that may be necessary. The report “Report On The Welfare Of Farmed Fish” can be found on this web site.

<http://www.nal.usda.gov/awic/>

Web site of the Animal Welfare Information Centre, which provides information for improved animal care and use in research, teaching, and testing. The publication “Responsible Care and Health Maintenance of Fish in Commercial Aquaculture” by T.E. Schwedler and S.K. Johnson can be found on this web site.

<http://www.psyeta.org/>

Web site of Psychologists for the Ethical Treatment of Animals (PSYETA). The publication “A Survey of Animal Care Considerations for Fish Handling” by Henrik Kreiberg can be found on this web site.

8.4 Aquaculture organizations

<http://www.aquaculturecouncilwa.com/>

Web site of the Aquaculture Council of Western Australia. The Guidelines on Aquatic Animal Welfare for the Aquaculture Industry in Western Australia can be found on this web site.

<http://www.britishtROUT.co.uk/bta.htm>

British Trout Association, Bow Business Centre, London E3 2SE. United Kingdom.

Scottish Salmon Growers Association, Drummond House, Scott Street, Perth PH1 5EJ, United Kingdom.

www.ifa.ie

Irish Salmon Growers Association.

<http://www.bim.ie>

Bord iascaigh Mhara (Irish Sea Fisheries Board)

<http://www.marine.ie>

Irish Marine Institute

<http://www.aquariumcouncil.org>

Marine Aquarium Council

<http://www.feap.info>

Federation of European Aquaculture Producers. The FEAP has a position paper on animal protection in fish aquaculture (2003).

8.5 Fisheries organizations

http://www.fisheries.org/html/Public_Affairs/Sound_Science/Guidelines2004.shtml

American Fisheries Society, American Institute of Fisheries Research Biologists & American Society of Ichthyologists and Herpetologists (2004) *Guidelines for the Use of Fishes in Research*. American Fisheries Society: Bethesda.

<http://www.ifm.org.uk/index.html>

Web site of the Institute of Fisheries Management (IFM). It provides the "Stillwater Coarse Fisheries Code of Practice by the Institute of Fisheries Management (IFM)". *The Management of Intensively Stocked Stillwater Coarse Fisheries Including the Codes of practice*. Nottingham: Institute of Fisheries Management, 39 pp.

<http://www.le.ac.uk/biology/fsbi/>

The FSBI (The Fisheries Society of the British Isles) is the premier Society in the British Isles, and increasingly in Europe, catering for the interests of professional fish biologists and fisheries managers. The Society is affiliated to the United Kingdom Institute of Biology and the Society for Science and Technology.

<http://www.leicester.ac.uk/biology/fsbi/welfare.pdf>

FSBI (2002) *Fish Welfare. Briefing Paper 2*. Cambridge: Fisheries Society of the British Isles

<http://www.efishbusiness.co.uk/>

eFishBusiness, England & Wales.

<http://www.vet.ed.ac.uk/animalwelfare/>

University of Edinburgh Animal Welfare Research Group, Easter Bush Veterinary Centre, Easter Bush, Roslin, Midlothian, EH25 9RG, U.K

<http://www.vet.ed.ac.uk/animalwelfare/Fish%20pain/welfare.htm#FARMED>

Web pages written by University of Edinburgh Animal Welfare Research Group M.Sc.students in 2002: WELFARE OF FARMED FISH and

<http://www.vet.ed.ac.uk/animalwelfare/Fish%20pain/welfare.htm#ANGLED>

WELFARE OF ANGLED FISH .

<http://www.atc.stir.ac.uk/staq/>

Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, United Kingdom.

8.6 Governmental and intergovernmental organizations

<http://www.fao.org/fi>

Web site of the Food and Agriculture Organization of the United Nations (FAO) and FAO fisheries department. The FAO is one of the largest specialised agencies in the United Nations system and the lead agency for agriculture, forestry, fisheries and rural development. The Code of Conduct for Responsible Fisheries can be found on the web site of FAO's Fisheries Department: <http://www.fao.org/fi/agreem/codecond/ficonde.asp>.

The web site of the EIFAC – European Inland Fisheries Advisory Commission, FAO.

<http://www.fao.org/fi/body/eifac/eifac.asp>

http://europa.eu.int/comm/fisheries/policy_en.htm

Web site of the European Commission's Directorate-General for Fisheries. It provides the Common Fisheries Policy (CFP).

http://europa.eu.int/comm/fisheries/doc_et_publ/liste_publi/farmedfish_en.pdf

Report on "Farmed fish and welfare" produced by the directorate-general of Fisheries of the European Commission.

<http://www.coe.int/> and http://www.coe.int/T/E/Legal_Affairs/

The site of the Council of Europe including information on the Standing Convention on the Protection of Animals.

http://www.efsa.eu.int/science/ahaw/ahaw_opinions/495_en.html.

Opinion of the European Food Safety Authority (EFSA) Scientific Panel on Animal Health and Welfare on a request from the Commission related to welfare aspects of the main systems of stunning and killing the main commercial species of animals.

<http://www.defra.gov.uk/>

Web site of the Department of Environment Food and Rural Affairs. It provides a wide variety of information on animal health and welfare. Amongst other things the report "Government Response to the Farm Animal Welfare Council's Report on the Welfare of Farmed Fish" is available on this web site.

<http://www.defra.gov.uk/Science/Publications/Report%20of%20the%20Workshop%20on%20Farmed%20Fish%20Welfare.pdf>

DEFRA Science Directorate (2002) *Report of the Workshop on Farmed Fish Welfare held on Monday 28th October 2002*. Department for Environment, Food and Rural Affairs. 37pp.

<http://www.fawc.org.uk/reports/fish/fishrtoc.htm>

FAWC (1996) *Report on the Welfare of farmed Fish*. The Farm Animal Welfare Council, 52 pp.

<http://www.verbraucherministerium.de>

Homepage of the German Ministry of Consumer Protection, Food and Agriculture

<http://www.verbraucherministerium.de/englisch/tierschutzbericht/tierschutzbericht-2001.htm#03>

Review of the German regulations and legislation on animal welfare.

<http://www.minlnv.nl/>

Web site of the Dutch Ministry of Agriculture, Nature Management and Fisheries. It provides information regarding the Dutch policy to commercial fisheries. It provides several publications on commercial fisheries in the Netherlands and on the welfare of fish.

<http://www.nal.usda.gov/awic/pubs/Fishwelfare/fishwelfare.htm>

Web site of the U.S. Department of Agriculture, Agricultural Research Service, National Agricultural Library, Animal Welfare Information Center, Beltsville, Maryland. It provides detailed information regarding fish welfare from 1970-2003 (Erickson, 2003).

<http://www.homeoffice.gov.uk/comrace/animals/licensing.html>

The Home Office of the United Kingdom Government operates the licensing and inspection systems according to the Animals (Scientific Procedures) Act 1986

In Ireland the following organizations are involved in Handling of fishes:

- Dept of Communication, Marine & Natural Resources www.dcmnr.gov.ie
- An Bord Iascaigh Mhara – commercial & aquaculture www.bim.ie
- Marine Institute – aquaculture and experimental www.marine.ie
- Central Fisheries Board www.cfb.ie
- Animal Welfare Committee, Veterinary Ireland www.veterinary-ireland.org
- Dept. of Agriculture & Food www.agriculture.gov.ie

8.7 Scientific organizations

<http://www.ICES.dk>

Web site of the International Council for the Exploration of the Sea.

<http://199.245.200.110/>

Web site of the American Society of Ichthyologists and Herpetologists. The American Society of Ichthyologists and Herpetologists is dedicated to the scientific study of fishes, amphibians and reptiles. The publication “Guidelines for Use of Fishes in Field Research” can be found on this web site.

<http://oslovet.veths.no/>

Web site of the Norwegian Reference Centre for Laboratory Animal Science & Alternatives. It provides the Norwegian Animal Welfare Act.

<http://www.aps.uoguelph.ca/~aquacentre/>

Web site of the Aquaculture Centre, a collaborative between the University of Guelph and the Ontario Ministry of Agriculture and Food. Amongst other things it provides a reference source of scientific literature on welfare of fish.

<http://www.ifb-potsdam.de/>

Web site of the Institut für Binnenfischerei in Potsdam near Berlin. Main objectives of the research work done by this institute are questions of the ecologically oriented fisheries management of waters as well as problem definitions of the environmental compatible and market-focused aquaculture; also questions of animal welfare in fisheries and aquaculture included. The Ifb does not represent the German Institut für Binnenfischerei. It is an Institut financed by several Länder of Germany.

<http://www.curtin.edu.au/>

Web site of the Curtin University of Technology. The Australian code of practice for the care and use of animals for scientific purposes can be found on this site.

<http://www.hafro.is/catag/>

<http://www.nal.usda.gov/awic/pubs/Fishwelfare/Thorsteinsson.pdf>

Site with information for those who want to obtain information on tagging or marking of fish. The CATAG (Concerted Action for Tagging of Fishes) group has collected information on: tags, tagging, experiments where tags are used, legislation concerning tagging, health and behaviour changes, which can be induced by tagging and models used to work on data from tagging experiments.

<http://www.bfa-fisch.de>

Homepage of the Federal Research Center for Fisheries. Information on animal welfare is included.

<http://www.ntva.no/>

The Norwegian Academy of Technological Sciences has the objective of promoting research and education in technology and related sciences. The Holmenkollen Guidelines for Sustainable Aquaculture 1998 can be found on their web site.

http://www.vet.ed.ac.uk/animalwelfare/animal_pain.htm

Web site of the Royal School of Veterinary Studies, providing information on the perception of pain in fish.

<http://www.cefas.co.uk/homepage.htm#>

The Fish Health Inspectorate, CEFAS Weymouth Laboratory, Barrack Road, The Nothe, Weymouth, Dorset DT4 8UB, United Kingdom

<http://uwadmnweb.uwo.edu/Zoology/faculty/Rose/Rose%20Rev%20Fish%20Sci.pdf>

Paper by J.D. Rose (2002), The neurobehavioral nature of fishes and the question of awareness and pain. *Reviews in Fisheries Science* 10: 1-38.

8.8 Codes and guidelines

<http://www.defra.gov.uk/animalh/welfare/fawc/fish/fishrtoc.htm>

The Report on the Welfare of Farmed Fish by the FAWC.

http://www.geocities.com/b_c_aquaculture/code.html

Draft Code of Practice for the salmon farming industry by the B.C. Salmon Farmers Association.

<http://www.anglersnet.co.uk/saa/Code2002.htm>

The Code of Conduct for European Aquaculture by the Federation of European Aquaculture Producers (FEAP). the original web site for the Code of Conduct is:

http://www.feap.info/feap/code/default_en.asp

<http://www.aquamedia.org>

<http://www.ntva.no/rapport/aqua/report.htm>

The Holmenkollen Guidelines for Sustainable Aquaculture 1998.

<http://www.fao.org/fi/agreem/codecond/ficond.asp>

The Code of Conduct for Responsible Fisheries by the FAO Fisheries and Aquaculture Department.

<ftp://ftp.fao.org/docrep/fao/005/v9878e/V9878E00.pdf>

FAO Technical Guidelines for Responsible Fisheries No. 5: Aquaculture development (FAO 1997).

<ftp://ftp.fao.org/docrep/fao/003/W4493e/W4493e00.pdf>

<http://www.recfishoz.com/CoP%20web%202001.htm>

The National Code of Practice for Recreational and Sport Fishing by eleven national member associations of Recfish Australia.

http://www.intrafish.com/code-practise/conduct_eu_aqua.pdf

The Code of Conduct for Coarse Anglers by the National Angling Alliance.

http://www.fisheries.org/html/Public_Affairs/Sound_Science/Guidelines2004.shtml

The Guidelines for the Use of Fishes in Research (Nickum *et al.*, 2004) were developed to help scientists develop realistic understandings of fish in the design and conduct of research on fish and fish habitats. The Guidelines provide a structure that ensures appropriate attention to valid experimental design and procedures while ensuring humane treatment of fish. Written by the 11-member Uses of Fishes in Research Committee, the Guidelines provide recommendations on field and laboratory activities, such as sampling, holding, and handling fishes; information on administrative matters, including regulations and permits; and advice on ethical questions such as perceptions of pain or discomfort that may be experienced by experimental subjects. The 53-page document now includes many listings of web sites of value.

The Guidelines have been developed for use by researchers within the United States; therefore, the roles, responsibilities, and information needs of Institutional Animal Care and Use Committees are given specific attention. However, the principles described in the Guidelines are applicable to research on fishes everywhere.

<http://www.anglersnet.co.uk/code.pdf>

Code of Conduct for Coarse Anglers. London: National Angling Alliance, 40 pp.

www.cfb.ie/fishing_in_ireland/CatchandRelease.htm

Catch and Release for Atlantic Salmon in Ireland, Central Fisheries Board Leaflet No. 1.

The Canadian Council on Animal Care has produced a detailed series of Guidelines on: the care of fish in research, teaching and testing (Griffin & Gauthier, 2003).

8.9 Other

<http://www.cabi-publishing.org/Bookshop/ReadingRoom/085199296x.asp>

An online version of "Animal Welfare And Meat Science" by N.G. Gregory, amongst other things dealing with the welfare aspects of commercial fisheries and aquaculture.

http://www.oie.int/eng/normes/fcode/A_summry.htm

An overview of international guidelines and recommendations on infectious, aquatic diseases.

<http://www.primate.wisc.edu/pin/welfare.html>

Overview of American regulations on animal welfare.

<http://www.uni-giessen.de/tierschutz/>

<http://www.vetinfo.demon.nl/aw/flist.html>

A reference source of scientific literature on welfare of fish.

<http://www.vetmed.ucdavis.edu/CCAB/fish.htm>

A reference source of literature on welfare of fish.

<http://homepage.tinet.ie/~pnowlan/>

LAST – Laboratory Animal Science & Teaching in Ireland.

APPENDIX

BIBLIOGRAPHY

1 References for "Animal welfare"

- Algers, B., 2004. See Chapter 7 *Cited References*.
- Anon. 1997. *A report on the animal welfare aspects of fish farming*. Animal Welfare Foundation of Canada.
- Anonymous. 1979. See Chapter 7 *Cited References*.
- Baeverfjord, G. 1998. Ethics and animal welfare in intensive aquaculture production. *Farming Marine Fish beyond the year 2000: Technological Solutions for Biological Challenges*. ICES C.M. L: 18, 3p.
- Battye, J. 1994. Ethics and Animal Welfare - Where do we go from here? In: *Animal Welfare in the 21st Century: Ethical, Educational and Scientific Challenges* (Eds. R.M. Baker, D.J. Mellor and A.M. Nicol, p. 3-10).
- Bernoth E. M. & Wormuth H. J. 1990. Animal protection aspects on killing fish. *Deutsche Tierärztliche Wochenschrift* 97, 154-157.
- Bernoth E. M. 1991. Intensive culture of freshwater fish. *Deutsche Tierärztliche Wochenschrift* 98, 312-316.
- Bower, J., 1990. *Farm animal welfare in the 1990s. Some hopes and hindrances*. Oxford, England, Education Services, p. 16.
- Brambell, F.W.R., 1965. *Report of the technical committee to enquire into the welfare of animals kept under intensive husbandry systems*. Command Report 2836, Her Majesty's Stationary Office, London.
- Broom, D.M., 1993. A usable definition of animal welfare. *Journal of Agricultural and Environmental Ethics* (Supplement 2) 6, 15-25.
- Broom, D.M., 1998. Welfare stress, and the evolution of feelings. *Advances in the Study of Behaviour*, 27, 371-403.
- Broom, D.M., 1998. Fish welfare and the public perception of farmed fish. Report Aquavision '98. The second Nutreco aquaculture business conference Stavanger forum. Addressing the challenges to maintain a sustainable 10-20% annual growth of the aquaculture industry, pp 89-91.
- Broom, D.M. & Johnson, K.G. (eds.), 1993. *Stress and Animal Welfare*. Chapman & Hall, London.
- Casebolt, D.B., Speare, D.J. & Horney, B.S., 1998. Care and use of fish as laboratory animals: current state of knowledge. *Laboratory Animal Science*, 48, 124-136.
- Cawley, G., 1993. Welfare aspects of aquatic veterinary medicine. In: Brown, L. (ed.) *Aquaculture for veterinarians: fish husbandry and medicine*, Pergamon Press, Chicago, USA, p. 169-173.
- Chandroo, K.P., 1999. Animal welfare and Aquaculture. *Ontario Aquaculture Association Newsletter*, March-April, 3.
- Chandroo, K.P., 2000. Assessing the welfare status of farmed rainbow trout (*Oncorhynchus mykiss*) with electromyogram telemetry. Masters Thesis, University of Guelph.
- Council of Europe, 2000. *Draft Recommendation Concerning Farmed Fish (5th revision), 23^d October 2000*. Standing Committee of the European Convention for the Protection of Animals Kept for Farming Purposes (T-AP), Strasbourg.
- Crawley, G., 1993. Welfare aspects of aquatic veterinary medicine. In: Brown, L. (ed.) *Aquaculture for veterinarians*. Oxford: Pergamon Press.
- Curtis, S.E., 1985. What constitutes animal well-being? In: *Animal Stress* (ed. by G.P. Moberg), pp. 1-13. American Physiological Society, Bethesda, Maryland.
- Curtis, S.E. & Stricklin, W.R., 1991. The importance of animal cognition in agricultural animal production systems: an overview. *Journal of Animal Science*, 69, 5001-5007.
- Dantzer, R., 1993. Research perspectives in farm animal welfare: the concept of stress. *Journal of Agricultural and Environmental Ethics* (Supplement 2), 6, 86-92.
- Dantzer, R., Mormede, P., Henry, J.P., 1983. Physiological assessment of adaptation in farm animals. In: *Farm Animal Housing and Welfare*. S.H. Baxter, M.R. Baxter and J.A.C MacCormack (Eds.). Martinus Nijhoff Publishers, p. 8-19.
- Davenport, K., 1993. See Chapter 7 *Cited References*.
- Dawkins, M.S., 1980. *Animal Suffering: the Science of Animal Welfare*. Chapman and Hall, London.
- Dawkins, M.S., 1995. *Unravelling Animal Behaviour*. 2nd Edition, Longman, London.
- Dawkins, M.S., 1998. Evolution and animal welfare. *The Quarterly Review of Biology*, 73, 305-328.
- Degrazia, D., 1999. Animal ethics around the turn of the twenty-first century. *Journal of Agricultural and Environmental Ethics*, 11, 111-129.
- Dol, M., Fentener van Vlissingen, M., Kasanmoentalib, S., Visser & T., Zwart, H.,

1999. Recognizing the intrinsic value of animals. In: *Beyond animal welfare*. Animals in Philosophy and Science volume 2. Assen: Van Gorcum.
- Duncan, I.J.H., 1990. Animal welfare what is it and how can we measure it? In: *Farm and Animal Welfare in the 1990's*: A symposium sponsored by Alberta Institute of Agrologists, Edmonton Branch, p. 15.
- Duncan, I.J.H., 1993. Welfare is to do with what animals feel. *Journal of Agricultural and Environmental Ethics*, 6, 8-14.
- Duncan, I.J.H., 1996. Animal Welfare Defined in Terms of Feelings. *Acta Agriculturae Scandinavica*, Section A, Animal Science, Supplementum 27, 29-35.
- Duncan, I.J.H. & Fraser, D., 1997. See Chapter 7 *Cited References*.
- Duncan, I.J.H. & Petherick, J.C., 1991. The implications of cognitive processes for animal welfare. *Journal of Animal Science*, 69, 5017-5022.
- Erickson, H.S., 2003. See Chapter 7 *Cited References*.
- Ewbank, R., 1993. Animal welfare: a historical overview. *Journal of Agricultural and Environmental Ethics* (Supplement 1), 6, 82-86.
- FAWC, 1996. *Report on the welfare of farmed fish*. Farm Animal Welfare Council, Surrey.
- Fish Veterinary Society, 1993. *Guidelines on the welfare of farmed fish*. Fish Veterinary Society Publication.
- Flecknell, P.A., Molony, V., 1997. *Pain and Injury*. *Animal Welfare*, pp. 63-73, CAB International. University Press, Cambridge.
- Fraser, D., 1993. Assessing animal well-being: Common sense, uncommon science. *Food Animal Well-Being*. 1993 Conference Proceedings and Deliberations. p. 37-54. Purdue, Univ., West Lafayette, IN.
- Fraser, D., 1999. Animal ethics and animal welfare science: Bridging the two cultures. *Applied Animal Behavioural Science*, 65, 171-189.
- Fraser, D. & Duncan, I.J.H., 1998. 'Pleasures', 'Pains' and animal welfare: toward a natural history of affect. *Animal Welfare*, 7, 383-396.
- Fraser, D. & Leonard, M.L., 1993. *Farm animal welfare*. *Animal production in Canada*. University of Alberta, Edmonton, p. 340.
- Fraser, D. & Matthews, L.R., 1997. Preference and motivation testing. *Animal Welfare*, pp. 159-173. CAB International. University Press, Cambridge.
- Fraser, D. *et al.* 1997. See Chapter 7 *Cited References*.
- FSBI, 2002. See Chapter 7 *Cited References*.
- Gonyou, H.W., 1993. Animal welfare: definitions and assessment. *Journal of Agricultural and Environmental Ethics* (Supplement 2) 6, 37-43.
- Gregersen, F., 1999. Research on fish welfare: small efforts prevent stress and aggression. *Fiskeriforskning Info.*, Norwegian Institute of Fisheries and Aquaculture Ltd. May, No. 6.
- Griffin, G. & Gauthier, C., 2003. Guidelines development and scientific uncertainty: use of previous case studies to promote efficient production of guidelines on the care and use of fish in research, teaching and testing. *Animal welfare* 13 Supplement: 181-186.
- Håstein, T., 2004. See Chapter 7 *Cited References*.
- Hatt, J., 1998. Animal rights growing concern for aquaculturists. *Northern Aquaculture* May, 3.
- Hill, J.R., Sainsbury, W.B., 1990. *Farm animal welfare: Who cares? How?* The Cambridge Centre for Animal Health and Welfare, Cambridge, UK.
- Humphrey, N.,K., 1981. Having Feelings, and Showing Feelings. Self-Awareness in Domesticated Animals, 37-39. Universities Federation for Animal Welfare. Potters Bar, England.
- Hurnik, J.F. & Lehman, H., 1988. Ethics and farm animal welfare. *Journal of Agricultural Ethics*, 1, 305-318.
- Hurnik, J.F., 1993. Ethics and Animal Agriculture. *Journal of Agricultural and Environmental Ethics* (Supplement 1) 6, 21-35.
- Jensen, K.K. & Sørensen, J.T., 1999. The idea of "ethical accounting" for a livestock farm. *Journal of Agricultural and Environmental Ethics*, 11, 85-100.
- Kristiansen, T.S. & Juell, J.E., 2002. How can we document the welfare of farmed fish"? In *Seafarming Today and Tomorrow*. Pp 285-286. Compiled: B. Basurco & M. Saroglia. *European Aquaculture Society Special Publication* 32, August 2002.
- Lymbery, P., 1992. *The welfare of farmed fish*. *Compassion in World Farming*, Hants. Cattleshall Printing Services.
- Manteca, X., 1998. Neurophysiology and assessment of welfare. *Meat Science* (Supplement 1), 49, S205-S218.
- McGlone, J.J., 1993. What is animal welfare? *Journal of Agricultural and Environmental Ethics* (Supplement 2) 6, 26-36.
- Mench, J.A., 1993. Assessing animal welfare: an overview. *Journal of Agricultural and Environmental Ethics* (Supplement 2) 6, 68-75.
- Mench, J.A., Madson, G.J., 1997. Behaviour. *Animal Welfare*, 127-141. CAB International. University Press, Cambridge.
- Mendl, M., 1991. Some problems with the concept of a cut-off point for determining when an animal's welfare is at risk. *Applied Animal Behaviour Science*, 31, 139-146.

- Midtlyng, P.J., 1997. Vaccinated fish welfare: protection versus side-effects. *Developments of Biological Standardization*, 90, 371-379.
- Moberg, G.P., 1985. Biological response to stress: key to assessment of animal well-being? *Animal Stress*, 27-49. American Physiological Society, Bethesda, Maryland.
- Moberg, G.P., 1993. Using risk assessment to define domestic animal welfare. *Journal of Agricultural and Environmental Ethics* (Supplement 2) 6, 1-7.
- Needham, E.A., Lehman, H., 1991. Farming salmon ethically. *Journal of Agricultural and Environmental Ethics*, 14, 78-81.
- Needham, T., 1990. Do fish have rights? *Canadian Aquaculture* September/October, 45-46.
- Neukirch, M., 1994. Legitimate aspects relevant to animal protection on killing fish. *Deutsche Tierärztliche Wochenschrift*, 101, 316-319.
- Neuman, I.S.A., 1990. *Welfare of fishes in aquaculture*. Utrecht: Rijksuniversiteit Utrecht, 21p.
- Newman, S., 1994. Quantitative and molecular genetic effects on animal well-being: adaptive mechanisms. *Journal of Animal Science*, 72, 1641-1653.
- Nickum, J.G., H. L. Bart, Jr, P. R. Bowser, I. E. Greer, C. Hubbs, J. A. Jenkins, J. R. MacMillan, J. W. Rachlin, J. D. Rose, P. W. Sorensen & J. R. Tomasso, 2004. *Guidelines for the use of fishes in research*. American Fisheries Society, American Institute of Fisheries research Biologists, American Society of Ichthyologists and Herpetologists.
- OIE, 2004. Proceedings Global conference on animal welfare: an OIE initiative, Paris, 23025 February 2004. Luxembourg: Office for Official Publications of the European Communities. 333 p.
- Peters, G., 1990. Problems concerning animal protection laws in connection with mass culture of fishes. *Deutsche Tierärztliche Wochenschrift*, 97, 157-160.
- Pottinger, T.G., 1995. *Fish welfare literature review: Prepared for the Angling Governing Bodies Liaison Group and the British Fields Sports Society*. Ambleside: Institute of Freshwater Ecology, 83p.
- Price, E.O., 1985. Evolutionary and ontogenetic determinants of animal suffering and well-being. *Animal Stress* 15-25. American Physiological Society, Bethesda, Maryland.
- Raat, A.J.P. Raat & Van den Bos, R. (eds.), 1999. *Welzijn van Vissen [Welfare of Fishes]*. Tilburg. Tilburg University Press. 211 pp (in Dutch).
- Rodd, R., 1990. *Biology, Ethics, and Animals*. Clarendon Press, New York, p. 272.
- Rollin, B., 2000. Animal ethics - an emerging force to be reckoned with. *Northern Aquaculture* 6, 9.
- Rollin, B.E., 1993. Animal welfare, science, and value. *Journal of Agricultural and Environmental Ethics* (Supplement 2), 6, 44-50.
- Rollin, B.E., 1996. Bad ethics, good ethics and the genetic engineering of animals in agriculture. *Journal of Animal Science*, 74, 535-554.
- RSPAC, 2002. Welfare Standards for Farmed Atlantic Salmon, issue 2. 2002.
- Sandoe, P. & Simonsen, H.B., 1992. Assessing animal welfare: where does science end and philosophy begin? *Animal Welfare*, 1, 257-267.
- Sauer, N., 1993. *Tierschutz bei Fischen*. Giessen: Universität Giessen: 290 p.
- Sauer, N. & Manz, D., 1994. Tierschutzbestände bei Fischen. *Tierärztliche Umschau*, 49 (10), 653-658
- Schwedler, T.E., Johnson, S.K., 1997. Responsible care and health maintenance of fish in commercial aquaculture. *Animal Welfare Issues Compendium*, United States Department of Agriculture.
- Sherwin, C., 2001. See Chapter 7 Cited References.
- Singer, P., 1975. *Animal Liberation - A new ethics for our treatment of animals*. Random House Publishers, New York, p. 301.
- Spedding, C.R.W., 1993. Animal welfare policy in Europe. *Journal of Agricultural and Environmental Ethics* (Supplement 1) 6, 110-117.
- Sumner, L.W., 1988. Animal welfare and animal rights. *Journal of Medical Philosophy*, 13, 159-175.
- Van den Bos, R., (ed.) 1996. *Welzijn van dieren en Wetenschap : Door welke bril kijkt u? : Essays over de plaats van het welzijn van dieren in de wetenschap en de gevolgen hiervan voor het dierenwelzijnsbeleid*. [Animal welfare and science: From what perspective do you approach: essay on the position of animal welfare in science and the consequences of this to animal welfare policy]. Tilburg: Tilburg University Press, 88 p (in Dutch).
- Van den Bos, R., (ed.) 1997. *Welzijn van dieren en dierenwelzijnsbeleid. Essays over doelstellingen, instrumenten en evaluatie van dierenwelzijnsbeleid in Nederland*. [Animal welfare and animal welfare policy. Essays on aims, instruments and evaluation of animal welfare policy in The Netherlands]. Tilburg: Tilburg University Press, 97 p (in Dutch).
- Webster, J., 1984. Problems of confinement on farms / in laboratories and in the home. *Priorities in Animal Welfare*. The British Veterinary Association Animal Welfare Foundation, 55-61.

- Webster, J., 1995. *Animal Welfare: a cool eye towards eden*. Blackwell Science Ltd., London.
- Wemelsfelder, F., 1997. Investigating the animal's point of view. An inquiry into a subject-based

method of measurement in the field of animal welfare. *Animal Consciousness and Animal Ethics*. Van Gorcum, Assen. pp. 73-89.

- Wunderlich, G., 1991. The ethics of animal agriculture. *Food Review*, 14, 24-27.

2. References for "Painperception of fish"

2.1 Painperception

- Arena, P.C. & Richardson, K.C., 1990. The relief of pain in cold-blooded vertebrates. Australian Council for the Care of Animals in Research and Teaching-News, Canberra, A.C.T.: Australian Council for the Care of Animals in Research & Teaching (ACCART), Autumn 1990, v. 3, 1, 1-4.
- Bateson, P., 1991. Assessment of pain in animals. *Animal Behaviour*, 42, 827-839.
- Bateson, P., 1992. Do animals feel pain? *New Scientist*, 30-33.
- Bennett, G.J., 2000. Update on the neurophysiology of pain transmission and modulation: focus on the NMDA-receptor. *Journal of Pain and Symptom Management*, Supplement. 19, S2-S6.
- Bolles, R.C. & Fanselow, M.S., 1980 Perceptual-defensive-recuperative model of fear and pain. *Behavioral and Brain Sciences*, 3, 291-323.
- Braithwaite, V.A. & Huntingford, F.A., 2004. See Chapter 7 Cited references.
- Brown, L.A., 1985. The clinical measurement of pain, distress and discomfort in fish. Symposia Series I, *The detection and relief of pain in animals*. pp. 98-100. The British Veterinary Association Animal Welfare Foundation.
- Brown, L.A., Scott, P.W. & Greenwood, A.G., 1986. Pain in fish: A review. *Exotic animals in the eighties. Proceedings from the 25th Anniversary Symposium of the British Veterinary Zoological Society*, 18-20 April 1986. British Veterinary Zoological Society, place of publication not given. 1987: 1-64, chapter pagination: 107-110.
- Chandroo, K.P., Duncan, I.J.H. & Moccia R.D., 2004. Can fish suffer?: perspectives on sentience, pain, fear and stress. *Applied Animal Behaviour Science*, 86, 225-250.
- Chapman, R.C. & Nakamura, Y., 1999. A passion of the soul: an introduction to pain for conciousness researchers. *Consciousness and Cognition*, 8, 391-422.
- Chervova, L.S., 1996. Pain sensitivity and behavior of fishes. *Journal of Ichthyology*, 37, 1, 98-102.
- Erdmann, C., 1999. *The ability of fishes to feel pain and to suffer*. 154pp.
- Duncan, I.J.H., 2004. See Chapter 7 Cited References.
- Gregory, N., 1999. Do fish feel pain? *Surveillance*, 26, 8-10.
- Kestin, S.C., 1993. *Pain and stress in fish*. RSPCA (Royal Society for the Prevention of Cruelty to Animals). Bristol: RSPCA, 36 p.
- Kitchell, R.L. & Guinan, M.J., 1990. The nature of pain in animals. *The Experimental Animal in Biomedical Research*, Volume 1, 185-204. CRC Press, Inc., Boca Raton.
- Kitchell, R.L. & Johnson, R.D., 1985. Assessment of pain in animals. In: *Animal Stress*, 113-140. American Physiological Society. Bethesda, Maryland.
- Klausewitz, W., 1995. Pain, fear & ability to suffer in fish. A problematic complex activated by the new German Animal Protection Law. *Fortschritte.-Fishereiwiss. Adv. Fish Science* 12, 5-21.
- Klausewitz, W., 1989. Problems of sensitivity to pain and ability to suffer in fish. *Fischoekologie* 1, 1, 65-90.
- Morton, D.B. & Griffiths, P.H.M., 1985. Guidelines on the recognition of pain, distress and discomfort in experimental animals and an hypothesis for assessment. *Veterinary Record*, 116, 431-436.
- Oidtmann, B. & Hoffmann, R.W., 2001. Schmerzen und leiden bei Fischen. *Berliner und Munchener tierärztliche Wochenschrift*, 114, 7-8, 277-282.
- Peters, G., 1988. Schmerz und Stress bei Fischen. *Deutsche Tierärztliche Wochenschrift*, 95, 2, 60-63.
- Rose, J.D., 2002. See Chapter 7 Cited References.
- Rose, J.D., 2003. See Chapter 7 Cited References.
- Rose, M. & Adams, D., 1989. Evidence for pain and suffering in other animals. *Animal Experimentation: The Consensus Changes*, 42-71. Chapman and Hall, New York.
- Sanford, J., Ewbank, R., Molony, V., Tavernor, W.D. & Uvarov, O., 1986. Guidelines for the recognition and assessment of pain in animals. *Veterinary Record* 118, 334-338.
- Sneddon, L.U., 2003. See Chapter 7 Cited References.
- Sneddon, L.U. et al., 2003. See Chapter 7 Cited References.
- Stoskopf, M.K., 1994. Pain and analgesia in birds, reptiles, amphibians, and fish. *Investigative Ophthalmology & Visual Science*, 35, 775-780.
- Treede, R.D., Kenshalo, D.R., Gracely, R.H., Jones, A.K.P., 1999. The cortical representation of pain. *Pain* 79, 105-111.
- Trigg, R., 1970. *Pain and emotion*. Oxford: Clarendon Press.

- Verheijen, F.J. & Buwalda, R.J.A., 1988. *Doen pijn en angst een gehaakte en gedrilde karper lijden?* Utrecht: Vakgroep Vergelijkende Fysiologie, Rijksuniversiteit Utrecht, 40 p.
- Wright, E.M., Marcella, K.L. & Woodson, J.F., 1985. Animal pain: Evaluation and control. *Lab Animal* 120, 21-26.
- Yue, S., Moccia, R.D. & Duncan, I.J.H., 2004. Investigating fear in domestic rainbow trout *Oncorhynchus mykiss*, using an avoidance learning task. *Applied Animal Behaviour Science (In press)*. Presentation International Society of Applied Ethology, 37th International Congress, Abano Terme, Italy. June 24-27, 2003.
- Zieglgänsberger, W., 1986. Central control of nociception. *Handbook of Physiology*. Volume 4; *The Nervous System*, 581-645. American Physiological Society. Bethesda, Maryland.
- Zimmerman, M., 1986. Assessing pain in farm animals. *Proceedings Commission of European Communities Workshop*. October 1986.
- 2.2 Anatomical, Pharmacological and Neurophysiological Studies Pertaining to Pain, Fear and Stress in Fish
- Bernstein, J.J., 1970. Anatomy and physiology of the central nervous system. *Fish Physiology*, Vol. IV, Academic Press, New York, p. 1-78.
- Bradford Jr., M.R., 1995. Comparative aspects of forebrain organization in the ray-finned fishes: touchstones or not? *Brain, Behavior and Evolution*, 46, 259-274.
- Butler, A.B., 2000. Topography and topology of the teleost telencephalon: a paradox resolved. *Neuroscience Letters*, 293, 95-98.
- Cameron, A.A., Snow, P.J. & Plenderleith, M.B., 1990. Organization of the spinal cord in four species of elasmobranch fish: Cytoarchitecture and distribution of serotonin and selected neuropeptides. *Journal of Comparative Neurology*, 297, 201-218.
- Carruth, L.L., Jones, R.E. & Norris, D.O., 2000. Cell density and intracellular translocation of glucocorticoid receptor-immunoreactive neurons in the Kokanee salmon (*Oncorhynchus nerka kennerlyi*) brain, with an emphasis on the olfactory system. *General and Comparative Endocrinology*, 117, 66-76.
- Coggeshall, R.E., Leonard, R.B., Applebaum, M.L. & Willis, W.D., 1978. Organization of peripheral nerves and spinal roots of the Atlantic stingray, *Dasyatis sabina*. *Journal of Neurophysiology*, 41, 97-107.
- Corrêa, S.A.L., Corrêa, F.M.A. & Hoffmann, A., 1998. Stereotaxic atlas of the telencephalon of the weekly electric fish *Gymnotus carapo*. *Journal of Neuroscience Methods*, 84, 93-100.
- Davis, R.E., Kassel, J., 1983. Behavioral functions of the teleostean telencephalon. *Fish Neurobiology*. Volume 2: *Higher Brain Areas and Functions*, pp. 238-263. University of Michigan Press, Ann Arbor.
- Demski, L.S., 1983. Behavioural effects of electrical stimulation of the brain. In: *Fish Neurobiology*. Volume 2: *Higher Brain Areas and Functions*, pp. 317-359. University of Michigan Press, Ann Arbor.
- Devor, A., 2000. Is the cerebellum like cerebellar-like structures? *Brain Research Reviews*, 34, 149-156.
- Echteler, S.M. & Saidel, W.M., 1981. Forebrain connections in the goldfish support telencephalic homologies with land vertebrates. *Science* 212, 683-684.
- Ehrensing, R.H. et al. 1982. See Chapter 7 Cited References.
- Fibiger, H.C. & Phillips, A.G., 1986. Reward, motivation, cognition: psychobiology of mesotelencephalic dopamine systems. *Handbook of Physiology*. Volume 4; *The Nervous System*, pp. 647-675. American Physiological Society. Bethesda, Maryland.
- Finger, T.E., 1980. Nonolfactory sensory pathway to the telencephalon in a teleost fish. *Science*, 210, 671-673.
- Finger, T.E., 1983. Organization of the teleost cerebellum. *Fish Neurobiology*. Volume 1: *Brain Stem and Sense Organs*, pp.261-284. University of Michigan Press, Ann Arbor.
- Goping, G., Pollard, H.B., Adeyemo, O.M. & Kuijpers, G.A.J., 1995. Effect of MPTP on dopaminergic neurons in the goldfish brain: a light and electron microscope study. *Brain Research* 687, 35-52.
- Guthrie, D.M., 1983. Integration and control by the central nervous system. *Control Processes in Fish Physiology*, pp. 130-154. Croom Helm, London.
- Hatta, K., Korn, H., 1999. Tonic inhibition alternates in paired neurons that set direction of fish escape reaction. *Proceedings of the National Academy of Sciences of the United States States of America*, 96, 12090-12095.
- Hildebrand, M., 1995. *Analysis of Vertebrate Structure*, Fourth edition. John Wiley & Sons, Inc., New York.
- Hon, W.K. & Ng, T.B., 1986. Hormones with adrenocorticotrophic and opiate-like activities from the carp (*Cyprinus carpio*) pituitary. *Comparative Biochemistry and Physiology*, 85C, 443-448.
- Jansen, G.A. & Green, N.M., 1970. Morphine metabolism and morphine tolerance in goldfish. *Anesthesiology*, 32, 231-235.
- Kestin, S.C., 1994. *Pain and stress in fish. A report prepared for the RSPCA*. Royal Society for the Prevention of Cruelty to Animals, 36 pp.

- Kotrschal K., van Staaden M. J. & Huber R. 1998. Fish brains: evolution and environmental relationships. *Reviews in Fish Biology and Fisheries* 8, 373-408.
- Le Moal, M. & Simon, H., 1991. Mesocorticolimbic dopaminergic network: functional and regulatory roles. *Physiological Reviews*, 71, 155.
- Lett, B.T. & Grant, V.L. 1989. The hedonic effects of amphetamine and pentobarbital in goldfish. *Pharmacology Biochemistry & Behavior*, 32, 355-356.
- Matthews, G. & Wickelgren, W.O., 1978. Trigeminal sensory neurons of the sea lamprey. *Journal of Comparative Physiology*, 123, 329-333.
- Mattioli, R., Aguilar C. & Vasconcelos, L., 1995. Reinforcing properties of the neuropeptide substance P in *Carassius auratus*: evidence of dopaminergic system involvement. *Pharmacology Biochemistry and Behavior*, 50, 77-81.
- Mattioli, R., Santangelo, E.M., Costa, A.C.C. & Vasconcelos, L., 1997. Substance P facilitates memory in goldfish in an appetitively motivated learning task. *Behavioural Brain Research*, 85, 117-120.
- Mok, E.Y.M. & Munro, A.D., 1998. Effects of dopaminergic drugs on locomotor activity in teleost fish of the genus *Oreochromis* (Cichlidae): involvement of the telencephalon. *Physiology and Behavior*, 64, 227-234.
- Munro, A.D., 1986. The effects of apomorphine, *d*-amphetamine and chlorpromazine on the aggressiveness of isolated *Aequidens pulcher* (Teleostei, Cichlidae). *Psychopharmacology*, 88, 124-128.
- Munro, A.D., Dodd, J.M., 1983. Forebrain of fishes: neuroendocrine control mechanisms. *Progress in Nonmammalian Brain III* (ed. by G. Nisticò & L. Bolis), pp. 2-78. CRC Press Inc, Florida.
- Ng, T.B. & Chan, T.H., 1990. Adrenocorticotropin-like and opiate-like materials in the brain of the red grouper *Epinephelus akaara* (Teleostei: serranidae). *Comparative Biochemistry and Physiology*, 95C, 159-162.
- Nieuwenhuys, R., 1982. An overview of the organization of the brain of actinopterygian fishes. *American Zoologist* 22, 287-310.
- Ohnishi, K., 1997. Effects of telencephalic ablation on short-term memory and attention in goldfish. *Behavioural Brain Research*, 86, 191-199.
- Overmier, J.B. & Hollis, K.L., 1983. The teleostean telencephalon in learning. *Fish Neurobiology*. Volume 2: *Higher Brain Areas and Functions* (ed. by R.E. Davis & R.G. Northcutt), pp. 265-283. University of Michigan Press, Ann Arbor.
- Overmier, J.B. & Papini, M.R., 1986. Factors modulating the effects of teleost telencephalon ablation on retention, relearning, and extinction of instrumental avoidance behavior. *Behavioral Neuroscience*, 100, 190-199.
- Pavlov, D.S. & Kasumyan, A.O., 1995. A review of Russian studies on the behavior and sensory systems of fishes. Report 3. *Journal of Ichthyology*, 35, 806-815.
- Peters, G. *et al.* 1980. See Chapter 7 *Cited References*.
- Piront, M. & Schmidt, R., 1988. Inhibition of long-term memory formation by anti-ependymin antisera after active shock-avoidance learning in goldfish. *Brain Research*, 442, 53-62.
- Pollard, H.B., Kuijpers, G.A., Adeyemo, O.M., Youdim, M.B.H. & Goping, G., 1996. The MPTP-induced parkinsonian syndrome in the goldfish is associated with major cell destruction in the forebrain and subtle changes in the optic tectum. *Experimental Neurology*, 142, 170-178.
- Prechtl, J.C., Von der Emde, G., Wolfart, J., Karamürsel, S., Akoev, G.N., Andrianov, Y.N. & Bullock, T.H., 1998. Sensory processing in the pallium of a mormyrid fish. *Journal of Neuroscience*, 18, 7381-7393.
- Previc, F.H., 1999. Dopamine and the origins of human intelligence. *Brain and Cognition* 41, 299-350.
- Rolls, E.T., 2000. Précis of The brain and emotion. *Behavioral and Brain Sciences*, 23, 177-234.
- Rovainen, C.M. & Yan, Q., 1985. Sensory responses of dorsal cells in the lamprey brain. *Journal of Comparative Physiology A*, 156, 181-183.
- Salas, C., Broglio, C., Rodríguez, F., López, J.C., Portavella, M. & Torres, B., 1996. Telencephalic ablation in goldfish impairs performance in a 'spatial constancy' problem but not in a cued one. *Behavioural Brain Research*, 79, 193-200.
- Salas, C., Rodríguez, F., Vargas, J.P., Durán, E. & Torres, B., 1996. Spatial learning and memory deficits after telencephalic ablation in goldfish trained in place and turn maze procedures. *Behavioral Neuroscience*, 110, 965-980.
- Savage, G., 1980. The fish telencephalon and its relation to learning. *Comparative Neurology of the Telencephalon* (ed. by S.O.E. Ebbesson), pp.129-174. Plenum Press, New York.
- Schmidt, R., 1995. Cell-adhesion molecules in memory formation. *Behavioural Brain Research*, 66, 65-72.
- Snow, P.J., Plenderleith, M.B. & Wright, L.L., 1993. Quantitative study of primary sensory neurone populations of three species of elasmobranch fish. *Journal of Comparative Neurology*, 334, 97-103.

- Sovrano, V.A., Rainoldi, C., Bisazza, A. & Vallortigara, G., 1999. Roots of brain specializations: preferential left-eye use during mirror-image inspection in six species of teleost fish. *Behavioural Brain Research*, 106, 175-180.
- Van de Kar, L.D. & Blair, M.L., 1999. Forebrain pathways mediating stress-induced hormone secretion. *Frontiers in Neuroendocrinology*, 20, 1-48.
- Vargas, J.P., Rodríguez, F., López, J.C., Arias, J.L. & Salas, C. 2000. Spatial learning-induced increase in the argyrophilic nucleolar organizer region of dorsolateral telencephalic neurons in goldfish. *Brain Research*, 865, 77-84.
- Wendelaar Bonga, S.E., 1997. See Chapter 7 *Cited References*.
- Wendelaar Bonga, S.E. & Balm, P.H.M., 1999. Histological and histopathological effects of stress. In: Balm, P.H.M. (ed.) *Stress physiology in Animals*. Sheffield, Academic Press: 178-204.
- Winberg, S., Carter, C.G., McCarthy, I.D., He, Z.Y., Nilsson, G.E. & Houlihan, D.F., 1993. Feeding rank and brain serotonergic activity in rainbow trout *Oncorhynchus mykiss*. *Journal of Experimental Biology*, 179, 197-211.
- Winberg, S. & Lepage, O., 1998. Elevation of brain 5-HT activity, POMC expression, and plasma cortisol in socially subordinate rainbow trout. *American Journal of Physiology*, 274, R645-R654.
- Winberg, S. & Nilsson, G.E., 1993. Time course of changes in brain serotonergic activity and brain tryptophan levels in dominant and subordinate juvenile arctic charr. *Journal of Experimental Biology*, 179, 181-195.
- Zottoli, S.J., Newman, B.C., Rieff, H.I. & Winters, D.C., 1999. Decrease in occurrence of fast startle responses after selective Mauthner cell ablation in goldfish (*Carassius auratus*). *Journal of Comparative Physiology*, A 184, 207-218.
- Zupanc, G.K.H., 1997. The preglomerular nucleus of gymnotiform fish: relay station for conveying information between telencephalon and diencephalon. *Brain Research*, 761, 179-191.
- 2.3 *Cognitive ability, fear and awareness in fish*
- Bermond, B., 1992. *Leed bij dieren of de mythe van het bewustzijn*. Amsterdam: Universiteit van Amsterdam.
- Bermond, B., 1996. Dierenleed: een neuropsychologische analyse van moedwil en misverstand. *Nederlands Tijdschrift voor de Psychologie*, 51, 121-134.
- Bermond, B., 1997. See Chapter 7 *Cited References*.
- Bisazza, A., De Santi, A. & Vallortigara, G., 1999. Laterality and cooperation: mosquitofish move closer to a predator when the companion is on their left side. *Animal Behaviour*, 57, 1145-1149.
- Bisazza, A., Pignatti, R. & Vallortigara, G., 1997. Detour tests reveal task- and stimulus-specific behavioural lateralization in mosquitofish (*Gambusia holbrooki*). *Behavioural Brain Research*, 89, 237-242.
- Bisazza, A., Rogers, L.J. & Vallortigara, G., 1998. Origins of cerebral asymmetry: a review of evidence of behavioural and brain lateralization in fishes, reptiles and amphibians. *Neuroscience and Biobehavioural Reviews*, 22, 411-426.
- Bunge, M. & Ardila, R., 1987. *Philosophy of Psychology*. Springer-Verlag, New York.
- Canfield, J.G. & Rose, G.J., 1993. Activation of mauthner neurons during prey capture. *Journal of Comparative Physiology A*, 172, 611-618.
- Cantalupo, C., Bisazza, A. & Vallortigara, G., 1995. Lateralization of predator-evasion response in a teleost fish (*Girardinus falcatus*). *Neuropsychologia*, 33, 1637-1646.
- Csányi, V., 1986. Ethological analysis of predator avoidance by the paradise fish (*Macropodus opercularis* L.): II. Key stimuli in avoidance learning. *Animal Learning & Behavior*, 14, 101-109.
- Csányi, V., 1993. How genetics and learning make a fish an individual: a case study on the paradise fish. *Perspectives in Ethology*. Volume 10; *Behavior and Evolution* (ed. by P.P.G. Bateson, P.H. Klopfer & N.S. Thompson), pp. 1-51. Plenum Press, New York.
- Csányi, V., Csizmadia, G. & Miklosi, A., 1989. Long-term memory and recognition of another species in the paradise fish. *Animal Behaviour*, 37, 908-911.
- Csányi, V. & Dóka, A., 1993. Learning interactions between prey and predator fish. *Marine and Freshwater Behaviour and Physiology*, 23, 63-78.
- Csányi, V. & Lovász, F., 1987. Key stimuli and the recognition of the physical environment by the paradise fish *Macropodus opercularis*. *Animal Learning & Behavior*, 15, 379-381.
- Davis, R.E. & Klinger, P.D., 1994. NMDA receptor antagonist MK-801 blocks learning of conditioned stimulus-unconditioned stimulus contiguity but no fear of conditioned stimulus in goldfish (*Carassius auratus* L.). *Behavioral Neuroscience*, 108, 935-940.
- Eaton, R.C., Lavender, W.A. & Wieland, C.M., 1981. Identification of Mauthner-initiated response in goldfish: evidence from simultaneous cinematography and

- electrophysiology. *Journal of Comparative Physiology*, 144, 521-531.
- Eaton, R.C., Lavender, W.A. & Wieland, C.M., 1982. Alternative neural pathways initiate fast-start responses following lesions of the Mauthner neuron in goldfish. *Journal of Comparative Physiology*, 145, 485-496.
- Eaton, R.C. & Nissanov, J., 1985. A review of Mauthner-initiated escape behaviour and its possible role in hatching in the immature zebrafish, *Brachydanio rerio*. *Environmental Biology of Fishes*, 12, 265-279.
- Facchin, L., Bisazza, A. & Vallortigara, G., 1999. What causes lateralization of detour behavior in fish? evidence for asymmetries in eye use. *Behavioural Brain Research*, 103, 229-234.
- Hall, D. & Suboski, D., 1995. Visual and olfactory stimuli in learned release of alarm reactions by zebra danio fish (*Brachydanio rerio*). *Neurobiology of Learning and Memory*, 63, 229-240.
- Harper, D.G. & Blake, R.W. 1990. Fast-start performance of rainbow trout *Salmo gairdneri* and northern pike *Esox Lucius*. *Journal of Experimental Biology*, 150, 321-342.
- Höglund, E., Balm, P.H. & Winberg, S., 2000. Skin darkening, a potential social signal in subordinate Arctic charr (*Salvelinus alpinus*): the regulatory role of brain monoamines and pro-opiomelanocortin-derived peptides. *Journal of Experimental Biology*, 203, 1711-1721.
- Höjesjö, J., Johnsson, J.I. & Axelsson, M., 1999. Behavioural and heart rate responses to food limitation and predation risk: an experimental study on rainbow trout. *Journal of Fish Biology*, 55, 1009-1019.
- Johnsson, J.I., 1997. Individual recognition affects aggression and dominance relations in rainbow trout, *Oncorhynchus mykiss*. *Ethology*, 103, 267-282.
- Johnsson, J.I. & Åkerman, A., 1998. Watch and learn: preview of the fighting ability of opponents alter contest behaviour in rainbow trout. *Animal Behaviour*, 56, 771-776.
- Jones, R.B., 1997. Fear and distress. *Animal Welfare* (ed. by M.C. Appleby & B.O. Hughes), pp. 75-87. CAB International. University Press, Cambridge.
- Kilian, A., Von Fersen, L. & Güntürkün, O., 2000. Lateralization of visuospatial processing in the bottlenose dolphin (*Tursiops truncatus*). *Behavioural Brain Research*, 116, 211-215.
- Lachlan, R.F., Crooks, L. & Laland, K.N., 1998. Who follows whom? Shoaling preferences and social learning of foraging information in guppies. *Animal Behaviour*, 56, 181-190.
- López, C.J., Broglio, C., Rodríguez, F., Thinus-Blanc, C. & Salas, C., 2000. Reversal learning deficit in a spatial task but not in a cued one after telencephalic ablation in goldfish. *Behavioural Brain Research*, 109, 91-98.
- López, J.C., Bingman, V.P., Rodríguez, F., Gómez, Y. & Salas, C., 2000. Dissociation of place and cue learning by telencephalic ablation in goldfish. *Behavioral Neuroscience*, 114, 687-699.
- Losey Jr., G.S. & Margules, L., 1974. Cleaning symbiosis provides a positive reinforcer for fish. *Science*, 184, 179-180.
- Mendl, M., 1999. Performing under pressure: stress and cognitive function. *Applied Animal Behaviour Science*, 65, 221-244.
- Metcalfe, N.B., Taylor, A.C. & Thorpe, J.E., 1995. Metabolic rate, social status and life-history strategies in Atlantic salmon. *Animal Behaviour*, 49, 431-436.
- Miklósi, Á. & Andrew, R.J., 1999. Right eye use associated with decision to bite in zebrafish. *Behavioural Brain Research*, 105, 199-205.
- Nesse, R.M., 2000. Is depression and adaptation? *Archives of General Psychiatry*, 57, 14-20.
- O'Connor, K.I., Metcalfe, N.B. & Taylor, A.C., 1999. Does darkening signal submission in territorial contests between juvenile Atlantic salmon, *Salmo salar*? *Animal Behaviour*, 58, 1269-1276.
- O'Connor, K.I., Metcalfe, N.B. & Taylor, A.C., 2000. Familiarity influences body darkening in territorial disputes between juvenile salmon. *Animal Behaviour*, 59, 1095-1101.
- Olla, B.L. & Davis, M.W., 1989. The role of learning and stress in predator avoidance of hatchery-reared coho salmon (*Oncorhynchus kisutch*) juveniles. *Aquaculture*, 76, 209-214.
- Øverli, Ø., Harris, C.A. & Winberg, S., 1999. Short-term effects of fights for social dominance and the establishment of dominant-subordinate relationships on brain monoamines and cortisol in rainbow trout. *Brain, Behaviour and Evolution*, 54, 263-275.
- Øverli, Ø., Winberg, S., Damsgård, B. & Jobling, M., 1998. Food intake and spontaneous swimming activity in Arctic char (*Salvelinus alpinus*): role of brain serotonergic activity and social interactions. *Canadian Journal of Zoology*, 76, 1366-1370.
- Pitcher, T.J., 1993. *Behaviour of Teleost Fishes*, 2nd edition. Chapman & Hall, London.
- Popper, A.N. & Carlson, T.J., 1998. Application of sound and other stimuli to control fish behavior. *Transactions of the American Fisheries Society*, 127, 673-707.
- Schaerer, S. & Kirschfeld, K., 2000. The role of background movement in goldfish vision. *Journal of Comparative Physiology A*, 186, 583-593.
- Topál, J. & Csányi, V., 1999. Interactive learning in the paradise fish (*Macropodus opercularis*): an ethological interpretation of the second-

- order conditioning paradigm. *Animal Cognition*, 2, 197-206.
- Vallortigara, G., Rogers, L.J. & Bisazza, A., 1999. Possible evolutionary origins of cognitive brain lateralization. *Brain Research Reviews*, 30, 164-175.
- Varner, G.E., 1998. In Nature's Interests? *Interests, Animal Rights, and Environmental Ethics. Environmental Ethics and Science Policy Series*, Oxford University Press, U.S.A.
- Wainwright, P.C., 1986. Motor correlates of learning behaviour: feeding on novel prey by pumpkinseed sunfish (*Lepomis gibbosus*). *Journal of Experimental Biology*, 126, 237-247.
- ### 3. References for "Commercial fisheries"
- #### 3.1 Electric fishing
- Ainslie, B.J. *et al.* 1998. See Chapter 7 *Cited References*.
- Bardygula-Nonn, L.G., Nonn, R. & Savitz, J., 1995. Influence of pulsed direct current electrofishing on mortality and injuries among four centrachid species. *North American Journal of Fisheries Management*, 15, 799-803.
- Barton, B.A. & Dwyer, W.P., 1997. Physiological stress effects of continuous and pulsed-DC electroshock on juvenile bull trout. *Journal of Fish Biology*, 51, 998-1008.
- Barton, B.A. & Grosh, R.S., 1996. Effect of AC electroshock on blood features in juvenile rainbow trout. *Journal of Fish Biology*, 49, 1330-1333.
- Carline, R.F., 2001. Effects of high-frequency pulsed-DC electrofishing on a wild brown trout population. *North American Journal of Fisheries Management*, 21, 571-579.
- Cho, G.K. *et al.* 2002. See Chapter 7 *Cited References*.
- Cowdell, B.R. & Valdez, R.A., 1994. Effects of pulsed DC electroshock on adult roundtail chub from the Colorado River in Colorado. *North American Journal of Fisheries Management*, 14, 659-660.
- Cowx, I & Lamarque, P., 1990. See Chapter 7 *Cited References*.
- Dalbey, S.R., McMahon, T.E. & Fredenberg, W., 1996. Effect of electrofishing pulse shape and electrofishing-induced spinal injury on long term growth and survival of wild rainbow trout. *North American Journal Fisheries Management*, 16, 560-569.
- Dolan, C.R. *et al.* 2002. See Chapter 7 *Cited References*.
- Dwyer, W.P. & Erdahl, D.A., 1995. Effects of electroshock voltage, wave form, and pulse rate on survival of cutthroat trout eggs. *North American Journal of Fisheries Management*, 15, 647-650.
- Dwyer, W.P., Fredenberg, W. & Erdahl, D.A., 1993. Influence of electroshock and mechanical shock on survival of trout eggs. *North American Journal of Fisheries Management*, 13, 839-843.
- Dwyer, W.P. *et al.* 2001. See Chapter 7 *Cited References*.
- Dwyer, W.P. & White, R.G., 1995. Influence of electroshock on short-term growth of adult rainbow trout and juvenile arctic grayling and cutthroat trout. *North American Journal of Fisheries Management*, 15, 148-151.
- Dwyer, W.P. & White, R.G., 1997. Effects of electroshock on juvenile arctic grayling and Yellowstone cutthroat trout growth, 100 days after treatment. *North American Journal of Fisheries Management*, 17, 174-177.
- Fredenberg, W.A., 1992. Evaluation of electrofishing-induced spinal injuries resulting from field electrofishing surveys in Montana. Montana Department of Fish, Wildlife, and Parks, Final Report, Helena.
- Gatz Jr., A.J., Loar, J.M. & Cada, G.F., 1986. Effects of repeated electroshocking on instantaneous growth of trout. *North American Journal of Fisheries Management*, 6, 176-182.
- Godfrey, H., 1957. Mortalities among developing trout and salmon ova following shock by direct-current electrical fishing gear. *Journal of the Fisheries Research Board of Canada*, 14, 153-164.
- Habera, J.W., Strange, R.J., Carter, B.D. & Moore, S.E., 1996. Short-term mortality and injury of rainbow trout caused by three-pass AC electrofishing in a southern Appalachian stream. *North American Journal of Fisheries Management*, 16, 192-200.
- Habera, J.W. *et al.* 1999. See Chapter 7 *Cited References*.
- Hauck, F.R., 1949. Some harmful effects of the electric shocker on large rainbow trout. *Canadian Journal of Fisheries and Aquatic Sciences*, 43, 763-771.
- Wood-Gush, D.G.M., Dawkins, M.S. & Ewbank, R., 1981. *Self-Awareness in Domesticated Animals*. Universities Federation for Animal Welfare. Potters Bar, England.
- Zayan, R., 1991. The specificity of social stress. *Behavioural Processes*, 25, 81-93.

- Transactions of the American Fisheries Society*, 77, 61-64.
- Hollender, B.A. & Carline, R.F., 1994. Injury to wild brook trout by backpack electrofishing. *North American Journal of Fisheries Management*, 14, 643-649.
- Holmes, R., McBride, D.N., Viavant, T. & Reynolds, J.B., 1990. Electrofishing induced mortality and injury to rainbow trout, arctic grayling, humpback whitefish, least cisco, and northern pike. Alaska Department of Fish and Game. Division of Sport Fish, Fisher Manuscript 90-3, Anchorage.
- Hudy, M., 1985. Rainbow trout and brook trout mortality from high voltage AC electrofishing in a controlled environment. *North American Journal of Fisheries Management*, 5, 475-479.
- Jepsen, N. & Aarestrup, K., 1999. See Chapter 7 *Cited References*.
- Jepsen, N. & Berg, S., 2002. See Chapter 7 *Cited References*.
- Keefe, M. & Whitesel, T.A., 2000. See Chapter 7 *Cited References*.
- Kocovsky, P.M., Gowan, C., Fausch, K.D. & Riley, S.C., 1997. Spinal injury rates in three wild trout populations in Colorado after eight years of backpack electrofishing. *North American Journal of Fisheries Management*, 17, 308-313.
- Maxfield, G.H., Lander, R.H. & Liscom, K.L., 1971. Survival, growth, and fecundity of hatchery-reared rainbow trout after exposure to pulsating direct current. *Transactions of the American Fisheries Society*, 100, 546-552.
- McCrimmon, H.R. & Bidgood, B., 1965. Abnormal vertebrae in the rainbow trout with particular reference to electrofishing. *Transactions of the American Fisheries Society*, 94, 84-88.
- McMichael, G.A., 1993. Examination of electrofishing injury and short-term mortality in hatchery rainbow trout. *North American Journal of Fisheries Management*, 13, 229-233.
- McMichael, G.A., Fritts, A.L. & Pearsons, T.N., 1998. Electrofishing injury to stream salmonids; injury assessment at the sample, reach, and stream scales. *North American Journal of Fisheries Management*, 18, 894-904.
- Mesa, M. G. & Schreck, C.B., 1989. Electrofishing mark-recapture and depletion methodologies evoke behavioural and physiological changes in cutthroat trout. *Transactions of the American Fisheries Society*, 118, 644-658.
- Meyer, C.W. & Millar, D.D., 1990. Spinal injury and mortality in rainbow trout and brown trout collected by electroshocking. *Proceedings of the Annual Conference Western Association of Fish and Wildlife Agencies*, 70, 54-61.
- Mitton, C.J.A. & McDonald, D.G., 1994a. Consequences of pulsed DC electrofishing and air exposure to rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences*, 51, 1791-1797.
- Mitton, C.J.A. & McDonald, D.G., 1994b. Effects of electroshock, air exposure, and forced exercise on swim performance in rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences*, 51, 1799-1803.
- Muth, R.T. & Ruppert, J.B., 1996. Effects of two electrofishing currents on captive ripe razorback suckers and subsequent egg-hatching success. *North American Journal of Fisheries Management*, 16, 473-476.
- Muth, R.T. & Ruppert, J.B., 1997. Effects of electrofishing on captive embryos and larvae of razorback sucker. *North American Journal of Fisheries Management*, 17, 160-166.
- Newman, L.E. & Stone, F.G., 1992. Reduced viability in walleye eggs caused by exposure to pulsed DC electrofishing. U.S. Fish and Wildlife Service, Ashland Fishery Resources Office, Ashland, Wisconsin.
- Pratt, V.S., 1955. Fish mortality caused by electrical shockers. *Transactions of the American Fisheries Society*, 84, 93-96.
- Reynolds, J.B. & Kolz, A.L., 1988. Comment: electrofishing injury to large rainbow trout. *North American Journal of Fisheries Management*, 8, 516-517.
- Roach, S.M., 1999. See Chapter 7 *Cited References*.
- Rümmler, F., Schreckenbach, K. & Pfeiffer, M., 1998. Auswirkungen der Elektrofischerei auf Fische. *Fischer & Teichwirt*, 49, 88-92.
- Rümmler, F., Schreckenbach, K., Göthling, U. & Schiewe, S., (in press). Untersuchungen zu den Auswirkungen des Elektrofischfangs auf Fische. *Schriften des Instituts für Binnenfischerei*, Bd. 14, Potsdam-Sacrow.
- Ruppert, J.B. & Muth, R.T., 1997. Effects on electrofishing fields on captive juveniles of two endangered cyprinids. *North American Journal of Fisheries Management*, 17, 314-320.
- Schill, D.J. & Elle, F.S., 2000. See Chapter 7 *Cited References*.
- Schreck, C.B., Whaley, R.A., Bass, M.L., Maughan, O.E. & Solazzi, M., 1976. Physiological responses of rainbow trout (*Salmo gairdneri*) to electroshock. *Journal of the Fisheries Research of Canada*, 33, 76-84.
- Sharber, N.G. & Carothers, S.W., 1988. Influence of electrofishing pulse shape on spinal injuries in adult rainbow trout. *North American Journal of Fisheries Management*, 8, 117-122.
- Sharber, N.G., Carothers, S.W., Sharber, J.P., De Vos Jr., J.C. & House, D.A., 1994. Reducing

electrofishing-induced injury of rainbow trout. *North American Journal of Fisheries Management*, 14, 340-346.

- Sorensen, P.W., 1994. Effects of electroshocking on the sexual behavior of goldfish and brook trout. *North American Journal of Fisheries Management*, 14, 862-865.
- Spencer, S.L., 1967. Internal injuries of largemouth bass and bluegills caused by electricity. *The Progressive Fish-Culturists*, 29, 168-169.
- Suuronen, P. *et al.* 1996. See Chapter 7 *Cited References*.
- Thompson, K.G., Bergersen, E.P. & Nehring, R.B., 1997. Injuries to brown trout and rainbow trout induced by capture with pulsed direct current. *North American Journal of Fisheries Management*, 17, 141-153.
- Thompson, K.G., Bergersen, E.P., Nehring, R.B. & Bowden, D.C., 1997. Long-term effects of electrofishing on growth and body composition of brown trout and rainbow trout. *North American Journal of Fisheries Management*, 17, 154-159.
- Tipping, J.M. & Gilhuly, G.J., 1996. Survival of electroanesthetized adult steelhead and eggs of fall Chinook salmon. *North American Journal of Fisheries Management*, 16, 469-472.
- VanderKooi, S.P., Gale, W.L. & Maule, A.G., 2000. Preliminary study of gill Na⁺, K⁺-ATPase activity in juvenile spring Chinook salmon following electroshock or handling stress. *North American Journal of Fisheries Management*, 20, 528-533.
- VanderKooi, S.P., Maule, A.G. & Schreck, C.B., 2001. The effects of electroshock on immune function and disease progression in juvenile spring chinook salmon. *Transactions of the American Fisheries Society*, 130, 397-408.

3.2 Bycatch

- Berghahn, R., Waltemath, M. & Rijnsdorp, A.D., 1992. Mortality of fish from the bycatch of shrimp vessels in the North Sea. *Journal of Applied Ichthyology*, 8, 293-306.
- Kaimmer, S.M. & Trumble, R.J., 1998. See Chapter 7 *Cited References*.
- Pascoe, S., 1997. Bycatch management and the economics of discarding. *FAO Fisheries Biology Technical Paper* 370.
- Ross, M.R. & Hokenson, S.R., 1997. Short-term mortality of discarded finfish bycatch in the Gulf of Maine fishery for northern shrimp *Pandalus borealis*. *North American Journal of Fisheries Management*, 17, 902-909.
- Van Beek, F.A., Van Leeuwen, P.I. & Rijnsdorp, A.D., 1989. Survival of plaice and sole discards in the otter and beam trawl fisheries in the North Sea. International Council for the Exploration of the Sea, Copenhagen, Denmark, 20 pp.

3.3 Fishing gear escape

- Chopin, F.S. & Arimoto, T., 1995. See Chapter 7 *Cited References*.
- Sangster, G., 1992. The survival of fish escaping from fishing gears. Copenhagen-Denmark ICES, 9 pp.
- Sangster, G.I. *et al.* 1996. See Chapter 7 *Cited References*.

3.4 Physiological effects

- Candy, J.R., Carter, E.W., Quinn, T.P. & Ridell, B.E., 1996. Adult Chinook salmon behavior and survival after catch and release from purse-seine vessels in Johnstone Strait, British Columbia. *North American Journal of Fisheries Management*, 16, 521-529.
- Chopin, F.S. *et al.* 1996. See Chapter 7 *Cited References*.
- Canley, M.L., Mayr, S.D., Young, P.S. & Cech Jr., J.J., 2002. Swimming performances and physiological stress responses of splittail exposed to a fish screen. *North American Journal of Fisheries Management*, 22, 1241-1249.
- Davis, M.W. & Olla, B.L., 2001. Stress and delayed mortality induced in pacific halibut by exposure to hooking, net towing, elevated seawater temperature and air: implications for management of bycatch. *North American Journal of Fisheries Management*, 21, 725-732.
- Davis, M.W., Olla, B.L. & Schreck, C.B., 2001. Stress induced by hooking, net towing, elevated sea water temperature and air in sablefish: lack of concordance between mortality and physiological measures of stress. *Journal of Fish Biology*, 58, 1-15.
- Hopkins, T. E. & Cech, J.J., 1992. See Chapter 7 *Cited References*.

3.5 Mortality

- Anonymous, 1993. See Chapter 7 *Cited References*.
- Buchanan, S. *et al.* 2002. See Chapter 7 *Cited References*.
- Caddy, J.F., 1973. See Chapter 7 *Cited References*.
- Carr, H.A., Robinson, W.E., Sullivan, P.A., Caruso, P. & Cruse, J.M., 1992. Survival of juvenile Atlantic cod and American Plaice in the Northwestern Atlantic trawl fishery. MTS-'92:-Global Ocean Partnership. *Proceedings Marine Technology Society*, Washington, DC USA, pp. 316-321.
- Davis, M.W. & Olla, B.L., 2002. Mortality of lingcod towed in a net as related to fish length, seawater temperature, and air exposure: a laboratory bycatch study. *North American Journal of Fisheries Management*, 22, 1095-1104.

- DeAlteris, J.T. & Reifsteck, D.M., 1993. See Chapter 7 *Cited References*.
- Dieterman, D.J. *et al.* 2000. See Chapter 7 *Cited References*.
- Dunning, D.J. *et al.* 1989. See Chapter 7 *Cited References*.
- Farrington, M., Milliken, H., Lent, E. & Carr, H.A., 1998. Selectivity and Survival of Atlantic Cod (*Gadus morhua*) and Haddock (*Melanogrammus aeglefinus*) in the Northwest Atlantic Longline Fishery. National Marine Fisheries Service, Northeast Region, One Blackburn Drive, Gloucester, MA 01930.
- Fritz, K.R. & Johnson, D.L., 1987. See Chapter 7 *Cited References*.
- Gallinat, M.P., Ngu, H.H. & Shively, J.D., 1997. Short-term survival of lake trout released from commercial gill nets in Lake Superior. *North American Journal of Fisheries Management*, 17, 136-140.
- Hay, D.E. *et al.* 1986. See Chapter 7 *Cited References*.
- Lawson, P.W. & Sampson, D.B., 1996. Gear-related mortality in selective fisheries for ocean salmon. *North American Journal of Fisheries Management*, 16, 512-520.
- Lockwood, S.J. *et al.* 1983. See Chapter 7 *Cited References*.
- Main, J. & Sangster, G.I., 1990. See Chapter 7 *Cited References*.
- McLoughlin, R.J. *et al.* 1991. See Chapter 7 *Cited References*.
- Murphy, M.D., Heagy, R.F., Neugebauer, V.H., Gordon, M.D. & Hintz, J.L., 1995. Mortality of spotted seatrout released from gill-net or hook-and-line gear in Florida. *North American Journal of Fisheries Management*, 15, 748-753.
- Nelson, J.D. *et al.* 1989. See Chapter 7 *Cited References*.
- Orsi, J.A., Wertheimer, A.C. & Jaenicke, H.W., 1993. Influence of selected hook and lure types on catch, size and mortality of commercially troll-caught chinook salmon. *North American Journal of Fisheries Management*, 13, 709-722.
- Pálsson, O.K. *et al.* 2003. See Chapter 7 *Cited References*.
- Riemens, R.G., 1984. See Chapter 7 *Cited References*.
- Sangster, G.I. & Lehmann, K., 1993. See Chapter 7 *Cited References*.
- Smith, E.M. & Howell, P.T., 1987. See Chapter 7 *Cited References*.
- Soldal, A.V. & Isaksen, B., 1993. See Chapter 7 *Cited References*.
- Soldal, A.V. *et al.* 1991. See Chapter 7 *Cited References*.
- Stevens, B.G., 1990. See Chapter 7 *Cited References*.
- Suuronen, P. *et al.* 1993. See Chapter 7 *Cited References*.
- Thompson, R.B. & Hunter, C.J., 1973. See Chapter 7 *Cited References*.
- Thompson, R.B. *et al.* 1971. See Chapter 7 *Cited References*.
- Wassenberg, T.J. & Hill, B.J., 1989. See Chapter 7 *Cited References*.

3.6 Other

- Czajkowski, S.P., Coble, D.W., Copes, F.A., Bruch, R., Kamke & Peeters, P.J., 1996. Lower returns of tagged walleyes initially caught by trawling than by trap nets, fyke nets, or electroshocking. *North American Journal Fisheries Management*, 16, 453-456.
- Engås, A., Kyrkjebø Haugland, E., Tore & Øvredal, J., 1998. Reactions of cod (*Gadus morhua* L.) in the pre-vessel zone to an approaching trawler under different light conditions. *Hydrobiologia*, 371/372, 199-206.
- Lowry, N., Ferro, R.S.T., Ulmestrand, M.A.M. & Vold-Sodal, A., 1998. Third European Marine Science and Technology Conference. Luxembourg European Commission DG 12 Science, Research and Development, 6, 82-83.
- Maitland, P. S. & Lyle, A. A., 1992. Conservation of freshwater fish in the British Isles : Proposals for management. *Aquatic Conservation Marine Freshwater Ecosystems* 2, 165-183.
- Maitland, P. S., 1991. *The exploitation of freshwater fish in Europe : A review of fishing methods*. Council of Europe, 39 p.
- Maitland, P. S., 1995. *Freshwater fish of annexes II and IV of the EC Habitats Directive (92/43/EEC) : Implementation of the Directive on "Habitats" 92/43/EEC EC European Community*. Stirling: Fish Conservation Centre, 179 p.
- Muenkner, W., Kuhlmann, H. & Oehlenschlaeger, J., 1998. Investigations on the sensitivity of fish after the catch on board. *Inf.-Fischwirtsch.* vol. 45, no. 2, pp. 89-92.
- Olla, B.L. *et al.* 1997. See Chapter 7 *Cited References*.
- Wardle, C.S., 1976. Fish reactions to fishing gears. *Scott. Fish. Bull.*, 43, 16-20.
- North American Journal of Fisheries Management* 9, 74-80.
- Barnhart, R.A. & Roelofs, T.D., 1979. *Catch-and-release fishing as a management tool: A*

4. References for "Recreational fisheries"

4.1 Catch and release

- Barnhart, R.A., 1989. Symposium review; catch and release fishing, a decade of experience.

- national sportfishing symposium*, september 7-8, 1977. Arcata: Humboldt State University, 220 p.
- Bendock, T. & Alexandersdottir, M., 1990. Hook and release mortality of chinook salmon in the Kenai river recreational Fishery. *Fish. Data Ser.* No. 90-16, Alaska Department of Fish and Game.
- Bendock, T. & Alexandersdottir, M., 1991. Hook and release mortality in the Kenai river recreational Fishery. *Fish. Data Ser.* No. 91-39, Alaska Department of Fish and Game.
- Bendock, T. & Alexandersdottir, M., 1993. Hooking mortality of Chinook salmon released in the Kenai River, Alaska. *North American Journal of Fisheries Management*, 13, 540-549.
- Bugley, K. & Sherperd, G., 1991. Effect of catch-and-release angling on the survival of black seabass. *North American Journal of Fisheries Management*, 11, 468-471.
- Burkett, D.P., Mankin, P.C., Lewis, G.W., Childers, F. & Phillip, D.P., 1986. Hook-and-line vulnerability and multiple recapture of largemouth bass under a minimum total-length limit of 457 mm. *North American Journal of Fisheries Management*, 6, 109-112.
- Burkholder, A., 1992. Mortality of northern pike captured and released with sport fishing gear. *Fish. Data Ser.* 92-3. Alaska Department of Fish and Game.
- Catch & Release – the Future is in your Hands* – a 20 minute video available from the Central Fisheries Board, Ireland. Video produced by, and available from, the Fisheries Research Services, Marine Laboratory, Aberdeen, Scotland.
- Childress, W.M., 1989. Catch-and-release mortality of white and black crappie. Pages 175-186 in Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California.
- Clark Jr., R.D., 1983. Potential effects of voluntary catch and release of fish on recreational fisheries. *North American Journal of Fisheries Management*, 3, 306-314.
- Clark, R.A., 1991. Mortality of arctic grayling captured and released with sporting gear. *Fish. Data Ser.* No. 91-59. Alaska Department of Fish and Game.
- Cooke, S.J., Phillip, D.P., Schreer, J.F. & McKinley, R.S., 2000. Locomotory impairment of nesting male largemouth bass following catch-and-release angling. *North American Journal of Fisheries Management*, 20, 968-977.
- Cooke, S.J., Schreer, J.F. & McKinley, R.S., 1999. Evaluating catch-and-release angling practices from the fish's perspective. *Evaluating-the-benefits-of-recreational-fisheries* Pitcher, T.J. (ed.), 7, 78-87.
- Ferguson, R.A. & Tufts, B.L., 1992. See Chapter 7 *Cited References*.
- Gjernes, T., Kronlund, A.R. & Mulligan, T.J., 1993. Mortality of chinook and coho salmon in their first year of ocean life following catch and release by anglers. *North American Journal of Fisheries Management*, 13, 524-539.
- Goeman, T.J., 1991. Walleye mortality during a live release tournament on Mille Lacs, Minnesota. *North American Journal of Fisheries Management*, 11, 57-61.
- Harrell, R.M., 1988. Catch and release mortality of striped bass caught with artificial lures and baits. *Proceedings of the Annual Conference Southeastern Association of Fish and Wildlife Agencies*, 41, 70-75.
- Haux, C. & Sjöbeck, M.L., 1985. Physiological stress responses in a wild fish population of perch (*Perch fluvialitis*) after capture and subsequent recovery. *Marine Environmental Research*, 15, 77-95.
- Hegen, H.E. *et al.* 1987. See Chapter 7 *Cited References*.
- Hubbard, W.D. & Miranda, L.E., 1991. Mortality of white crappie after catch and release. *Proceedings of the 43rd Annual Conference of Southeast Association of Fish and Wildlife Agencies*, 43, 49-55.
- Jackson, J.J. & Willis, D.W., 1991. Short-term mortality of smallmouth bass caught during a live-release tournament at Lake Oahe, South Dakota. *Prairie Nat.*, 2, 201-204.
- Jolley, J.W. & Irby Jr., E.W., 1979. Survival of tagged and released Atlantic sailfish (*Istiophorus platypterus*: Istiophoridae) determined with acoustical telemetry. *Ull. Mar. Sci.*, 29, 155-169.
- Kieffer, J.D., Kubacki, M.R., Phelan, F.J.S., Phillip, D.P. & Tufts, B.L., 1994. Effects of catch and releasing angling on nesting male smallmouth bass. *Transactions of the American Fisheries Society*, 124, 70-76.
- Lee, D.P., 1989. Mortality of tournament caught and released black bass in California. Pages 207-216 in Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California.
- Nelson, K.L., 1998. See Chapter 7 *Cited References*.
- Phillip, D.P., Toline, C.A., Kubacki, M.F., Phillip, D.B.F. & Phelan, F.J.S., 1997. The impact of catch-and-release angling on the reproductive success of smallmouth bass and largemouth bass. *North American Journal of Fisheries Management*, 17, 557-567.

- Siewert, H.F. & Cave, J.B., 1990. Survival of released bluegill, *Lepomis macrochirus*, caught on artificial flies, worms, and spinner lures. *Journal of Freshwater Ecology*, 5, 407-411.
- Steger, T.M., Grizzle, J.M., Weathers, K. & Newman, M., 1994. Bacterial diseases and mortality of angler-caught largemouth bass released after tournaments on Walter F. George Reservoir, Alabama/Georgia. *North American Journal of Fisheries Management*, 14, 435-441
- Thorstad, E.B. *et al.* 2003. See Chapter 7 *Cited References*.
- Vincent-Lang, D. *et al.* 1993. See Chapter 7 *Cited References*.
- Weidlein, W.D., 1989. Mortality of released sublegalized smallmouth bass, catch-and-release implications. Pages 217-228 in Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California.
- 4.2 *Physiological effects/ stress*
- Beggs, G.L., Holeyton, G.F. & Crossman, E.J., 1980. Some physiological consequences of angling stress in muskellunge, *Esox masquinongy* Mitchill. *Journal of Fish Biology*, 17, 649-659.
- Booth, R.K., Kieffer, J.D., Davidson, K., Bielak, A.T. & Tufts, B.L., 1995. Effects of late-season catch and release angling on anaerobic metabolism, acid-base status, survival and gamete viability in wild Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 52, 283-290.
- Broadhurst, M.K. & Barker, D.T., 2000. Effects of capture by hook and line on plasma cortisol, scale loss and survival in juvenile mulloway (*Agrysomus hololeidotus*). *Archive of Fishery and Marine Research*, 48, 1-10.
- Broadhurst, M.K., Barker, D.T. & Kennelly, S.J., 1999. Scale-loss and survival of juvenile yellowfin bream, *Acanthopagrus australis*, after simulated escape from a Nordmoere-grid guiding panel and release from capture by hook and line. *Bulletin of Marine Science*, 64, 255-268.
- Brobbel, M.A., Wilkie, M.P., Davidson, K., Kieffer, J.D., Bielak, A.T. & Tufts, B.L., 1996. Physiological effects of catch and release angling in Atlantic salmon (*Salmo salar*) at different stages of freshwater migration. *Canadian Journal of Fisheries and Aquatic Sciences*, 53, 2036-2043.
- Carmichael, G.J., Wedemeyer, G.A., McCraren, J.P. & Millard, J.L., 1983. Physiological effects of handling and hauling stress on smallmouth bass. *The Progressive Fish Culturist*, 45, 110-113.
- Carragher, J.F. & Pankhurst, N.W., 1991. See Chapter 7 *Cited References*.
- Cooke, S.J., Bunt, C.M. & McKinley, R.S., 1998. See Chapter 7 *Cited References*.
- Cooke, S.J. *et al.* 2001. See Chapter 7 *Cited References*.
- Davidson, G.W. *et al.* 1997. See Chapter 7 *Cited References*.
- Ferguson, R.A. & Tufts, B.L., 1992. See Chapter 7 *Cited References*.
- Gallman, E.A. *et al.* 1999. See Chapter 7 *Cited References*.
- Gustavson, A.W. *et al.* 1991. See Chapter 7 *Cited References*.
- Klinger, H., 1988. Schmerz und Stress beim geangelten Fisch. *Fischer und Teichwirt*, 39, 10-13.
- Klausewitz, W., 1990. Gutachten zur Anwendung des Setzkeschers beim Angelvorgang. (unpublished).
- Klausewitz, W., 1989. Über Schmerzempfinden und Leidensfähigkeit der Fische. *Fischökologie* 1: 65-90.
- Klausewitz, W., 1995. Schmerzen, Angst und Leidensfähigkeit bei Fischen- ein durch das novellierte deutsche Tierschutzgesetz aktualisierter Problemkomplex. *Fortschritte d. Fischereiwiss.* 12: 5-21.
- Koßmann, H. & Pfeiffer, W., 1997. Fische im Setzkescher. Eine Verhaltensstudie zur Hälterung von geangelten Fischen Videofilm mit Begleittext. 6 S., Herausgeber VDSF-Landesverband Rheinland Pfalz.
- Lowe, T.E. & Wells, R.M.G., 1996. See Chapter 7 *Cited References*.
- Perrier, C. Terrier, M. & Perrier, H. 1978. A time-course study of the effects of angling stress on cyclic AMP, lactate and glucose plasma levels in the rainbow trout (*Salmo gairdneri* Richardson) during a 64 hour recovery period. *Comparative Biochemistry Physiology*, 60A, 217-219.
- Poole, W.R., Nolan, D. & Tully, O. 2000. Extrapolating baseline blood cortisol levels in trout: modelling capture effects in wild sea trout *Salmo trutta* (L.) infested with *Lepeophtheirus salmonis* (Krøyer). *Aquaculture Research* 31 (11): 835-841.
- Raat, A. J. P. *et al.* 1997. See Chapter 7 *Cited References*.
- Ridgway-Mark, S. & Shuter-Brian, J., 1997. Predicting the effects of angling for nesting male smallmouth bass on production of age-0 fish with an individual-based model. *North American Journal of Fisheries Management*, 17, 568-580.
- Ridley, C.A., Isely, J.J. & Tomasso Jr., J.R., 1997. Effects of supplemental feeding on physical characteristics and physiological responses to angling of striped bass. *North American*

- Journal of Fisheries Management*, 17, 791-794.
- Schreckenbach, K. & Thürmer, C., 1999. Einfluss von Wasserströmungen auf die Streßreaktionen von geangelten Rotfedern (*Scardinius erythrophthalmus*) bei der Lebendhaltung im Setzkescher. Untersuchungen des Instituts für Binnenfischerei e. V. Potsdam Sacrow (unpublished).
- Schreckenbach, K. & Thürmer, C., 2000. Stressreaktionen von geangelten Süßwasserfischen bei ihrer Lebendhaltung im Mustersetzkescher „mosella“. Gutachten Inst. Binnenfischerei Potsd. Sacrow, 16 p. (unpublished).
- Schreckenbach, K. & Wedekind, H., 1996. Einfluß des Angelns und der Setzkescherhaltung auf die Streßreaktionen von Regenbogenforellen (*Oncorhynchus mykiss*) und Plötzen (*Rutilus rutilus*). Gutachten im Auftr. d. Fischereiverbandes Nordrhein-Westfalen e. V.; Institut f. Binnenf. Potsdam-Sacrow e. V. (unpublished).
- Schreckenbach, K. & Wedekind, H., 1998. Einfluß der Angelfischerei und der Behandlung nach dem Fang auf die Streßreaktionen und die Qualität von Regenbogenforellen und Rotaugen. *Allgemeine Fischereizeitung - Fischwaid* 6: 16-18.
- Schreckenbach, K. & Wedekind, H., 1999. Belastungen von Speisefischen bei der Lebendhaltung und Auswirkungen auf die Fleischqualität. *Fischer & Teichwirt* 6: 223-224.
- Schreckenbach, K. & Wedekind, H., 2000. Einfluß der Angelfischerei und der Behandlung nach dem Fang auf die Streßreaktionen und die Fleischqualität von Regenbogenforellen (*Oncorhynchus mykiss*) und Plötzen (*Rutilus rutilus*) und Rotfedern (*Scardinius erythrophthalmus*). *Schriftenreihe Fischerei & Naturschutz* 2: 22-40.
- Schreer, J.F. et al. 2001. See Chapter 7 Cited References.
- Sobchuck, M.S. & Dawson, N., 1988. Physiological response of walleye (*Stizostedion vitreum*) to hooking/playing and hooking/holding stress. *Ontario Fisheries Technical Report Series*, 25, 15.
- Tomasso, A.O., Isely, J.J. & Tomasso Jr., J.R., 1996. Physiological response and mortality of striped bass angled in freshwater. *Transactions of the American Fisheries Society*, 125, 321-325.
- Wedekind, H. & Schreckenbach, K., 1996. Einfluss der Angelfischerei einschließlich der Behandlung nach dem Fang auf die Fleischqualität von Süßwasserfischen. Gutachten im Auftrag der Min. f. Umwelt, Raumordnung und Landw. Nordrhein-Westfalen; 74 pp.; Institut f. Binnenfischerei Potsdam-Sacrow e. V. (unpublished).
- Wilkie, M.P., Davidson, K., Brobbel, M.A., Kieffer, J.D., Booth, R.K., Bielak, A.T. & Tufts, B.L., 1996. Physiology and survival of wild Atlantic salmon following angling in warm summer waters. *Transactions of the American Fisheries Society*, 125, 572-580.
- Wydoski, R.S. et al. 1976. See Chapter 7 Cited References.
- ### 4.3 Mortality
- Barwick, D.H., 1985. See Chapter 7 Cited References.
- Beamesderfer, R.C.P. & North, J.A., 1995. Growth, natural mortality, and predicted response to fishing for largemouth bass and smallmouth bass populations in North America. *North American Journal of Fisheries Management*, 15, 688-704.
- Bennet, D.H., Dunsmoor, L.K., Rohrer, R.E. & Rieman, B.E., 1989. Mortality of tournament caught largemouth and smallmouth bass in Idaho lakes and reservoirs. *California Fish and Game*, 75, 20-26.
- Bettoli, P.W. & Osborne, R.S., 1998. See Chapter 7 Cited References.
- Burdick, B. & Wydoski, R., 1989. Effects of hooking mortality on a bluegill fishery in a western reservoir. In: Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California: 187-196
- Clapp, D.F. & Clark Jr., R.D., 1989. See Chapter 7 Cited References.
- Dedual, M., 1996. Observed mortality of rainbow trout caught by different angling techniques in Lake Taupo, New Zealand. *North American Journal of Fisheries Management*, 16, 357-363.
- Dextrase, A.J. & Ball, H.E., 1991. Hooking mortality of lake trout angled through the ice. *North American Journal of Fisheries Management*, 11, 477-479.
- Diodati, P.J., 1991. Estimating mortality of hook and released striped bass. Massachusetts Division of Marine Fisheries, Project AFC-22, Final Report, Salem.
- Dorazio, R.M., 1995. Mortality estimates of striped bass caught in Albemarle Sound and Roanoke River, North Carolina. *North American Journal of Fisheries Management*, 15, 290-299.
- Dotson, T., 1982. See Chapter 7 Cited References.
- DuBois, R.B., Margeneau, T.L., Stewart, R.S., Cunningham, P.K. & Rasmussen, P.W. 1994. Hooking mortality of northern pike angled through ice. *North American Journal of Fisheries Management*, 14, 769-775.

- Dunmall, K.M., Cooke, S.J., Schreer, J.F. & McKinley, R.S., 2001. The effect of scented lures on the hooking injury and mortality of smallmouth bass caught by novice and experienced anglers. *North American Journal of Fisheries Management*, 21, 242-248.
- Faccin, A., 1983. Hooking mortality of fly-caught Duncan River rainbow trout (*Salmo gairdneri*) in Harper Lake, British Columbia. *Fish Tech. Circ.* 58. British Columbia Fish and Wildlife.
- Falk, M.R. & Gillmann, D.V., 1975. Mortality data for angled Artic grayling and northern pike from the Great Slave Lake area, Northwest Territories. Technical Report Series CEN/D-75-1. Department of Environmental Fisheries and Marine Services, Winnipeg, Manitoba.
- Falk, M.R. *et al.* 1974. See Chapter 7 *Cited References*.
- Fielder, D.G. & Johnson, B.A., 1994. Walleye mortality during live-release tournaments on Lake Oahe, South Dakota. *North American Journal of Fisheries Management*, 14, 776-780.
- Fletcher, D.H., 1987. Hooking mortality of walleyes captured in Porcupine Bay, Washington. *North American Journal of Fisheries Management*, 7, 594-596.
- Green, D.M., Schonhoff III, B.J. & Youngs, W.D., 1989. Evaluation of hooking mortality of smallmouth and largemouth bass. Pages 229-240 in Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California.
- Hartley, R.A. & Moring, J.R., 1991. See Chapter 7 *Cited References*.
- Hartley, R.A. & Moring, J.R., 1995. Differences in mortality between largemouth and smallmouth bass caught in tournaments. *North American Journal of Fisheries Management*, 15, 666-670.
- Haw, F., 1963. Comparative hooking mortality between treble and single hooks on chinook salmon. *Pacific Marine Fisheries Commission Annual Report*, 5, 22-24.
- Hoffman, G.C., Coble, D.W., Frie, R.V., Copes, F.A., Bruch, R.M. & Kamke, K.K., 1997. Walleye and sauger mortality associated with live-release tournaments on the Lake Winnebago system, Wisconsin. *North American Journal of Fisheries Management*, 16, 364-370.
- Horton, H.F. & Wilson-Jacobs, R., 1985. A review of hooking mortality of coho (*Oncorhynchus kisutch*) and chinook (*O. tshawytscha*) salmon and steelhead trout (*Salmo gairdneri*). Portland, OR: Pacific Marine Fisheries Commission.
- Hulbert, P.J. & Engstrom-Heg, R.E., 1980. See Chapter 7 *Cited References*.
- Hunsaker, D. *et al.* P., 1970. See Chapter 7 *Cited References*.
- Hysmith, B.T., Moczygemba, J.H. & Wilde, G.R., 1994. Hooking mortality of striped bass in Lake Texoma, Texas Oklahoma. *Proceedings of the Annual Conference Southeastern Association of Fish and Wildlife Agencies*, 46, 413-420.
- Klein, W.D., 1965. Mortality of rainbow trout caught on single and treble hooks and released. *The Progressive Fish Culturist*, 27, 171-172.
- Lasater, J.E. & Haw, F., 1961. Comparative hooking mortality between treble and single hooks on silver salmon. *Pacific Marine Fisheries Communications Bulletin*, 5, 73-76.
- Loftus, A.J., 1941. An evaluation of lake trout (*Salvelinus namaycush*) hooking mortality in the upper Great Lakes. Michigan Department of Natural Resources, 15 pp.
- Loftus, A.J. *et al.* 1988. See Chapter 7 *Cited References*.
- Malchoff, M.H. & Heins, S.W., 1997. Short-term mortality of weakfish caught on single-barb hooks. *North American Journal of Fisheries Management*, 17, 477-481.
- Mason, J.W. & Hunt, R.L., 1967. Mortality rates of deeply hooked rainbow trout. *The Progressive Fish Culturist*, 29, 87-91.
- Matlock, G.C. & Dailey, J.A., 1981. Survival of hook-caught spotted seatrout held in cages. *Texas Parks and Wildlife Department, Management Data Series* 15.
- Matlock, G.C. *et al.* 1993. See Chapter 7 *Cited References*.
- May, E., 1990. An evaluation of angler induced mortality of striped bass in Maryland. AFC-18-1, Maryland Department of Natural Resources.
- Meals, K.O. & Esteban Miranda, L., 1994. Size-related mortality of tournament-caught largemouth bass. *North American Journal of Fisheries Management*, 14, 460-463.
- Milne, D.J. & Ball, E.A.R., 1956. See Chapter 7 *Cited References*.
- Mongillo, P.E., 1984. A summary of salmonid hooking mortality. Washington Department of Game, 31 pp.
- Muoneke, M.I. & Childress, W.M., 1994. Hooking mortality: A review for recreational fisheries. *Reviews in Fisheries Science*, 2, 123-157.
- Muoneke, M.I., 1992a. Hooking mortality of white crappie, *Pomoxis annularis* Rafinesque, and spotted bass, *Micropterus punctatus* Rafinesque, in Texas Reservoirs. *Aquaculture and Fisheries Management*, 23, 87-93.
- Muoneke, M.I., 1992b. Seasonal hooking mortality of bluegills caught on natural baits. *North American Journal of Fisheries Management*, 12, 645-649.

- Newman, D.L. & Storck, T.W., 1986. Angler catch, growth, and hooking mortality of tiger muskellunge in small centrarchid-dominated impoundments. *American Fisheries Society Special Publications*, 15, 346-351.
- Nuhfer, A.J. & Alexander, G.R., 1992. See Chapter 7 *Cited References*.
- Nuhfer, A.J. & Alexander, G.R., 1994. Growth, survival, and vulnerability to angling of three wild brook trout strains exposed to different levels of angler exploitation. *North American Journal of Fisheries Management*, 14, 423-434.
- Ott Jr., R.A. & Storey, K.W., 1993. See Chapter 7 *Cited References*.
- Parker, R.R. *et al.* 1959. See Chapter 7 *Cited References*.
- Pauley, G.B. & Thomas, G.L., 1993. Mortality of anadromous coastal cutthroat trout caught with artificial lures and natural bait. *North American Journal of Fisheries Management*, 13, 337-345.
- Payer, R.D. *et al.* 1989. See Chapter 7 *Cited References*.
- Pelzman, R.J., 1978. Hooking mortality of juvenile largemouth bass *Micropterus salmoides*. *California Fish and Game*, 64, 185-188.
- Persons, S.E. & Hirsch, S.A., 1994. Hooking mortality of lake trout angled through ice by jigging and set-lining. *North American Journal of Fisheries Management*, 14, 664-668.
- Rutecki, T.L. & Meyers, T.R., 1992. Mortality of juvenile sablefish captured by hand-jigging and traps. *North American Journal of Fisheries Management*, 12, 836-837.
- Rutledge, W.P. & Pritchard, D.L., 1977. Hooking mortality of largemouth bass captured by artificial lures and natural baits. Pages 103-107 in Barnhart, R.A., Roelofs, T.D., editors. *Catch-and-release fishing – a decade of experience*. Humboldt State University, Arcata, California.
- Schaefer, W.F., 1989. Hooking mortality of walleyes in a Northwestern Ontario Lake. *North American Journal of Fisheries Management*, 9, 193-194.
- Schill, D.J., 1996. Hooking mortality of bait-caught rainbow trout in a Idaho trout stream and a hatchery: implications for special-regulation management. *North American Journal of Fisheries Management*, 16, 348-356.
- Schill, D.J. *et al.* 1986. See Chapter 7 *Cited References*.
- Schisler, G.J. & Bergersen, E.P., 1996. Postrelease hooking mortality of rainbow trout caught on scented artificial baits. *North American Journal of Fisheries Management*, 16, 570-578.
- Schramm, H.L., Haydt Jr., P.J. & Portier, K.M., 1987. Evaluation of prerelease, postrelease, and total mortality of largemouth bass during tournaments in two Florida lakes. *North American Journal of Fisheries Management*, 7, 394-402.
- Schwalme, K. & Mackay, W.C., 1985. See Chapter 7 *Cited References*.
- Taylor, M.J. & White, K.R., 1992. A meta-analysis of hooking mortality of nonanadromous trout. *North American Journal of Fisheries Management*, 12, 760-767.
- Titus, R.G. & Vanicek, C.D., 1988. See Chapter 7 *Cited References*.
- Warner, K. & Johnson, P.R. 1978. Mortality of landlocked Atlantic salmon hooked on four types of fishing gear at the hatchery. *The Progressive Fish Culturist*, 41, 99-102.
- Warner, K., 1979. See Chapter 7 *Cited References*.
- Wertheimer, A., 1988. See Chapter 7 *Cited References*.
- Wilde, G.R., 1998. Tournament associated mortality in black bass. *Fisheries*, 23, 12-22.
- Wydoski, R.S. *et al.* 1976. See Chapter 7 *Cited References*.
- #### 4.4 Retention in keepnets/confinement
- Cooke, S.J. & Hogle, W.J., 2000. See Chapter 7 *Cited References*.
- Drossé, H., 1991. Lebendhaltung gefangener Fische im Setzkescher ist Tierquälerei. *Monatsschrift Deutsches Recht*, 45, 278.
- Drossé, H., 1992. Die Lebendhaltung gefangener Fische im Setzkescher aus tierschutzrechtlicher Sicht. *Fischökologie Aktuell*, 6, 16-24.
- Drossé, H., 1994. Die Lebendhaltung gefangener Fische im Setzkescher. *Monatsschrift Deutsches Recht*, 242-245.
- Harsányi, A., 1992. Anwendung des Setzkeschers in der Fischerei. *Österreichs Fischerei*, 45, 18-29.
- Hartley, R.A. & Moring, J.R., 1993. See Chapter 7 *Cited References*.
- Kossmann, H. & Pfeiffer, W., 1996. *Fische im Setzkescher: Eine Verhaltensstudie zur Haltung von geangelten Fischen*. Ockenheim: Verband Deutscher Sportfischer e.V., Landesverband Rheinland-Pfalz, 6 p.
- Pankhurst, N.W. & Sharples, D.F., 1992. Effects of capture and confinement on plasma cortisol concentrations in the snapper, *Pagrus auratus*. *Australian Journal of Marine and Freshwater Research*, 43, 345-356.
- Plumb, J.A. *et al.*, 1988. See Chapter 7 *Cited References*.
- Pottinger, T.G. 1997. See Chapter 7 *Cited References*.
- Pottinger, T.G., 1997. See Chapter 7 *Cited References*.
- Pottinger, T.G., 1998. See Chapter 7 *Cited References*.

Raat, A.J.P. *et al.* 1997. See Chapter 7 *Cited References*.

Schulz, D., 1992. Tierschutzrelevante untersuchungen zur Lebendhaltung gefangener Rotaugen im Setzkescher. *Fischökologie Aktuell*, 6, 2-13.

4.5 Other

Anderson, W.G. *et al.* 1998. See Chapter 7 *Cited References*.

Beukema, J.J., 1969. Angling experiments with carp (*Cyprinus carpio* L.) I. Differences between wild, domesticated, and hybrid strains. *Netherlands Journal of Zoology* 19, 596-609.

Beukema, J.J., 1970a. See Chapter 7 *Cited References*.

Beukema, J.J., 1970b. See Chapter 7 *Cited References*.

Beukema, J.J., 1974. Experimental test of a basic assumption of capture-recapture method in pond populations of carp *Cyprinus carpio* L. *Journal of Fish Biology*, 6, 317-329.

5. References for "Aquaculture"

5.1 Handling

Aabel, J.P., 1991. Stress of Atlantic salmon caused by handling and grading. *Aquaculture and the Environment*, 14:1.

Barton, B.A., 2000. See Chapter 7 *Cited References*.

Barton, B.A., Bollig, H., Hauskins, B.L. & Jansen, C.R., 2000. Juvenile pallid (*Scaphirhynchus albus*) and hybrid pallid x shovelnose (*S.albus x platyrhynchus*) sturgeons exhibit low physiological responses to acute handling and severe confinement. *Comparative Biochemistry and Physiology*, 126, 125-134.

Barton, B.A., Zitzow, Z.E., 1995. Physiological responses of juvenile walleyes to handling stress with recovery in saline water. *The Progressive Fish-Culturist*, 57, 267-276.

Berka, R., 1986. See Chapter 7 *Cited References*.

Cech, J.J., Bartholow, S.D., Young, P.S., Hopkins, T.E., 1996. Striped bass exercise and handling stress in freshwater: physiological responses to recovery environment. *Transactions of the American Fisheries Society*, 125, 308-320.

Cleary, J.J., Pankhurst, N.W. & Battaglione, S.C., 2000. The effect of capture and handling stress on plasma steroid levels and gonadal condition in wild and farmed snapper *Pagrus auratus* (Sparidae). *Journal of the World Aquaculture Society*, 31, 558-569.

Cubero, L. & Molinero, A., 1997. Handling, confinement and anaesthetic exposure induces changes in the blood and tissue immune characteristics of gilthead sea bream. *Diseases of Aquatic Organisms*, 31, 89-94.

Coble, D.W., 1988. Effects of angling on bluegill populations: management implications. *North American Journal of Fisheries Management*, 9, 81-85.

Drossé, H., 1990. Tierquälerei beim Wettangeln. *Neue Zeitschrift. Strafrecht*, 10, 72-73.

Medway, Lord 1980. *Report of the panel of enquiry into shooting and angling (1976-1979)*. Panel of enquiry into shooting and angling, Horsham, Sussex, UK: Royal Society of the Prevention of Cruelty to Animals.

Raat, A.J.P., 1985. See Chapter 7 *Cited References*.

Raat, A.J.P., 1987. Effects of crowding and availability of food on growth and angling vulnerability in the two sexes of scaled and mirror patterned carp (*Cyprinus carpio* L.). *Netherlands Journal of Zoology*, 37, 1-25.

Raat, A.J.P., 1991a. Production, growth, condition and angling vulnerability of zander, *Stizostedion lucioperca* (L.), in relation to the availability of prey fish in ponds. *Aquaculture and Fisheries Management*, 22, 93-104.

Davis, L.E. & Schreck, C.B., 1997. The energetic response to handling stress in juvenile coho salmon. *Transactions of the American Fisheries Society*, 126, 248-258.

Foo, J.T.W. & Lam T.J., 1993. Serum cortisol response to handling stress and the effect of cortisol implantation on testosterone level in the tilapia, *Oreochromis mossambicus*. *Aquaculture*, 115, 145-158.

Frisch, A.J. & Anderson, T.A., 2000. The response of coral trout (*Plectropomus leopardus*) to capture, handling and transport and shallow water stress. *Fish Physiology and Biochemistry*, 23, 23-34.

Grutter, A.S. & Pankhurst, N.W., 2000. See Chapter 7 *Cited References*.

Kreiberg, H. 1994. A survey of animal care considerations for fish handling. *Humane Innovations and Alternatives*, 8, 592-595.

Kristjansson, B.A., Hardarson, Th., Skulason, S. & Sveinsson, T., 1995. Effects on growth of Arctic charr of handling, anaesthesia and repeated formalin bathing shortly after first erogenous feeding. *Journal. of Fish Biology*, 46, 163-165.

Melotti, P., Roncarati, A., Garella, E., Carnevali, O., Mosconi, G. & Polzonetti-Magni, A., 1992. Effects of handling and capture stress on plasma glucose, cortisol and androgen levels in brown trout (*Salmo trutta morpha fario*). *Journal of Applied Ichthyology*, 8, 234-239.

Reubush, K.J., Heath, A.G., 1996. Metabolic responses to acute handling by fingerlings inland and anadromous striped bass. *Journal of Fish Biology*, 49, 830-841.

- Rotllant, J. & Tort, L., 1997. See Chapter 7 *Cited References*.
- Thomas, P. & Robertson, L., 1991. Plasma cortisol and glucose stress responses of red drum (*Scianops ocellatus*) to handling and shallow water stressors and anesthesia with MS-222, quinaldine sulfate and metomidate. *Aquaculture* 96, 69-86.
- Waring, C.P., Stagg, P.M. & Poxton, M.G. 1992. The effect of handling on flounder (*Platichthys flesus* L.) and Atlantic salmon (*Salmo salar* L.). *Journal of Fish Biology* 41, 131-144.
- Waring, C.P. *et al.* 1996. See Chapter 7 *Cited References*.
- Wedemeyer, G.A., 1976. Physiological response of juvenile coho salmon (*Oncorhynchus kisutch*) and rainbow trout (*Salmo gairdneri*) to handling and crowding stress in intensive fish culture. *Journal of the Fisheries Research Board of Canada*, 33, 2699-2702.
- Wolffrom, T., 2004. See Chapter 7 *Cited References*.
- ## 5.2 Slaughter
- Anonymous, 2002. See Chapter 7 *Cited References*.
- Azam K. *et al.* 1989. See Chapter 7 *Cited References*.
- Azam, K., Mackie, I.M. & Smith, J., 1990. Effect of stunning method on the time of onset, duration and resolution of rigor in rainbow trout (*Salmo gairdneri*) as measured by visual observation and analyses for lactic acid, nucleotide-degradation products and glycogen. *Proceedings of the meeting of the Commission C2 Chilling and freezing of new fish products*. IIF Paris France Institut International du Froid 1990 pp. 351-357.
- Berg T. *et al.* 1997. See Chapter 7 *Cited References*.
- Bernoth, E.M., Wormuth, H.J., 1990. Tierschutzaspekte bei der Tötung von Fischen. *Deutsche tierärztliche Wochenschrift*, 97, 154-157.
- Bretzinger, Ch., 2001. Influence of different pre-slaughter stunning methods on stress reaction and product quality of rainbow trout (*Oncorhynchus mykiss*). Muenchen-FRG Hieronymus, 189 pp.
- Byrne, H., 2002. See Chapter 7 *Cited References*.
- Einen, O., Waagan, B. & Thamassen, M.S., 1998. Starvation prior to slaughter in Atlantic salmon (*Salmo salar*) - Effects on weight loss, body shape, slaughter- and fillet- yield, proximate and fatty acid composition. *Aquaculture*, 166, 85-104.
- Erikson, U., 2000. Potential effects of preslaughter fasting, handling and transport. In: *Farmed Fish Quality*. Kestin, S.C. and Warriss, P.D. (eds.). Blackwell Science: Oxford.
- Erikson, U., Sigholt, T., Rustad, T., Einarsdottir, I.E. & Jorgensen, L., 1999. Contribution of bleeding to total handling stress during slaughter of Atlantic salmon. *Aquaculture International*, 7,101-115.
- Council Directive 93/119/EC, 1993 on the protection of animals at the time of slaughter or killing. *Official Journal of the European Communities* No 1.340/21.
- Council Directive 98/37/EC, 1993 on the approximation of the laws of the Member States relating to machinery. *Official Journal of the European Communities*, L207, 1-46.
- Cooke, M. (2001): Ethical considerations for the production of farmed fish- the retailer's viewpoint In: *Farmed Fish Quality*, S.C. Kestin & P.D. Warriss (eds.), Blackwell, Oxford, UK, pp. 116-119.
- Croft, P.G., 1952. The effect of electrical stimulation of the brain on the perception of pain. *Journal Mental Science* 42, 421-426.
- FAWC, 1996. *Report on the welfare of fish*, PB 2765, MAFF, UK
- Flight, W.G.F. & Verheijen, F.J., 1993. The 'neck-cut' (spinal transaction): Not a humane way to slaughter eel, *Anguilla anguilla* (L.). *Aquaculture and Fisheries Management*, 24, 523-528.
- Horrex, H., 1999. Humane Slaughtering. *Fish Farmer*, September/October 1999. pp. 42.
- Keene, J.L., Noakes, D.L.G., Moccia, R.D. & Soto C.G., 1998. The efficacy of clove oil as an anaesthetic for rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquaculture Research*, 29, 89-101.
- Kestin, S.C., Van de Vis, J.W. & Robb, D.F.H., 2001. A simple protocol for assessing brain function in fish and the effectiveness of stunning and killing methods used on fish. *Veterinary Record*, 150, 320-307.
- Kestin, S.C., Van de Vis, J.W. & Robb, D.H.F., 2002. Protocol for assessing brain function in fish and the effectiveness of methods used to stun and kill them. *Veterinary Record*, 150, 302-307.
- Kestin S.C., Wotton S.B. & Gregory N.G., 1991. Effect of slaughter by removal from water on visual evoked activity in the brain and reflex movement of rainbow trout (*Oncorhynchus mykiss*). *Veterinary Record* 128, 443-446.
- Kestin, S.C. *et al.* 1995. See Chapter 7 *Cited References*.
- Kuhlmann, H., Münkner, W., 1996. Gutachterliche Stellungnahme zum tierschutzgerechten Betäuben Töten von Aalen in grösseren Menge. *Fischer und Teichwirt*, 47, 493-495.
- Kuhlmann, H., Muenker, W., Oehlenschlaeger, J. & van de Vis, H., 2001. On the stunning and slaughtering of eels (*Anguilla anguilla*) in

- accordance with animal welfare. Inf. Fischwirtsch. *Fischereiforsch.*, 48, 2: 82-89.
- Kuhlmann, H., Münkner, W., Van de Vis, J.W., Oehlenschläger J., Koch, M., 2000. Untersuchungen zur anästhesierenden Wirkung von Aqui-S und chemisch verwandten Verbindungen beim Aal (*Anguilla anguilla*). *Archiv für Lebensmittelhygiene*, Hannover, 51, 60-62
- Lambooij, E., Kloosterboer, R.J., Gerritzen, M, C., Veldman, M. & Van de Vis, J.W., 2004. Feasibility of electrical stunning followed by decapitation or chilling of African catfish (*Clarias gariepinus*) assessment of behavioural, neural and product quality parameters. *Submitted for publication in Aquaculture Research*
- Lambooij, E., Van de Vis, J.W., Kloosterboer & R.J., Pieterse, C., 2000. Assessment of electrical and mechanical stunning of farmed eel (*Anguilla anguilla*, L.). *Proc. I.C.o.M.S.T.* 370-371.
- Lambooij, E., Van de Vis, J.W., Kloosterboer, R.J., Pieterse, C., 2001. Ontwikkeling protocol voor beoordeling bedwelmings- en dodingsprocessen bij de paling (*Anguilla anguilla*, L.). RIVO Rapportnummer C063/01.
- Lambooij, E., van de Vis, J.W., Kloosterboer, R.J., & Pieterse, C., 2002b. Welfare aspects of live chilling and freezing of farmed eel (*Anguilla anguilla*, L.): neural and behavioural assessment. *Aquaculture* 210, 159-169.
- Lambooij, E. *et al.* 2002. See Chapter 7 *Cited References*.
- Lines, J.A., Robb., D.H, Kestin, S.C. , Crook, S.C. & Benson, T., 2003. Electric stunning: a humane slaughter method for trout. *Aquacultural Engineering*, 28, 141-154.
- Lymbery, P, 2002. *In too deep - The welfare of intensively farmed fish*. CIWF Trust, Hampshire, UK, 56 pp.
- Magnussen, O.M. & Johansen, S., 1990. Influence of handling and slaughtering on quality of farmed fish. *Proceedings of the meeting of the Commission C2 Chilling and freezing of new fish products*. IIF Paris France Institut International du Froid 1990 pp. 259-264.
- Morzel, M. & Van de Vis, J.W., 2003. See Chapter 7 *Cited References*.
- Morzel M. *et al.* 2003. See Chapter 7 *Cited References*.
- Oehlenschläger, J., Kestin, S.C., Tejada, M., Sorensen, N.K., Torrissen, O.J., Nesvabda, P. & Van Rijnsingen, L.C.M., 1998. Optimisation of harvest procedures of farmed fish with respect to quality and welfare. *Third European Marine Science and Technology Conference MAST Conference*, Lisbon, 23-27 May 1998.
- Parisi, G., Mecatti, M., Lupi, P., Scappini, F. & Poli, B.M., 2002. Comparison of five slaughter methods for European seabass. Changes in isometric contraction force and pH during the first 24 hours post mortem. *Aquaculture Europe 2002, European Aquaculture Society Special Publication 32*.
- Poli, B.M. *et al.* 2002. See Chapter 7 *Cited References*.
- RDA, 2003. [Criteria for killing of eel and African catfish] (*in Dutch*). Advice prepared by the Council of Animal Affairs for the preparation of legislation by the Dutch Ministry of Agriculture, Nature Management and Food Quality. Advies RDA2003/05, P.O.Box 90428, 2509 LK The Hague, The Netherlands, 28 pp.
- Robb, D.F.H. & Kestin, S.C., 2002. See Chapter 7 *Cited References*.
- Robb D.H.F. *et al.* 2000. See Chapter 7 *Cited References*.
- Roth, B. *et al.*, 2002. See Chapter 7 *Cited References*.
- Robb, D.F.H., O'Callaghan, M. Lines, J.A. & Kestin, S.C., 2002. Electrical stunning of rainbow trout (*Oncorhynchus mykiss*): factors that affect stun duration. *Aquaculture*, 205, 359-371.
- Robb, D.F.H. & Roth, B., 2002. See Chapter 7 *Cited References*.
- Robb D.H.F. *et al.* 2000. See Chapter 7 *Cited References*.
- Robb, D.F., Wotton, S.B. & Van de Vis, J.W., 2002. Preslaughter electrical stunning of eels. *Aquaculture Research*, 33, 37-42.
- Robb, D.H.F., 2000. *Humane Slaughter*. Paper presented at British Trout Farming conference, Sparsholt, Hampshire. September 2000.
- Robb, D.H.F., Kestin, S.C. & Lines, J.A., 2000. Progress with humane slaughter. *Fish Farmer*, Nov/Dec. 2000, pp. 41.
- Sigurisdottir S. *et al.* 2001. See Chapter 7 *Cited References*.
- Simmons, N.J., 1989. Electrical stunning of pigs. The effect of different stunning methods on the incidence of blood splash, broken bones and PSE occurrence. MSc Thesis, University of Bristol, UK, p.9
- Skjervold, P.O. *et al.* 2001. See Chapter 7 *Cited References*.
- Skjervold, P.O., Fjaera, S.O. & Oestby, P.B., 1999. Rigor in Atlantic salmon as affected by crowding stress prior to chilling before slaughter. *Aquaculture*, 175, 93-101.
- Solomon, M.B., Van Laack, R.L.J.M and Eastridge, J.S., 1998. Biophysical basis of pale, soft and exudative (PSE) pork and poultry muscle a review. *Journal of Muscle Foods*, 9, 1-11.
- TierSchlV, 1997. Verordnung zum Schutz von Tieren in Zusammenhang mit der Schlachtung und Tötung (Tierschutz-Schlachtverordnung) vom 3-3-1997, *BGB I*, Nr. 13, 6-3-1997.

- TierSchlV, 1999. Verordnung zur Änderung der Tierschlachtung verordnung vom 25-11-1999, *BGBI*, Nr. 54, 10-12-1999.
- Undeland, I., 2001. Lipid oxidation in fatty fish during processing and storage. In *Farmed Fish Quality* (eds. S.C. Kestin and P.D. Warriss), Blackwell, Oxford, UK, pp. 261-275.
- Van de Vis, J.W., 1997. Effect of commercial slaughter methods on quality and welfare of eels. In: *Welfare of Fish at Slaughter*. Workshop Minutes. Ed: Robb, D. School of Veterinary Science, Langford, Bristol.
- Van de Vis, J.W., Lambooi, E., Pieterse, C. & Kloosterboer, R.J., 2001. A method to assess unconsciousness and insensitivity in eel (*Anguilla anguilla*, L.) for development of humane slaughter methods. *Ann. Soc. Sci. Faeroensis* Suppl. XXVII, 181-186.
- Van de Vis, J.W., Oehlenschlaeger, J., Kuhlmann, H., Muenker, W., Robb, D.H.F. & Schelvis-Smit, A.A.M., 1999. Effect of the commercial and experimental slaughter of eels (*Anguilla anguilla*) on quality and welfare. *Farmed Fish Quality* (eds. S.C. Kestin and P.D. Warriss), Blackwell Science, 2001, pp 234-248.
- Van de Vis, J.W., 2001. See Chapter 7 *Cited References*.
- Van de Vis, J. W., Oehlenschläger, J., Kuhlmann, H., Münkner, W., Robb, D.F.H., & Schelvis-Smit, A.A.M. , 2001. Commercial and experimental slaughter of eel (*Anguilla anguilla*, L.): effect on quality and welfare. In: *Farmed Fish Quality* (eds. S.C. Kestin and P.D. Warriss), Blackwell, Oxford, UK, pp. 234-257.
- Van de Vis, J.W. *et al.* 2003a. See Chapter 7 *Cited References*.
- Van de Vis, J.W. *et al.* 2003b. See Chapter 7 *Cited References*.
- Van de Vis, J.W., Lambooi, E., Kloosterboer, R.J., Gerritzen, M.A. & Pieterse, C., 2002. Criteria for assessment of slaughter methods of eel (*Anguilla anguilla*) and African catfish (*Clarias gariepinus*). Oral presentation at the 32nd West European Fish Technologists' Association Meeting, Ireland, May 2002.
- Verheijen, F.J. & Flight, W.F.G., 1997. Decapitation and brining: experimental tests show that after these commercial methods for slaughtering of eel, *Anguilla anguilla* L., death is not instantaneous. *Aquaculture Research*, 28, 361-366.
- ### 5.3 Stocking density
- Barcellos, L.J.G., Nicolaiewsky, S., de Souza, S.M.G. & Lulhier, F., 1999. The effects of stocking density and social interaction on acute stress response in Nile tilapia *Oreochromis niloticus* (L.) fingerlings. *Aquaculture Research*, 30, 887-892.
- Barnett, C.W. & Pankhurst, N.W., 1998. See Chapter 7 *Cited References*.
- Clearwater, S.J. & Pankhurst, N.W., 1997. See Chapter 7 *Cited References*.
- El-Sayed, A.F.M., 2001. Intensive culture of Nile tilapia *Oreochromis niloticus* fry: Effects of stocking density and feeding levels on fish performance. *Aquaculture 2001: Book of Abstracts* 143 JM Parker Coliseum Louisiana State University Baton Rouge LA 70803 USA World Aquaculture Society 2001 206.
- Ewing, R.D. & Ewing, S.K. 1995. Review of the effects of rearing density on the survival to adulthood for Pacific salmon. *The Progressive Fish-Culturist*, 57, 1-25.
- Hossain, M.A.R., Beveridge M.C.M., & Haylor G.S., 1998. The effects of density, light and shelter on the growth and survival of African catfish (*Clarias gariepinus* Burchell, 1822) fingerlings. *Aquaculture* 160, 251-258.
- Jørgensen, E.H. *et al.* 1993. See Chapter 7 *Cited References*.
- Mazur, C.F. & Iwama, G.K., 1993. See Chapter 7 *Cited References*.
- Montero, D. *et al.* 1999. See Chapter 7 *Cited References*.
- Ortuño, J. *et al.* 2001. See Chapter 7 *Cited References*.
- Pickering, A.D. & Stewart, A., 1984. See Chapter 7 *Cited References*.
- Procarione, L.S. *et al.* 1999. See Chapter 7 *Cited References*.
- Ross R.M. & Watten B.J., 1998. Importance of rearing-unit and stocking density to the behavior, growth and metabolism of lake trout (*Salvelinus namaycush*). *Aquacultural Engineering*, 19, 41-56.
- Ruane, N.M., Carballo, E.C. & Komen, J. 2002. Increased stocking density influences the acute physiological stress response of common carp (*Cyprinus carpio* L.). *Aquaculture Research*, 33, 777-784.
- Skjervold, P.O. *et al.* 2001. See Chapter 7 *Cited References*.
- Tort, L. *et al.* 1996. See Chapter 7 *Cited References*.
- Vijayan, M.M. *et al.* 1990. See Chapter 7 *Cited References*.
- Vijayan, M.M. & Leatherland, J.F., 1988. Effect of stocking density on the growth and stress-response in brook charr, *Salvelinus fontinalis*. *Aquaculture*, 75, 159-170.
- Vijayan, M.M. *et al.* 1997. See Chapter 7 *Cited References*.
- Yin, Z. *et al.* 1995. See Chapter 7 *Cited References*.
- ### 5.4 Physiological effects/ stress
- Adams, M.S., 1990. Status and use of biological indicators for evaluating the effects of stress

- on fish. *American Fisheries Society Symposium*, 8, 1-8.
- Adams, S.M. (ed.), 1980. *Biological indicators of stress in fish*. Bethesda: American Fisheries Society. *American Fisheries Society Symposium*, 8, 191 p.
- Asterita, M.F., 1985. *The physiology of stress*. Human Sciences Press Inc., New York, p. 264.
- Barcellos, L.J.G., Nicholaiewsky S., Souza S.M.G. & Lulhier F. 1999. Plasmatic levels of cortisol in the response to acute stress in Nile tilapia, *Oreochromis niloticus* (L.), previously exposed to chronic stress. *Aquaculture Research*, 30, 437-444.
- Barton B.A., 1997. Stress in finfish: past, present and future - a historical perspective. In: *Fish Stress and Health in Aquaculture* (ed. by G.K. Iwama, A.D. Pickering, J.P. Sumpter & C.B. Schreck), pp.1-33. Society for Experimental Biology; seminar series 62. Cambridge University Press, Cambridge.
- Barton B.A. & Schreck, C.B., 1987. Metabolic cost of acute physical stress in juvenile steelhead. *Transactions of the American Fisheries Society*, 116, 257-263.
- Barton, B.A. & Iwama G.K., 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases*, 3-16.
- Brown, J.A., Edwards D.J. & Whitehead, C., 1989. Cortisol and thyroid hormone responses to acid stress in the brown trout. *Journal of Fish Biology*, 35, 73-84.
- Campbell, P.M., Pottinger, T.G., Sumpter, J.P., 1992. Stress reduces the quality of gametes produced by rainbow trout. *Biology of Reproduction*, 47, 1140-1150.
- Chrousos, C.P. & Gold, P.W., 1992. The concepts of stress and stress system disorders. Overview of physical and behavioral homeostasis. *JAMA*, 267, 1244-1252.
- Gamperl, A.K., Vijayan, M.M. & Boutilier, R.G., 1994. Experimental control of stress hormone levels in fishes: techniques and applications. *Reviews in Fish Biology and Fisheries*, 4, 215-255.
- Goede, R.W. & Barton, B.A., 1990. Organismic indices and an autopsy-based assessment as indicators of health and condition of fish. In: *Biological Indicators of Stress in Fish*, Adams, S.M. (ed.). American Fisheries Society Symposium 8, Bethesda, Maryland, 93-108.
- Gregory, T.R.W.C.M., 1999. The effects of chronic plasma cortisol elevation on the feeding behaviour, growth, competitive ability, and swimming performance of juvenile rainbow trout. *Physiological and Biochemical Zoology*, 72, 286-295.
- Harbuz, M.S. & Lightman, S.J., 1992. Stress and the hypothalomo-pituitary-interrenal axis: acute, chronic and immunological activation. *Journal of Endocrinology*, 134, 327-329.
- Harrell, R.M., 1992. Stress mitigation by use of salt and anesthetic for wild striped bass captured for brood stock. *The Progressive Fish Culturist*, 54, 228-233.
- Iger, Y., Balm, P.H.M., Jenner, H.A. & Wendelaar Bonga, S.E., 1995. Cortisol induces stress-related changes in the skin of rainbow trout (*Oncorhynchus mykiss*). *General Comparative Endocrinology*, 97, 188-198.
- Iwama, G.K., Morgan, J.D. & Barton, B.A., 1995. Simple field methods for monitoring stress and general condition of fish. *Aquaculture Research*, 26, 273-282.
- Iwama, G.K., Pickering, A.D., Sumpter, J.P. & Schreck, C.B., 1997. *Fish stress and health in aquaculture*. Society for Experimental Biology, Seminar series 62. Cambridge: Cambridge University Press. 278 p.
- Jeney, G., Galeotti, M., Volpatti, D., Jeney, Z. & Anderson, D., 1997. Prevention of stress in rainbow trout (*Oncorhynchus mykiss*) fed diets containing different doses of glucan. *Aquaculture*, 154, 1-15.
- Levine, S., 1985. A definition of stress? In: *Animal Stress*, Moberg, G.P. American Physiological Society, Bethesda, Maryland, p. 51-69.
- Lines, J.A. & Frost, A.R., 1999. Review of opportunities for low stress and selective control of fish. *Aquacultural Engineering*, 20, 211-230.
- Mazeaud, M.M., Mazeaud, F. & Donaldson E.M., 1977. Primary and secondary effects of stress in fish: Some new data with a general review. *Transactions of the American Fisheries Society*, 106, 201-212.
- McDonald, D.G. & Robinson, J.G., 1993. Physiological responses of lake trout to stress: effects of water hardness and genotype. *Transactions of the American Fisheries Society*, 122, 1146-1155.
- Moberg, G.P., 1985. Animal stress. In: *Animal Stress*, Moberg, G.P. American Physiological Society, Bethesda, Maryland. p. 324.
- Olla, B.L., Davis, M.W. & Schreck, C.B. 1995. Stress-induced impairment of predator evasion and non-predator mortality in pacific salmon. *Aquaculture Research*, 26, 393-398.
- Olla, B.L. *et al.* 1992. See Chapter 7 *Cited References*.
- Pagnotta, A., Brooks, L. & Milligan, L., 1994. The potential regulatory roles of cortisol in recovery from exhaustive exercise in rainbow trout. *Canadian Journal of Zoology*, 72, 2136-2146.

- Peters, G., 1991. *Stressoren und Stresseffekte in der Fischintensivhaltung*. Deutsche Veterinar Gesellschaft. KTBL-Schrift, 344, 80-90.
- Peters, G. *et al.* 1980. See Chapter 7 *Cited References*.
- Peters, G., Klinger, H. & Delventhal, H. 1983. Diagnosis of stress in the European eel. *Rapports Proces-verbal Réunion Conseil international Exploration Mer* 182, 79-86.
- Pickering, A.D., 1981. *Stress and fish*. London: Academic Press, 367 p.
- Pickering, A.D., 1981. Introduction: The concept of biological stress. In: *Stress and Fish*, Pickering, A.D. Academic Press, London, p. 1-9.
- Pickering, A.D., 1993. Growth and stress in fish production. *Aquaculture*, 111, 51-63.
- Pickering, A.D., 1993. Stress and adaptation: husbandry and stress. In: *Recent Advances in Aquaculture IV*. Muir, J.F. and R.J. Roberts (Eds.), Blackwell Scientific Publications, London, p. 155-169.
- Pickering, A.D., 1998. See Chapter 7 *Cited References*.
- Pickering, A.D. & Pottinger, T.G., 1989. Stress responses and disease resistance in salmonid fish: effects of chronic elevation of plasma cortisol. *Fish Physiology and Biochemistry*, 7, 253-258.
- Pickering, A.D., Pottinger, T.G. & Christie P., 1982. Recovery of the brown trout, *Salmo trutta* L., from acute handling stress: a time-course study. *Journal of Fish Biology*, 20, 229-244.
- Pottinger, T.G. & Pickering, A.D., 1997. Genetic basis to the stress response: selective breeding for stress-tolerant fish. In: *Fish Stress and Health in Aquaculture* (ed. by G.K. Iwama, A.D. Pickering, J.P. Sumpter & C.B. Schreck), pp. 171-193. Society for Experimental Biology; seminar series 62. Cambridge University Press, Cambridge.
- Rice, J.A., 1990. Bioenergetics modeling approaches to evaluation of stress in fishes. In: *Biological Indicators of Stress in Fish*, Adams, S.M. (Ed.) American Fisheries Society Symposium 8, Bethesda, Maryland, p. 80-92.
- Robertson, L., Thomas, P., Arnold, C.R. & Trant, J.M., 1987. Plasma cortisol and secondary stress responses of red drum (*Scianops ocellatus*) to handling, transport, rearing, density and a disease outbreak. *The Progressive Fish Culturist*, 49, 1-12.
- Ruane, N.M., 2002. See Chapter 7 *Cited References*.
- Ruane, N.M., Wendelaar Bonga, S.E. & Balm, P.H.M., 1999. Differences between rainbow trout and brown trout in the regulation of the pituitary-interrenal axis and physiological performance during confinement. *General and Comparative Endocrinology*, 115, 210-219.
- Rueda, P.A., 2004. Towards assessment of welfare in African catfish, *Clarias gariepinus*: the first step. PhD thesis, Fish Culture & Fisheries Group, Wageningen Institute for Animal Sciences, Wageningen University, The Netherlands.
- Scherer, E., 1992. Behavioural responses as indicators of environmental alterations: approaches, results, developments. *Journal of Applied Ichthyology*, 8, 122-131.
- Schreck, C.B., 1981. Stress and compensation in teleostean fishes: response to social and physical factors. In: *Stress and Fish* (ed. by A.D. Pickering), pp. 295-321. Academic Press Inc., New York.
- Schreck, C.B., 1982. Stress and rearing of salmonids. *Aquaculture*, 28, 241-249.
- Schreck, C.B., 1990. See Chapter 7 *Cited References*.
- Schreck, C.B., 2000. Accumulation and long-term effects of stress in fish. Pp 147-158 in *The Biology of Animal Stress*. Eds. G.P. Moberg & J.A. Mench. CABI Publishing.
- Schreck, C.B. *et al.*, 2001. See Chapter 7 *Cited References*.
- Schreck, C.B., Olla, B.L. & Davis, M.W., 1997. Behavioural responses to stress. In: *Fish Stress and Health in Aquaculture* (ed. by G.K. Iwama, A.D. Pickering, J.P. Sumpter & C.B. Schreck), pp. 145-170. Society for Experimental Biology; seminar series 62. Cambridge University Press, Cambridge.
- Sharpe, C.S., Thompson, D.A., Blankenship, H.L. & Schreck, C.B., 1997. Effects of routine handling and tagging procedures on physiological stress responses in juvenile chinook salmon. *The Progressive Fish-Culturist*, 60, 81-87.
- Sniezsko, S.F., 1974. The effects of environmental stress on outbreaks of infectious diseases of fishes. *Journal of Fish Biology*, 6, 197-208.
- Strange, R.J., Cech, J.J., 1992. Reduced swimming performance of striped bass after confinement stress. *Transactions of the American Fisheries Society*, 121, 206-210.
- Strange, R.J., Schreck, C.B., 1978. Anesthetic and handling stress on survival and cortisol concentration in yearling chinook salmon (*Oncorhynchus tshawytscha*). *Journal of the Fisheries Research Board of Canada*, 35, 345-349.
- Sumpter, J.P., 1997. The endocrinology of stress. In: *Fish Stress and Health in Aquaculture* (ed. by G.K. Iwama, A.D. Pickering, J.P. Sumpter & C.B. Schreck), pp. 95-118. Society for Experimental Biology; seminar series 62. Cambridge University Press, Cambridge.

- Thomas, P., 1990. Molecular and biochemical responses of fish to stressors and their potential use in environmental monitoring. In: *Biological Indicators of Stress in Fish*, Adams, S.M. (Ed.) *American Fisheries Society Symposium* 8, Bethesda, Maryland, p. 9-28.
- Van Raaij, M.I.M., Pit, D.S.S., Balm, P.H.M., Steffens, A.B. & Van den Thillart, G., 1996. Behavioral strategy and the physiological stress response in rainbow trout exposed to severe hypoxia. *Hormones and Behavior* 30, 85-92.
- Vijayan, M.M. & Moon, T.W., 1992. Acute handling stress alters hepatic glycogen metabolism in food deprived rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 2260-2266.
- Wardle, C.S., 1981. Physiological stress in captive fish. In: *Aquarium Systems*, Hawkins, A.D. (Ed.), Academic Press, London, p. 403-414.
- Wedemeyer, G.A., 1996. See Chapter 7 *Cited References*.
- Wedemeyer, G.A. & McLeay, D.J., 1981. Methods for determining the tolerance of fishes to environmental stressors. In: *Stress and Fish* (ed. by A.D. Pickering), pp. 247-275. Academic Press Inc., New York.
- Weyts, F.A.A. *et al.* 1999. See Chapter 7 *Cited References*.
- 5.5 Tagging**
- Davenport, J. *et al.* 2002. Chapters 6 (Legislation) & 7 (Fish welfare and health in relation to tagging) In: Thorsteinsson V. (coord.) 2002. *Tagging Methods for Stock Assessment and Research in Fisheries. Report of Concerted Action FAIR CT.96.1394 (CATAG)*. Reykjavik. Marine Research Institute Technical Report No. 79. pp 183.
- Jepsen, N., Davis, L.E., Schreck, C.B. & Siddens, B., 2001. The physiological response of chinook salmon to two methods of radio-tagging. *Transactions of the American Fisheries Society*, 130, 3: 495-500.
- Sharpe, C.S., Thompson, D.A., Blankenship, H.L. & Schreck, C.B., 1998. Effects of routine handling and tagging procedures on physiological stress responses in juvenile chinook salmon. *The Progressive Fish-Culturist*, 60, 81-87.
- Thorsteinsson, V. 2002. See Chapter 7 *Cited References*.
- 5.6 Transportation**
- Bandein, J. & Leatherland, J.F., 1997. See Chapter 7 *Cited References*.
- Barnett, C.W. & Pankhurst, N.W., 1998. See Chapter 7 *Cited References*.
- Barton, B.A., 2000. See Chapter 7 *Cited References*.
- Barton, B.A. & Peter, R.E., 1982. Plasma cortisol stress response in fingerling rainbow trout, *Salmo gairdneri* Richardson, to various transport conditions, anaesthesia, and cold shock. *Journal of Fish Biology*, 20, 39-51.
- Carmichael, G.J., 1984. Long distance truck transport of intensively reared largemouth bass. *The Progressive Fish-Culturist*, 46, 111-115.
- Congleton, J.L., LaVoie, W.J., Schreck, C.B. & Davis, L.E., 2000. Stress indices in migrating juvenile chinook salmon and steelhead of wild and hatchery origin before and after barge transportation. *Transactions of the American Fisheries Society*, 129, 946-961.
- Davis, K.B. & Parker, N.C., 1986. Plasma corticosteroid stress response of fourteen species of warmwater fish to transportation. *Transactions of the American Fisheries Society*, 115, 495-499.
- Erikson, U. *et al.* 1997. See Chapter 7 *Cited References*.
- Garcia, L.M.B. *et al.* 2000. See Chapter 7 *Cited References*.
- Iversen, M. *et al.* 1998. See Chapter 7 *Cited References*.
- Johnson, S.L., Solazzi, M.F. & Nickelson, T.E., 1990. Effects on survival and homing of trucking hatchery yearling coho salmon to release sites. *North American Journal of Fisheries Management*, 10, 427-433.
- Korovin, V.A., Zybin, A.S. & Legomin, V.B., 1982. Response of juvenile fishes to stress factors associated with transfers during fish farming. *Journal of Ichthyology*, 22, 98-102.
- Paterson, B.D. *et al.* 2003. See Chapter 7 *Cited References*.
- Robertson, L. *et al.* 1988. See Chapter 7 *Cited References*.
- Rouger, Y. *et al.* 1998. See Chapter 7 *Cited References*.
- Sandodden, R. *et al.* 2001. See Chapter 7 *Cited References*.
- Schreck, C.B. *et al.* 1995. See Chapter 7 *Cited References*.
- Schreck, C.B., Solazzi, M.F., Johnson, S.L. & Nickelson, T.E., 1989. Transportation stress affects performance of coho salmon, *Oncorhynchus kisutch*. *Aquaculture*, 82, 15-20.
- Specker, J.L. & Schreck, C.B., 1980. Stress responses to transportation and fitness for marine survival in coho salmon (*Oncorhynchus kisutch*) smolts. *Canadian Journal of Fisheries and Aquatic Sciences*, 37, 765-769.
- Wedemeyer, G.A., 1992. Transporting and handling smolts. *World Aquaculture*, 23, 47-50.

5.7 Social interaction/Behaviour

- Abbott, J.C. & Dill, L.M., 1985. Patterns of aggressive attack in juvenile steelhead trout (*Salmo gairdneri*). *Canadian Journal of Fisheries and Aquatic Sciences*, 42, 1702-1706.
- Alanärä, A. 1997. Feeding behaviour and growth performance in fish within a dominance hierarchy. In: *Voluntary Food Intake in Fish*, Eds D. Houlihan, A. Kiessling & T. Boujard, pp 10.
- Alanärä, A. & Brännäs, E., 1996. See Chapter 7 *Cited References*.
- Bardach, J., Magnuson, R., May, R. & Reihart, J., (Eds.). 1980. *Fish behaviour and its use in the capture and culture of fishes*. Proceedings of the Conference on the Physiological and Behavioral Manipulation of Food Fish as Production and Management Tools, Ballagio, Italy, Nov 3, 1977, 512 p.
- Bell, A., 2002. The effects of some management practices on the behaviour and welfare of farmed Atlantic salmon in sea cages. PhD thesis, University of Stirling.
- Chandroo, K.P., Moccia, R.D. & McKinley, R.S., 2000. Utilization of physiological telemetry to monitor behavioural responses of rainbow trout, *Oncorhynchus mykiss* (Walbaum), to captive culture conditions. *Bulletin of the Aquaculture Association of Canada*, 99, 34-36.
- Cooke, S.J., Chandroo, K.P., Beddow, T.A., Moccia, R.D. & McKinley, R.S., 2000. Swimming activity and energetic expenditure of captive rainbow trout, *Oncorhynchus mykiss* (Walbaum), estimated by electromyogram telemetry. *Aquaculture Research*, 31, 495-505.
- Cordero Martinez, F.J., Beveridge, M.C.M. & Muir, J.F., 1994. A note on the behaviour of adult Atlantic halibut, *Hippoglossus hippoglossus* (L.), in cages. *Aquaculture and Fisheries Management*, 25, 475-481.
- Cubitt, K.F., 2002. Individual growth patterns in farmed Atlantic salmon: the role of social interactions. PhD Theses, University of Glasgow.
- Ejike, C. & Schreck, C.B., 1980. See Chapter 7 *Cited References*.
- Furevik, D.M., Bjordal, Å., Huse, I. & Ferno, A., 1993. Surface activity of Atlantic salmon (*Salmo salar* L.) in net pens. *Aquaculture*, 110, 119-128.
- Greaves, K., 2001. See Chapter 7 *Cited References*.
- Greaves, K. & Tuene, S., 2001. See Chapter 7 *Cited References*.
- Hulscher-Emeis, T.M., 1992. The variable colour patterns of *Tilapia zillii* (Cichlidae): integrating ethology, chromatophore regulation and the physiology of stress. *Netherlands Journal Zoology*, 42, 525-560.
- Laidley, C.W. & Leatherland, J.F., 1988. See Chapter 7 *Cited References*.
- Landless, P. J., 1976. Demand-feeding behaviour of rainbow trout. *Aquaculture*, 7, 11-25.
- Lochmann, S.E., Perschbacher P.W., Merry G.E. & Fijan N., 1998. Aggressive biting among channel catfish in pool studies. *The Progressive Fish-Culturist* 60, 119-126.
- Metcalfe, N.B., Huntingford, F.A. & Thorpe, J.E., 1987. The influence of predation risk on the feeding motivation and foraging strategy of juvenile Atlantic salmon. *Animal Behaviour*, 35, 901-911.
- Noakes, D.L.G. & Leatherland, J.F., 1977. Social dominance and interrenal cell activity in rainbow trout, *Salmo gairdneri* (Pisces, Salmonidae). *Environmental Biology of Fishes*, 2, 131-136.
- Olla, B.L. & Davis, M.W., 1989. The role of learning and stress in predator avoidance hatchery-reared coho salmon (*Oncorhynchus kisutch*) juveniles. *Aquaculture*, 76, 209-214.
- Øverli, Ø. *et al.* 1999. See Chapter 7 *Cited References*.
- Peters, G. *et al.* 1988. See Chapter 7 *Cited References*.
- Pottinger, T.G. & Pickering, A.D., 1992. See Chapter 7 *Cited References*.
- Sloman, K.A. *et al.* 2000. See Chapter 7 *Cited References*.
- Srivastava, R.K., Brown, J.A. & Allen, J., 1991. The influence of wave frequency and wave height on the behaviour of rainbow trout (*Oncorhynchus mykiss*) in cages. *Aquaculture*, 97, 143-153.
- Sutterlin, A.M. & Stevens, E.D., 1992. Thermal behaviour of rainbow trout and arctic char in cages moored in stratified water. *Aquaculture*, 102, 65-75.
- Wilson, J.L. & Roys, L.L., 1993. Behavioral interactions in juvenile channel catfish, *Ictalurus punctatus*. *Journal of Applied Aquaculture*, 3, 363-381.

5.8 Food

- Andrew, J.E., Noble, C., Kadri, S., Jewell, H. & Huntingford, F.A., 2002. The effects of demand feeding on swimming speed and feeding responses in Atlantic salmon (*Salmo salar* L.), gilthead sea bream (*Sparus auratus* L.) and European seabass (*Dicentrarchus labrax* L.) in sea cages. *Aquaculture research*, 33, 501-507.
- Barton, B.A., Schreck, C.B. & Fowler, L.G., 1988. Fasting and diet content affecting stress-induced changes in plasma glucose and cortisol in juvenile chinook salmon. *The Progressive Fish-Culturist*, 50, 16-22.
- Sakakura, Y., Koshio, S., Iida, Y., Tsukamoto, K., Kida, T. & Blom, J., 1998. Dietary vitamin C

- improves the quality of yellowtail (*Seriola quinqueradiata*) seedlings. *Aquaculture*, 161, 427-436.
- Simmons, L., Moccia, R.D., Bureau, D.P., Sivak, J.G. & Herbert, K., 1999. Dietary methionine requirement of juvenile arctic charr *Salvelinus alpinus* (L.). *Aquaculture Nutrition* 5.
- Talbot, C., Corneillie, S. & Korsøen, Ø., 1999. Pattern of feed intake in four species of fish under commercial farming conditions: implications for feeding management. *Aquaculture Research*, 30, 509-518.
- ### 5.9 Light regime
- Barahona-Fernandes, M.H., 1979. See Chapter 7 Cited References.
- Bolla, S. & Holmefjord, I., 1988. See Chapter 7 Cited References.
- Downing, G. & Litvak, M.K., 1999. See Chapter 7 Cited References.
- Duray, M. & Kohno, H., 1988. See Chapter 7 Cited References.
- Fernö, A., Huse, I., Juell, J.E. & Bjordal, A. 1995. Vertical distribution of Atlantic salmon (*Salmo salar* L.) in net pens: Trade-off between surface light avoidance and food attraction. *Aquaculture*, 132, 285-296.
- Marliave J. B. 1977. Effects of three artificial lighting regimes on survival of laboratory-reared larvae of the sailfin sculpin. *The Progressive Fish-Culturist*, 39, 117-118.
- McHugh, J.J., 1978. Effects of light shock and handling shock on striped bass fry. *The Progressive Fish-Culturist*, 40, 82.
- Mork, O.I. & Gulbrandsen, J., 1994. Vertical activity of four salmonid species in response to changes between darkness and two intensities of light. *Aquaculture*, 127, 317-328.
- Puvanendran, V. & Brown, J.A., 2002. Foraging, growth and survival of Atlantic cod larvae in different light intensities and photoperiods. *Aquaculture*, 214, 131-151.
- Tandler, A. & Helps, S., 1985. See Chapter 7 Cited References.
- ### 5.10 Health
- Anderson, D.P. 1990. Immunological indicators: effects of environmental stress on immune protection and disease outbreaks. In: *Biological indicators of Stress in Fish*, Adams, S.M. (Ed.) American Fisheries Society Symposium 8, Bethesda, Maryland, p. 38-50.
- Ashton, N. *et al.* 1969. See Chapter 7 Cited References.
- Austin, B. & Austin, D.A., 1987. *Bacterial fish pathogens: disease in farmed and wild fish*. Chichester: Ed. Ellis Horwood Ltd., 364 p.
- Bjerkås, E. & Bjørnstad, E., 1999. See Chapter 7 Cited References.
- Bjerkås, E. *et al.* 2001. See Chapter 7 Cited References.
- Bjerkås, E. *et al.* 2000. See Chapter 7 Cited References.
- Bruno, D.W. & Raynard, R.S., 1994. The effect of water temperature on eye opacity in Atlantic salmon (*Salmo salar* L.). *Bulletin of the European Association of Fish Pathologist*, 14, 86-88.
- Cullen, A.P. *et al.* 1994. Chapter 8 Cited References.
- Fraser, P.J. *et al.* 1990. See Chapter 7 Cited References.
- Garcia-Garbi, N. *et al.* 1998. See Chapter 7 Cited References.
- Griffin, B.R., Davis, K. B. & Schlenk, D., 1999. Effect of simulated copper sulphate therapy on stress indicators in channel catfish. *Journal of Aquatic Animal Health* 11, 231-236.
- Griffin, B.R., Davis, K.B., Darwish, A. & Straus, D.L., 2002. Effect of exposure to potassium permanganate on stress indicators in channel catfish. *Journal of the World Aquaculture Society*, 33, 1-9.
- Hargis, W.J., 1991. See Chapter 7 Cited References.
- Iwata, M., Komatsu, S., Collie, N.L., Nishioka, R.S. & Bern, H.A., 1987. Ocular cataract and seawater adaptation in salmonids. *Aquaculture*, 66, 315-327.
- Kincaid, H.L., 1989. See Chapter 7 Cited References.
- Mazur, C.F. & Iwama, G.K., 1993. See Chapter 7 Cited References.
- Pankhurst, N.W. & Dedual, M., 1994. See Chapter 7 Cited References.
- Peters, G., 1988. Stress macht auch Fische krank. *Naturwissensch. Rundschau* 41, 303-309.
- Peters, G., M. Faisal, Lang, T. & Ahmed, I., 1988. Stress caused by social interaction and its effect on susceptibility to *Aeromonas hydrophila* infection in rainbow trout *Salmo gairdneri*. *Diseases of Aquatic Organisms*, 4, 83-89.
- Peters, G., Nüssgen, A. & Raabe, A., 1991. Social stress induces structural and functional alterations of phagocytes in rainbow trout (*Onchorhynchus mykiss*). *Fish and Shellfish Immunology*, 1, 17-31.
- Plumb, J.A. 1994. See Chapter 7 Cited References.
- Reichenbach-Klinke, H.H., 1980. *Krankheiten und Schädigungen der Fische*. Stuttgart: Gustav Fisher Verlag. 472 p.
- Schreck, C.B., Maule, A.G. & Kaattari, S.L., 1993. Stress and disease resistance. In: *Recent Advances in Aquaculture* IV. Muir, J.F. and R.J. Roberts (Eds.), Blackwell Scientific Publications, London, p.170-175.
- Snieszko, S.F., 1974. See Chapter 7 Cited References.

- Stangeland K. *et al.* 1996. See Chapter 7 *Cited References*.
- Thorburn, M.A., Teare, G.F., Martin S.W. & Moccia, R.D., 2001. Group-level factors associated with chemotherapeutic treatment regimens in land-based trout farms in Ontario, Canada. *Preventive Veterinary Medicine*, 50, 165-176.
- Vazzana, M. *et al.* 2002. See Chapter 7 *Cited References*.
- Waagbø, R. *et al.* 1996. See Chapter 7 *Cited References*.
- Waagbø, R. *et al.* 1998. See Chapter 7 *Cited References*.
- Wall, A. & Bjerkas, E., 1999. A simplified method of scoring cataracts in fish. *Bulletin of the European Association of Fish Pathologist*, 19, 162-165.
- Wall, A.E. & Richards, R.H., 2000. Occurrence of cataracts in triploid Atlantic salmon (*Salmo salar*) on four farms in Scotland. *Veterinary Record*, 131, 553-557.
- Wedemeyer, G.A., 1997. Effects of rearing conditions on the health and physiological quality of fish in intensive culture. In: *Fish Stress and Health in Aquaculture* (ed. by G.K. Iwama, A.D. Pickering, J.P. Sumpter & C.B. Schreck), pp. 35-71. Society for Experimental Biology; seminar series 62. Cambridge University Press, Cambridge.
- Winfree, R.A. *et al.* 1998. See Chapter 7 *Cited References*.
- Yildiz, H.Y. & Pulatsu, S., 1999. Evaluation of the secondary stress response in healthy Nile tilapia (*Oreochromis niloticus* L.) after treatment with a mixture of formalin, malachite green and methylene blue. *Aquaculture Research*, 30, 379-383.
- 5.11 Other**
- Balarin, J.D., 1984. Tilapia. In: *Evolution of Domesticated Animals*, I.L. Mason (Ed.). Longman, London/Condon Group Ltd., New York, p. 391-397.
- Bhujel R.C., 2000. A review of strategies for the management of Nile tilapia (*Oreochromis niloticus*) broodfish in seed production systems, especially hapa-based systems. *Aquaculture*, 181, 37-59.
- Bullock, A.M., 1988. Solar ultraviolet radiation: A potential environmental hazard in the cultivation of farmed finfish. In: *Recent Advances in Aquaculture*, Muir, J.F. and R.J. Roberts (Eds.), Croom Helm Publishers, London, p. 139-224.
- Burgov, L., 1992. Rainbow trout culture in submersible cages near offshore oil platforms. *Aquaculture*, 100, 167-174.
- Casebolt, D.B., Speare, D.J. & Horney, B.S., 1998. Care and use of fish as laboratory animals: current state of knowledge. *Laboratory Animal Science*, 48, 124-136.
- CCAS (Canadian Council on Animal Care), 1994. *Resource* Vol. 18, No. 2, pp. 1-2.
- Davis, K.B. & Parker, N.C., 1990. See Chapter 7 *Cited References*.
- European Commission, 2001. *Facts and figures on the CFB. Basic data on the Common Fisheries Policy*. Luxembourg: Office for Official Publications of the European Communities, 28 pp.
- Grizzle, J.M., Chen, J., Williams, J.C. & Spano, J.S., 1992. Skin injuries and serum enzyme activities of channel catfish (*Ictalurus punctatus*) harvested by fish pumps. *Aquaculture*, 107, 333-346.
- Israeli-Weinstein, D. & Kimmel, E., 1998. See Chapter 7 *Cited References*.
- Kertz, A.F., 1996. Animal care and use: an issue now and in the future. *Journal of Animal Science*, 74 257-261.
- Kindischi, G. & Kirkpatrick, B., 1992. A small electric grid for preventing escapism of fish. *Journal of Applied Aquaculture*, 1, 103-109.
- Klontz, G.W., 1995. Care of fish in biological research. *Journal of Animal Science*, 73, 3485-3492.
- Monaghan, P. & Wood-Grush, D., 1990. Managing the behaviour of animals. Chapter 5. Chapman and Hall, London.
- Morton, A. 1994. Fish farming: The final frontier. *Can-Ag-Fax* 1: 1-3.
- National Research Council, 1991. *Education and training in the care and use of laboratory animals: A guide for developing institutional programs*. National Academy Press, Washington, D.C. p. 139.
- Nickum, J.G. 1988. Guidelines for use of fishes in field research. *Fisheries*, 13, 16-22.
- Olfert, E.D., Cross, B.M. & McWilliam, A.A., 1993a. *Guide to the care and use of experimental animals*, Vol. 1. Canadian Council on Animal Care, Ottawa, pp. 51-74 and 115-124.
- Olfert, E.D., Cross, B.M. & McWilliam, A.A., 1993b. *Guide to the care and use of experimental animals*, Vol. 2. Canadian Council on Animal Care, Ottawa, p. 1-10.
- Pickering, A.D., 1992. Rainbow trout husbandry: management of the stress response. *Aquaculture*, 100, 125-139.
- Scapigliati, G. *et al.* 1999. See Chapter 7 *Cited References*.
- Wagner, E.J. *et al.* 1997. See Chapter 7 *Cited References*.

This document presents the results of the EIFAC Ad Hoc Working Party on Handling of Fishes in Fisheries and Aquaculture, including the report of its workshop held from 24 to 26 March 2004 in Utrecht, Netherlands. The welfare of fishes has become a subject of attention in recent years. Concerns have been raised about the potential for infringed fish welfare in fisheries, aquaculture and fisheries research. The Ad Hoc Working Party agreed that handling of fishes must be carried out in a responsible way. Fishes show reactions to injurious stimuli. These stimuli result in physiological stress responses. Depending on the duration and intensity of the exposure to the stressor, the stress response varies from acute to chronic. Handling causing injuries and damage to the fishes should be avoided in fisheries, aquaculture and fisheries research. To ensure the welfare of fish in commercial fisheries, recreational fisheries and aquaculture, sufficient legislation and regulation is necessary. All EIFAC member countries discussed in this report have legislation and regulations concerning animal welfare. However, some of the legislation and regulations do not provide specific attention to fish welfare.

ISBN 978-92-5-106016-2 ISSN 0258-6096



9 789251 060162

TR/M/I0218E/1/06.08